3 1761 06704360 4

F.

MEDICAL LABORATORY METHODS AND TESTS

BY

HERBERT FRENCH

M.A., M.D. (Oxon.), F.R.C.P. (Lond.)

ASSISTANT PHYSICIAN, GUY'S HOSPITAL, ETC.

THIRD EDITION

168921
25/1/22

LONDON
BAILLIÈRE, TINDALL AND COX
8, HENRIETTA STREET, COVENT GARDEN

1913

(Reprinted)

[ALL RIGHTS RESERVED]

Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

PREFACE TO THE THIRD EDITION

THE fact that this small handbook has reached its third edition is a gratifying proof that it is of service. New methods and tests have been introduced, and old ones modified, but the primary object of the work has been maintained ; it confines itself to chemical and microscopical methods that are of established value in clinical medicine.

HERBERT FRENCH.

62, WIMPOLE STREET, LONDON, W.,

December, 1911.

PREFACE TO THE FIRST EDITION

THIS volume has been written in response to repeated complaints that there was no *small* book dealing with the chemical and microscopical tests and investigations which are most useful to medical men. The object has been to detail the commoner methods, pointing out the conclusions which may be drawn from the various tests, but laying stress upon the fallacies to which each is liable. The book deals, not with the examination of patients, but with that of fluids or substances obtained from them, and bedside methods have been excluded. It is intended to be a small handbook for the medical laboratory.

HERBERT FRENCH.

GUY'S HOSPITAL,

March 19, 1904.

CONTENTS

CHAPTER	PAGE
I. EXAMINATION OF THE URINE - - -	I
II. EXAMINATION OF THE BLOOD - - -	65
III. EXAMINATION OF THE SPUTUM - - -	107
IV. EXAMINATION OF PUS - - -	117
V. EXAMINATION OF THE GASTRIC CONTENTS - -	128
VI. EXAMINATION OF THE FÆCES - - -	140
VII. MICROSCOPICAL EXAMINATION IN AFFECTIONS OF THE SKIN - - - - -	154
VIII. EXAMINATION OF SEROUS EXUDATIONS, CEREBRO- SPINAL AND CYSTIC FLUIDS - - -	158
IX. PRESERVATION AND STAINING OF MORBID ANATOMY SPECIMENS - - - - -	165
X. EXAMINATION OF SEMINAL STAINS - - -	172
XI. TESTS FOR THE COMMONER POISONS - -	174
INDEX - - - - -	180

LIST OF ILLUSTRATIONS

FIG.		PAGE
1.	'Envelope' Crystals of Calcium Oxalate	7
2.	Diagram of a Calcium Oxalate Crystal	7
3.	Uric Acid Crystals	8
4.	Ammonio-Magnesium Phosphate Crystals	9
5.	Stellar Phosphate Crystals	10
6.	Magnesium Phosphate Crystals	10
7.	'Thorn-apple' Crystals	10
8.	Tyrosin Crystals	11
9.	Leucin Spherules	11
10.	Cystin Crystals	12
11.	Squamous Cells	14
12.	Bladder Cells	14
13.	Renal Tubule Cells	15
14.	Tailed Cells from Renal Pelvis	16
15.	Pus Cells, Fresh and Degenerate	17
16.	Renal Tube-Casts	18
17.	Spermatozoa	20
18.	Bilharzia Hæmatobia Ovum	24
19.	Spectra of Blood in Urine	31
20.	Phenyl-glucosazone Crystals	38
21.	Phenyl-lactosazone Crystals	38
22.	Diagram of Urobilin Spectrum	46
23.	Diagram of Hæmatoporphyrin Spectra	52
24.	Esbach's Albuminometer	53
25.	Pavy's Estimation of Sugar	55
26.	Gans' Apparatus	56
27.	Southall's Ureometer and Pipette	58
28.	Blood-counting Chamber	66
29.	Hæmocytometer Squares and Blood Corpuscles	66

FIG.		PAGE
30.	Hæmocytometer Pipettes - - -	67
31.	Counting Red Corpuscles - - -	69
32.	Haldane's Hæmoglobinometer - - -	74
33.	" " " introducing Coal-gas -	75
34.	Bottle for Staining Blood Films in - - -	78
35.	Cigarette-paper Method of making Blood Film -	79
36.	Normocytes - - - <i>Plate I., facing</i>	88
37.	Badly Fixed and Crenated Red Cells " "	88
38.	Megalocytes and Microcytes - " "	88
39.	Poikilocytes - - - " "	88
40.	Nucleated Red Corpuscles - " "	88
41.	Punctate Basophilia - - - " "	88
42.	Small Lymphocyte - - - " "	88
43.	Large (Hyaline) Lymphocyte - " "	88
44.	Polymorphonuclear Leucocyte - " "	88
45.	Coarsely Granular Eosinophile Leucocyte " "	88
46.	Myelocyte - - - " "	88
47.	Eosinophile Myelocyte - - - " "	88
48.	Basophile Leucocyte - - - " "	88
49.	Malaria Parasite : Ring Form - " "	88
50.	Filaria Embryo and Red Cells - " "	88
51.	Malaria Parasite : Crescent Form " "	88
52.	Spirilla of Relapsing Fever and Red Cells " "	88
53.	Widal Tube - - - - -	100
54.	Bell's Calcimeter - - - - -	105
55.	Elastic Fibres from Sputum - - - - -	109
56.	Tetrads and Pus Cells - - - - -	114
57.	Charcot-Leyden Crystals and Pus Cells - - -	116
58.	Curschmann's Spiral - - - - -	116
59.	Tubercle Bacilli and Pus Cells - <i>Plate II., facing</i>	120
60.	Staphylococci - - - " "	120
61.	Streptococci - - - " "	120
62.	Gonococci - - - " "	120
63.	Pneumococci - - - " "	120
64.	Diphtheria Bacilli - - - " "	120
65.	Anthrax Bacilli - - - " "	120
66.	Tetanus Bacilli - - - " "	120
67.	Actinomyces - - - " "	120
68.	Cholera Vibrio - - - " "	120

FIG.				PAGE
69.	Plague Bacilli	-	-	120
70.	Vincent's Angina Organisms	-	<i>Plate II., facing</i>	120
71.	Sarcinæ ventriculi	-	,,	138
72.	Yeast Cells	-	-	138
73.	Cholesterin Crystals	-	-	144
74.	Amœbæ of Dysentery	-	-	145
75.	Head of Tænia Solium	-	-	147
76.	,, ,, Mediocanellata	-	-	147
77.	,, Bothriocephalus Latus	-	-	147
78.	Oxyuris Vermicularis	-	-	148
79.	Ankylostomum Duodenale	-	-	149
80.	Trichocephalus Dispar	-	-	150
81.	Ovum of Trichocephalus Dispar	-	-	152
82.	,, ,, Tænia Solium	-	-	152
83.	,, ,, Ascaris Lumbricoides	-	-	152
84.	Spores on Hair in Tinea Tonsurans	-	-	155
85.	,, ,, ,, Circinata	-	-	155
86.	Sarcoptes Hominis (Female)	-	-	157
87.	Hydatid Hooklets	-	-	162
88.	Apparatus for Marsh's Test	-	-	175

CHAPTER I

EXAMINATION OF THE URINE

I. GENERAL CHARACTERISTICS OF URINE.

THE **amount** passed in twenty-four hours is, roughly, 1,500 c.c., or 50 ounces ; but varies within wide limits. Free perspiration diminishes it, as in summer and after exercise ; it is increased in winter and after copious drinking. As passed, it is usually quite clear. On standing, a faint cloud of 'mucus' often forms, especially in women. A dense pink precipitate of urates is common in summer urine ; a white precipitate of earthy phosphates is not unusual in the 'alkaline tide' which follows a full meal. Some common pathological changes in amount are :

Increase in—

Granular kidney,	18 to 100
ounces,	
Diabetes mellitus,	100 to 600
ounces,	
Diabetes insipidus,	100 to 800
ounces,	
Hysteria,	} Up to 100
After epileptic fit,	
After head injury,	
Recovery from	
anasarca,	} ounces or more.

Decrease in—

Fevers, such as pneumonia—
<i>e.g.</i> , 25 ounces.
Acute nephritis — <i>e.g.</i> , 10
ounces, or even none at all.
Heart failure— <i>e.g.</i> , 20 ounces,
10 ounces, or less.
After catheterization and other
local operations it may be
entirely suppressed.

The **specific gravity** is determined by the urinometer, the scale being read at the point where the graduated stem leaves the fluid. If there be insufficient urine to float the urinometer, add an equal bulk of distilled water ; multiply the last two figures of the specific gravity of the mixture by 2 to give that of the original urine. In health it varies from 1002 in very cold weather, or after much drinking, or in those who are very nervous, up to 1030 after profuse perspiration, whilst in some women it is often as high as 1040 without any evidence of existent or subsequent disease.

There are various contrivances, consisting, as a rule, of small glass bulbs of different colours and of different densities, by means of which the specific gravity of very small amounts of urine can be taken when the determination is very important, as in the case of examination for life insurance.

It is sometimes useful to know that if one multiplies the last two figures of the specific gravity by 2·2, one has a rough measure of the number of grammes of solids there are in each litre of urine. Thus if the specific gravity be 1018, then the specimen contains 39·6 grammes of solid per 1000 c.c. ; if the total urine passed in twenty-four hours be also known, it is easy to calculate the total quantity of solids that are being excreted in it.

Pathological Changes in Specific Gravity :

Unduly low in—

Diabetes insipidus, 1002 or 1004.
Hysteria, 1004.
Post-epileptic, 1004.
Granular kidney, 1008 to 1012.

Unduly high in—

Heart failure, 1030 to 1035.
Acute fevers, 1030 to 1035.
Diabetes mellitus, 1035 to 1045.

The **reaction** is determined with litmus-paper, 'acid' urine turning it red, 'alkaline' blue; when both red and blue litmus-papers are turned purple the reaction is termed 'amphoteric.' The acidity is due to acid sodium phosphate; an alkaline tide follows the big meal of the day, owing to the increased acid secretion in the stomach. The urine is more acid in those who eat meat; in vegetarians it may be continuously alkaline. On standing, the acidity at first increases, then diminishes, and is changed to alkalinity by conversion of urea into ammonia by the fermentative action of the *Micrococcus ureæ*.

Pathological Changes in Reaction:

Increased Acidity.

In acute fevers—*e.g.*, pneumonia, acute rheumatism.
In the uric acid diathesis.

Alkalinity.

Suppuration in the bladder.

The **colour** of normal urine, due to urochrome, varies from pale straw-yellow (low specific gravity) to dark sherry (high specific gravity).

Pathological Changes in Colour (for tests, see pp. 29, 42, 49, 51, 52):

Pallor: in diabetes mellitus, diabetes insipidus, granular kidney, hysteria, epilepsy, head injury, excessive drinking, extreme nervousness.

Blood: may make the urine bright red, dark red, port-wine colour; brown, brown-black, or black.

Scarlet: after eosin; sometimes taken unknowingly in red sweets.

Red-Brown: after rhubarb, senna, chrysophanic acid, hæmatoxylin.

Greenish-yellow to greenish-black : bile pigments ; phenyl compounds, such as carbolic acid, salol, resorcin, naphthol, guaiacol, creosote, naphthalene, indican, arbutin, uva ursi, bromoform, thallin, cascara sagrada, buckthorn, melanin, alkapton.

Green : after methyl green or other similar aniline dye sometimes used for colouring sweets or foodstuffs ; or after methylene blue when the latter is present in quite small amounts, or is in conjunction with bile or pus in the urine.

Bright yellow : after santonin.

Dichroic : urobilin, red by transmitted, green by reflected, light.

Blue : after the administration of methylene blue.

The **smell** of normal urine is 'aromatic,' but becomes ammoniacal on standing.

Pathological Changes in Smell :

Ammoniacal : in cystitis.

Fæcal : from contamination with *Bacillus coli communis*.

Like violets : after the administration of turpentine.

Sweet : in diabetes mellitus.

After eating asparagus the urine has a very characteristic odour, due to methyl mercaptan.

II. URINE DEPOSITS.

It is often possible to obtain sufficient deposit for examination by merely allowing the urine to stand in a conical specimen-glass for some hours, and then

pipetting off that which has subsided to the bottom. It is better, however, to use a centrifugalizing machine to concentrate the lighter particles—blood-corpuscles, bacteria, and so forth.

The **colour** of the deposit may help one to form some idea of its main constituent—for example :

A *white deposit* may be due to phosphates, oxalates, pus, mucus; in some cases to starch grains which have fallen into the urine after the use of a dusting-powder; or, rarely, to urates which have come down in a urine in which there was no uroerythrin to colour them pink.

A *pink deposit* is nearly always due to urates, which, though colourless themselves, usually carry down with them the urinary pigment uroerythrin.

A *dingy brown deposit* is very suggestive of phosphates that have carried down with them blood pigment, and therefore points to hæmaturia.

A *brick-red deposit* like cayenne pepper is due to uric acid.

A *dark crimson deposit* is due to blood.

There may be many different constituents in the same deposit, however, and more than a naked-eye examination is required. Microscopical investigation is essential.

The Method of examining a Centrifugalized Deposit from a Urine microscopically. — One of the best ways of examining a urinary deposit microscopically is as follows: Prepare a stock bottle of normal saline solution (0·6 per cent. of common salt in water), to which enough methylene blue has been added to make the colour deep; have cover-

glasses, pipettes, and slides all ready cleaned, and some 2-inch squares of blotting-paper; pour off all the supernatant fluid from the tube containing the centrifugalized urine, leaving behind the deposit and as little urine as possible; transfer a single drop of the deposit to a microscope slide by means of a pipette; with a clean glass rod transfer 1 drop of the normal saline methylene blue solution to the same slide, putting it near, but not in, the drop of centrifugalized deposit. The glass rod can be kept in the stock bottle of normal saline methylene blue for the purpose, and it is most important not to get it contaminated by anything on the slide, or else tube-casts and so on may be transferred back to the stock bottle. Now take a cover-glass, and with its edge gently, but thoroughly, mix together the drop of urine deposit with that of methylene blue; then put on the cover glass; remove the surplus fluid by laying a square of blotting-paper firmly over the top of it. The preparation is then ready for examination under both the low power and the $\frac{1}{6}$ -inch objective. The blue stains the cellular elements present, bringing out nuclei, and at the same time forming a good background in which renal tube-casts, red blood-discs, and crystals can be seen. It is most important when examining with the microscope to shut down the diaphragm as much as possible; many things will then be visible which would escape detection if the full light were employed.

Although many different constituents will be found in the same deposit, it is more convenient to describe each separately; and for this purpose it is of service to divide them into three groups, namely:

- A. Crystalline deposits.
- B. Amorphous non-organized deposits.
- C. Organized deposits.

A. CRYSTALLINE DEPOSITS.

1. **Calcium Oxalate** forms colourless crystals really octahedra (Fig. 2), but looking like squares with a diagonal cross—so-called ‘envelope’ crystals (Fig. 1). They vary greatly in size, some being only just visible under the $\frac{1}{6}$ -inch objective, others easily seen with the low power. To the naked eye the deposit is white, sometimes presenting a ‘powdered-

FIG. 1.—‘Envelope’ Crystals of Calcium Oxalate.

(High power of microscope.)

FIG. 2.—Diagram to show Octahedral Character of a Calcium Oxalate Crystal.

wig’ appearance from the deposition of a dense snow-white layer of oxalate crystals on the top of more flocculent mucus; or they may produce characteristic fine striæ, like scratch lines along the sides of a conical specimen-glass as the result of slight unevennesses in the surface of the vesse becoming filled with the minute crystals.

Importance. — They occur in many healthy acid urines, and are particularly abundant after eating rhubarb, tomatoes, cauliflower, and some other vegetables. There are therefore by no means always of pathological significance.

When so persistent as to constitute ‘oxaluria,’ they are a sign of acid dyspepsia.

The presence of many calcium oxalate crystals leads to irritation of the urinary passages, with the result that the urine often contains excess of nucleoproteid at the same time (see p. 48), and, in the male, even considerable numbers of spermatozoa (see p. 20). The irritation caused by oxaluria may be an important factor in some cases of frequency of micturition and of nocturnal enuresis in girls or boys, or of spermatorrhœa in men.

When the other clinical signs suggest the presence of a renal or vesical stone, the occurrence of abundant envelope crystals, together with red corpuscles, affords confirmatory evidence in favour of a 'mulberry' calculus, especially if the crystals tend to be agglomerated together into microscopic masses instead of all being separated from one another.

2. **Uric Acid** crystals are coloured yellowish-brown by the urochrome, and have a variety of shapes, the commoner being 'whetstones' and 'rosettes' (Fig. 3). 'Dumb bells' and 'bundles' are rarer forms, due to incomplete crystallization. The crystals occur in acid urine as a deposit which looks to the naked eye like grains of cayenne pepper.

FIG. 3.—Uric Acid Crystals.
(Whetstone and Rosette Forms.)
(High power of microscope.)

Importance.—They indicate hyperacidity of the urine. The condition is not necessarily gouty, nor is the uric

acid necessarily in excess. The process by which the uric acid becomes precipitated is, briefly, as follows: The NaH_2PO_4 , to which the acidity of the urine is due, tends to remove Na from combination with uric acid, forming Na_2HPO_4 , and converting the soluble sodium quadriurate first into the less soluble sodium biurate, then removing the sodium even from the latter, so that the comparatively insoluble uric acid goes out of solution. It is when the NaH_2PO_4 is in considerable excess—for example, upon a diet rich in meat—that the uric acid is precipitated to any extent. The hyperacidity may cause irritability of the bladder or urethritis, and the crystals may form a calculus. Treatment should be directed to diminishing the acidity of the urine in order to keep the uric acid in solution.

3. **Ammonio-Magnesium Phosphate** (NH_4MgPO_4) gives rise to colourless ‘coffin-lid’ or ‘knife-rester’ crystals (Fig. 4).*

Importance. — They occur in urines which contain ammonia. To the naked eye the deposit is heavy, opaque white, and may be mistaken for pus. The ammoniacal change may have occurred after the urine was passed; the crystals then have no pathological import. If found in fresh urine, they indicate cystitis or other cause of decomposition within the urinary tract.

FIG. 4. — Ammonio-Magnesium Phosphate Crystals.
(High power of microscope).

* The term ‘triple’ is often used for this variety of phosphate.

4. Earthy Phosphates :

(a) **Calcium Phosphate.**—This is usually amorphous; rarely it assumes the form of colourless elongated prisms, occurring singly, or in starlike clusters of 'stellar phosphate' (Fig. 5).

(b) **Magnesium Phosphate.**—This is also amorphous as a rule, but occasionally it occurs in the form of large colourless prisms, not very unlike those of ammonio-magnesium phosphate. The points of distinction will be seen on comparing Figs. 4 and 6.

FIG. 5.—Stellar Phosphate Crystals.

(High power of microscope.)

Importance.—'Earthy,' as distinct from alkaline or

ammonio-magnesian, phosphates do not indicate disease.

They occur in faintly acid, neutral, or alkaline urines. They are abundant in the urines of vegetarians, or in the alkaline urine which is passed soon after a very big meal. After a big dinner, for example, the first urine voided

FIG. 6.—Magnesium Phosphate Crystals.

(High power of microscope.)

may be milky-looking, owing to precipitation of earthy phosphates.

FIG. 7.—'Thorn-apple' Crystals.

(High power of microscope.)

5. Sodium Urate and Ammonium Urate.—Imperfectly formed 'thorn-apple' crystals, seen under the high power as small spheres with numerous short spines (Fig. 7).

Importance.—Both forms are distinctly

rare. Sodium urate crystals have been noted in the urine of newly born infants. 'Thorn-apple' crystals in adults are due to ammonium urate, in alkaline urines, in association with ammonio-magnesium phosphate. They indicate decomposition, either after the urine has been passed, or, in cases of cystitis, within the urinary tract.

6. Tyrosin.—Sheaves of colourless needles (Fig. 8), to be distinguished from phenyl-glucosazone crystals by (1) their smaller size, (2) their being colourless.

Importance.—They are rare; are usually associated with leucin; and are most typically seen in cases of acute yellow atrophy of the liver. They may be found in other affections of the liver also—*e.g.*, in phosphorus poisoning and occasionally in pernicious anæmia.

FIG. 8.—Tyrosin Crystals.
(High power of microscope.)

7. Leucin.—Pale yellow spherical masses, often concentrically striated (Fig. 9).

FIG. 9.—Leucin Spherules.
(High power of microscope.)

Importance.—The same as that of tyrosin. Leucin and tyrosin are amido-compounds which, in health, should be converted into urea by the liver. Starch grains may be mistaken for leucin (see p. 14).

8. Cystin.—Colourless hexagonal plates in acid urine (Fig. 10).

Importance.—Cystin is a rare constituent of urine ; but, if present in one member of a family, it is liable to occur in others also owing to familial peculiarity in metabolism. The condition is rare. Cystin has given rise to calculi which have a peculiar greenish or yellowish hue and soapy feel.

FIG. 10.—Cystin Crystals.

(High power of microscope.)

B. AMORPHOUS NON-ORGANIZED DEPOSITS.

1. **Urates.**—Sodium and potassium urates are often found precipitated in concentrated acid urines, which, though clear when passed at body temperature, have become clouded on getting cold. The precipitation depends upon the inability of the urine to keep as many urates in solution when cold as when hot. They have great affinity for uroerythrin, which gives the deposit a characteristic pink colour. In the absence of uroerythrin urate deposits are white like phosphates. Under the microscope, granular particles are seen running together into mosslike clumps.

To confirm, warm the urine to body temperature ; urates redissolve. In case albumin be present, fill 3 inches of a test-tube with the urine and its deposit ; warm the upper 2 inches gently. Urates will clear up. Boil the top inch ; albumin, if present, will form a cloud, so that in the top inch there is a cloud of albumin ; in the middle inch clear urine in which urates have redissolved at body heat ; at the bottom the original deposit of urates.

Importance of Urates.—They indicate a concentrated

urine, but afford no clue to the cause of concentration. It may be physiological—for instance, in the summer-time, or after violent exercise or a Turkish bath, when the quantity of sweat lost by the skin causes the amount of urine to diminish; or it may be due to pathological conditions which diminish the total quantity of urine passed, such as acute fevers, heart failure, or excessive loss of fluid by vomiting or diarrhoea.

2. **Earthy Phosphates.**—*i.e.*, *Phosphates of Calcium and Magnesium.*—To the naked eye the deposit is white and more or less dense and opaque. It may be confused with thick mucus, pus, alkaline phosphates, or colourless urates. When blood is present, the methæmoglobin tinges the phosphates a dirty brown. The urine may be alkaline, amphoteric, or just acid. Under the high power granular particles not unlike those of amorphous urates are seen, with an occasional crystal of stellar (calcium) (Fig. 5, p. 10) or magnesium phosphate (Fig. 6, p. 10).

To some of the deposit in a test-tube add acetic acid; phosphates will redissolve; pus or urates will remain unaltered; mucus will become more marked. It is essential that acetic, and not nitric, acid should be used in performing the test; for nitric acid, though it would redissolve the phosphates, would at the same time precipitate any albumin present.

Phosphates, being less soluble in hot than in cold urines, often form a cloud on boiling. This clears up at once on the addition of acetic acid.

Importance of Earthy Phosphate Deposit.—It has no pathological significance; it indicates that the urine is not very acid.

3. **Starch Grains.**—These appear under the microscope as round or oval bodies, with crescentic markings. They may be detected by their blue colour on adding iodine.

Importance.—They are adventitious, from dusting-powders, but may be mistaken for leucin until the iodine test has been applied.

C. ORGANIZED SEDIMENTS.

1. Epithelium :

(a) **Squamous Cells.**—These are of large size, with a small central nucleus. They occur singly (Fig. 11), or in groups of two or three, or in sheets of twenty or more.

FIG. 11.—Squamous Cells.
(High power of microscope.)

Importance.—They are derived from the vagina. If very numerous, they indicate vaginitis and catarrh ; in smaller numbers they have no pathological significance.

(b) **Bladder Cells.**—Though variable in appearance, these are of two types. Those from the deep layer in the bladder are oval at one end, and at the other taper into a long or short tail (Fig. 12, B); the superficial cells are convex on their free surface, with concavities on their under side, into which the oval cells of the deep layer have fitted (Fig. 12, A).

FIG. 12.—Bladder Cells.
A, superficial ; B, deep.
(High power of microscope.)

Importance.—An occasional surface cell has no particular significance. If present in numbers, vesical catarrh is probable. When pyriform cells from the deeper layer are found, there is probably ulceration of the bladder.

(c) **Renal Cells.**—It is by no means always possible to say for certain from their microscopical characters what is the source of each and every epithelial cell in the urine. Degenerate leucocytes and degenerate renal cells look much alike. When the epithelium is comparatively fresh, however, two kinds of renal cells may be recognized with some certainty, namely—

(i.) Cells from the walls of the renal tubules.

(ii.) Cells from the renal pelvis.

(i.) **Cells from the Renal Tubules.**—These are about twice the size of leucocytes, and, when fresh, have a polygonal outline and central nucleus (Fig. 13, A). More frequently they have

FIG. 13.—Renal Tubule Cells.

A, fresh ; B, fatty.

(High power of microscope.)

undergone fatty degeneration, and appear irregular, and opaque with fine granules, amongst which the nucleus is no longer distinguishable (Fig. 13, B).

Importance.—They indicate a catarrhal condition of the kidney. They are particularly numerous in acute Bright's disease, but occur also in consecutive nephritis ; renal congestion from heart failure ; after infarction ; with renal calculus, tuberculous kidney, and with growth.

(ii.) **Cells from the Renal Pelvis.**—Cells from the renal pelvis have various shapes and sizes. Some are

FIG. 14. — Tailed Cells from Renal Pelvis.

• High power of microscope.)

ovoid and of comparatively large size; some are polygonal, resembling those from the renal tubules. But the most characteristic renal pelvic cell is long and narrow, with a single nucleus and a slender, wavy tail (Fig. 14). It might at first sight be mistaken for a pyriform cell from the deeper layers of the bladder mucosa, but the tail is much longer and more tapering.

Importance.—The presence of these tailed cells indicates excoriation of the renal pelvis, such as is caused by pyelitis, calculus, or a tuberculous lesion. The cells are always pathological.

(d) **Fragments of New Growth.**—It happens occasionally that fragments detached from a new growth in the kidney or bladder are found in the centrifugalized deposit. The cells stain well with methylene blue, and form continuous masses or clumps, sometimes with papillomatous projections. In the case of a bladder growth the particles are often large enough to be visible to the naked eye in the fresh urine; fragments of renal growth are usually much smaller. The variety of cell seen will depend upon whether the grow is a papilloma, a sarcoma, or a carcinoma. One word of warning is needed: it sometimes happens that leucocytes or other cells clump together in the centrifugalized deposit in such a way as to resemble fragments of tissue. It is only by familiarity with the microscopical characters of urinary

deposits that such artefacts can be distinguished from new growth.

Importance. — The finding of a fragment of new growth clinches the diagnosis. It must be remembered, however, that such fragments are not being broken off from the main growth constantly, so that in a suspicious case repeated examinations should be made if no fragment of new growth is found at first.

2. **Leucocytes.**—If fresh, these appear exactly like those of the blood, and are mostly polymorphonucleated: if the urine has stood, they become granular and fatty, but are distinguished from fatty renal cells by their smaller size (Fig. 15). Acetic acid renders the nucleus more distinct.

FIG. 15.—Pus Cells, Fresh and Degenerate.
(High power of microscope.)

Importance. — Small numbers indicate irritation within the urinary tract. They occur in Bright's disease, urethritis, vesical catarrh, and in association with blood in the urine. Large numbers indicate purulent inflammation, such as gonorrhœal urethritis, gleet, prostatitis, cystitis, renal growth, tubercle or calculus, or suppurative nephritis. Coli bacilluria causes slight albuminuria, together with considerable excess of leucocytes, barely amounting to macroscopic pus; the urine is generally opalescent, and the diagnosis, suggested by the pus cells, is confirmed by finding *Bacillus coli communis* in pure culture in a catheter specimen.

3. **Red Blood-Corpuscles.**—These often look like

those of fresh blood; in very dilute urines they may be laked, and appear as 'ghosts'; in concentrated urines they may be crenated, and variously crumpled.

Importance.—In a woman they may be derived from menstrual blood. Otherwise, their presence is the best test of hæmaturia. They may be derived from the kidney, the ureter, the bladder, the prostate, the urethra, a caruncle; the microscope affords no evidence of their source. Apart from menstruation, they are always pathological.

FIG. 16.—Renal Tube-Casts.
(High power of microscope.)

4. **Renal Tube-Casts.**—There are the following varieties (Fig. 16):

(1) Epithelial casts, in which outlines of renal cells are seen.

(2) Blood casts, composed of red corpuscles.

(3) Leucocyte casts, consisting of white corpuscles, which are distinguished from renal cells by their smaller size.

(4) Fatty casts, not very different to granular casts, but containing here and there small, highly refractile specks and globules of fat.

(5) Granular casts, having no cell elements, and a typical granular appearance.

(6) Hyaline and waxy casts, seen only in outline, the cast itself having no internal structure.

Many become broken and unrecognizable; some appear of different structure in different parts—*e.g.*, granular at one end, epithelial at the other.

Importance.—It would seem probable that, whereas blood and leucocyte casts depend on exudation from vessels, epithelial, fatty, and granular casts are progressive downward changes in casts derived from the renal tubule cells. The precise nature of the hyaline material of a hyaline cast is not known. It is the result of inflammatory changes, and forms the matrix in which are embedded the epithelial cells, fat globules, or granular débris of the other forms of casts. A simple hyaline cast is one consisting of matrix only, with nothing embedded in it. An occasional granular or hyaline cast is to be found in healthy urine that has been much centrifugalized. Small numbers may be found when the kidney is passively congested, from failure of the right side of the heart. Blood casts may occur in blood diseases, and with other causes of hæmorrhage within the kidney substance, such as infarction or venous thrombosis; but the presence of many casts points strongly to nephritis. No case of Bright's disease has been fully diagnosed until renal tube-casts have been found in the urine. Epithelial, blood, and leucocyte casts indicate acute renal inflammation; as the latter subsides, blood and leucocyte

casts disappear, epithelial casts grow fewer, and the fatty casts increase; in chronic nephritis hardly any epithelial casts are seen, fatty casts are few, granular and hyaline preponderate. Waxy casts are not essentially different to hyaline: they are more refractile and more easily seen, but have no significance that hyaline casts have not.

5. Spermatozoa.—These are recognized readily in urine. Under a high power each is seen to have an

ovoid head about two-thirds the size of a red corpuscle, and a threadlike tail, usually broken off short, but, if complete, about as long as eight red corpuscles (Fig. 17).

Importance.—Spermatozoa may be found in the urines of perfectly healthy adult males, but they should not be present continuously and in numbers.

FIG. 17.—Sperma-
tozoa.
High power of micro-
scope.)

In themselves they are not pathological, but they may point to the existence of some irritating constituent in the urine. The commonest such irritant is oxalate of lime. If many envelope crystals are present, it is not at all uncommon, in the male, to find at the same time excess of mucus, excess of nucleo-proteid (see p. 48), and many spermatozoa. Treatment of the oxaluria, by diet or otherwise, readily cures the condition.

6. Bacteria.—(a) *Non-Pathogenic Micro-organisms.*—Normal urine which is left exposed to the air in warm weather soon swarms with the *Micrococcus ureæ*, which converts urea into ammonium carbonate. Associated with it may be hosts of non-pathogenic bacilli.

Importance.—None, except that if not recognized they may be mistaken for organisms which cause disease. If invasion by adventitious organisms is to be inhibited, some antiseptic such as boric acid or thymol should be added to the specimen, the amount so added varying with the volume of urine which it is desired to keep sweet. It is noteworthy that some urines decompose far less readily than do others. Those from patients with uncomplicated phthisis may resist decomposition for days.

(b) *Pathogenic Micro-organisms.* — The pathogenic organisms which may be found are—

Staphylococci.

Streptococci.

Gonococci.

Tubercle bacilli.

Bacillus coli communis.

Typhoid bacilli.

In examining for any of these, a catheter specimen should be obtained, and a film of the centrifugalized deposit made (p. 117). To prevent this from washing off in the process of staining, it is well to albuminize the slides or cover-slips as follows: Mix some fresh white of egg with nine times its bulk of water and filter it; smear the filtered egg-white upon one surface of the cover-slip, and allow it to dry unexposed to dust. Slips albuminized in this way can be kept for future use. To find out which surface is albuminized, breathe on it; the unalbuminized surface becomes opaque from condensed moisture, the other remains transparent. Smear the urine deposit thickly on this surface; allow it to dry; fix the film by passing it through a Bunsen flame five times.

For *gonococcus*, *Bacillus coli communis*, *typhoid bacillus*, *streptococcus*, and *staphylococcus*, stain with carbon methylene blue or carbol fuchsin (p. 117). Staphylococci

and streptococci may also be stained by Gram's method (p. 118), the others not.

For *tubercle bacilli* stain by the Ziehl-Neelsen method (p. 111).

Importance. — All these organisms are pathogenic, and any of them may cause cystitis. Gonococci more often come from the urethra. Typhoid bacilli are sometimes excreted in what appears to be healthy urine long after convalescence from enteric fever, but cannot be distinguished by the microscope from *Bacillus coli communis*. The latter may be present in pure culture in cases of coli bacilluria, the symptoms of which may vary from those of nocturnal enuresis to those which simulate appendicitis or other acute abdominal condition. If thorough identification of the various bacteria is required, extensive cultural methods are essential. These are beyond the scope of a student's or practitioner's clinical laboratory. Tubercle bacilli, if present, indicate a tuberculous lesion of the kidney, ureter, or bladder. If absent, a tuberculous lesion is not excluded. With care, tubercle bacilli are found at the first examination in something like half the cases of tuberculosis of the kidney or bladder. In suspicious cases in which the first examination proves negative, the procedure should be repeated upon successive days. There is nearly always a good deal of pus in the urine also, but it is worthy of note that the bacilli have many times been found when there was no pus. Absence of pus, therefore, is not an absolute contra-indication to examination for tubercle bacilli.

A source of fallacy is the *smegma bacillus*, which also

stains by the Ziehl-Neelsen method. It will not retain the stain, however, if the film be first immersed for ten minutes in warm caustic soda solution containing 5 per cent. of alcohol, whereas tubercle bacilli will still stain as before. Those who are familiar with urinary examinations can distinguish smegma bacilli from tubercle bacilli morphologically, the former being longer and better defined than the latter. The term 'smegma' might suggest that the smegma bacillus was confined to the male. This is by no means the case. It occurs not infrequently in the normal urine of the female as well as in that of the male, and may be present in a catheter specimen. It is, therefore, a very real source of fallacy in examinations for tubercle bacilli in urine. Another method of avoiding the fallacy is to immerse the films, just before counterstaining in carbol methylene blue, in a solution of 95 per cent. spirit containing 5 per cent. of strong hydrochloric acid. If left in this for five minutes before counterstaining, the red colour will be discharged from the smegma, and not from the tubercle bacilli.

7. **Yeast Fungi** (Fig. 72, p. 138) have no pathological significance. In warm weather they commonly occur, particularly in diabetic urine which has stood open to the air.

8. **Fat Globules** may occur as highly refractile particles, soluble in ether, readily stained black by osmic acid, or red by Sudan III. (p. 99).

Importance.—In temperate climates it is very rare to find fat in the urine unless by accidental or intentional admixture of milk or oil—*e.g.*, after vomiting,

or the passage of an oiled catheter.[^] True chyluria results most commonly from infection with filaria in the tropics (p. 97); it is also found occasionally in cases of persistent subacute nephritis; and, along with chylous ascites, after rupture of, or obstruction to, the receptaculum chyli or thoracic duct.

9. **Bilharzia Hæmatobia** is the only parasite likely to affect the urine. The mature worm occupies the pelvic vessels. The eggs work their way through the walls of the latter, and reach the urine through the bladder wall, producing hæmaturia.

FIG. 18.—*Bilharzia Hæmatobia*
Ovum.

(Medium power of
microscope.)

Each ovum is $\frac{1}{200}$ inch long and half as broad. With the low power it resembles a melon seed, with a short spine at one end, or, in some cases, on one side (Fig. 18). The contained embryo will not hatch in urine, but if as much of the urine as possible is poured off from the centrifugalized deposit, some clean, tepid water poured on to the latter, and the

tube again centrifugalized, the embryos may be observed under the microscope bursting their chitinous shells and then swimming about free in the water by means of the innumerable cilia which cover them.

Importance.—The parasite is acquired in some way from water, chiefly in Egypt and South Africa. The diagnosis of the cause of the hæmaturia depends on finding the ova.

10. **Prostatic Threads.**—If the urine be poured into a glass cylinder, these may be seen by the naked eye floating about as fine threads; being composed of mucus, they appear structureless under the low power.

Importance.—They point to previous gonorrhœa.

11. **Mucus.**—This is a normal constituent of many urines, particularly in women. Under the microscope it appears as structureless wisps and bands, in which may be embedded crystals, leucocytes, and so forth. The wisps stain deeply with methylene blue.

Importance.—In most cases mucus has no pathological significance. If abundant it suggests irritation of the urinary passages, the nature of the irritant being indicated by the calcium oxalate crystals, pus cells, or other features of the urine that may be noted at the same time. It is easy, before experience is acquired, to mistake mucus for renal tube-casts, and this in two ways: First, parts of the mucus often have a cylindrical form—‘mucous cylindroids’; secondly, if in pressing down the cover-slip on to the slide the former slips slightly along the latter, the mucus becomes pressed out lengthways in straight castlike streaks; the latter will at once be detected for what they are, because they will be all lying with their long axes parallel to one another.

III. TESTS FOR SUBSTANCES IN SOLUTION.

1. **Albumin.**—Filter the urine if it be turbid; it is essential for the following test that it should be clear.

Heller’s Nitric Acid Test.—Pour a little non-fuming nitric acid into a test-tube; hold the latter slanting, and allow urine to flow gently from a pipette on to the surface of the acid. Watch the line of

junction ; an opaque white ring will appear at once if much albumin be present, more slowly with less. A brownish-red ring which often appears also is independent of the presence of albumin, and results from the oxidation of indican by the nitric acid.

Fallacies.—(1) If fuming nitric acid be used, the urea will be decomposed, giving off bubbles of CO_2 and N_2 . This will cause the urine to mix with the acid, so that the line of junction is destroyed, and even though albumin be present it may not be possible to distinguish the white ring.

(2) Albumose will give a similar white ring ; but it disappears on warming, to reappear on cooling, whereas that due to albumin remains on warming.

(3) If the urine be concentrated, urea nitrate may form a white ring. Dilute the urine with an equal bulk of water, and repeat the test. Urea nitrate will now remain in solution.

(4) After the administration of drugs containing resins—*e.g.*, copaiba—a white ring appears with nitric acid ; it is more diffuse than that of albumin. It is due to precipitation of the resin, which will redissolve on adding a little alcohol.

Test by Heating.—Fill a test-tube three parts full of urine; note the reaction to litmus-paper. If alkaline, just acidify with acetic acid ; the object of this is to prevent the formation of uncoagulable alkali-albumin on warming. Acetic acid is used because a stronger, such as nitric or hydrochloric, would form acid-albumin, which is likewise uncoagulable by heat. Boil the upper 1 inch thoroughly, and look through the test-tube at a dark background. If

the boiled inch is as clear as the unboiled urine, no albumin is present ; if it has become cloudy, the precipitate may be due to one or more of four things—namely :

- (i.) Albumin.
- (ii.) Phosphates.
- (iii.) Carbonates.
- (iv.) Nucleo-proteid.

To decide between these, first add 2 or 3 drops more acetic acid ; if the cloud clears up entirely, it is due to phosphates ; if, whilst clearing up, it produces brisk effervescence, it was due to carbonates, either alone or together with phosphates ; if it remains, albumin or nucleo-proteid, or both, are present. Now add 1 drop of strong nitric acid : this will clear up the cloud if it is due to nucleo-proteid, and if the cloud persists, albumin is present.

Fallacies.—(1) Phosphates (see above).

(2) Carbonates (see above).

(3) Nucleo-proteid (see above).

(4) If the urine be not acid, alkali-albumin may be formed, and no cloud appear, though albumin be present.

(5) When the amount of albumin present is small, it may be missed if the boiling be not continued for half a minute or more.

(6) In very rare cases—chiefly myeloid sarcoma—Bence-Jones's albumose gives a dense cloud with the boiling test, insoluble in acetic acid. The nature of the case is suggested, however, by the low temperature at which the coagulum appears (see p. 47).

The heat test is the best for albumin, and has the fewest fallacies.

The Salicyl-Sulphonic Acid Test.—It is essential to have a particularly small test-tube. Into this put 30 drops of the urine, and add a few drops of a saturated aqueous solution of salicyl-sulphonic acid. If no precipitate occurs, no proteid is present. If there is a precipitate, boil the tube; if the liquid in it becomes clear again, the precipitate was due to proteoses and not albumin; if the precipitate persists, it is due to albumin; indeed, with albumin it increases by coagulation on heating. The main advantage of the test is that it saves time when many examinations have to be made, as in hospital work; for if no precipitation occurs with the reagent in the cold, no heating is required; the reagent is very stable, and no particular care is required as to the exact amount of it used for each test.

The Magnesium Sulphate and Nitric Acid Test.—This is performed in the same way as the ordinary nitric acid test, but using nitric acid saturated with magnesium sulphate. It has the advantage of precipitating the white ring of albumin more rapidly than nitric acid alone does; but it has the disadvantage of precipitating other things also. If the test gives no white ring, albumin is absent; if it does give a white ring, the boiling test should be used to decide whether it is due to albumin or not.

There are many other tests for albumin, such as those with potassium ferrocyanide, picric acid, and so forth, but the above are far more reliable and delicate than any of the others.

Importance of Albumin.—The presence of albumin in the urine is seldom without significance, though far

from always meaning Bright's disease. In women it may be an accidental accompaniment at the menstrual period, or arise from vaginitis or some uterine discharge; in men it may be temporary after seminal emission; a so-called 'functional' albuminuria is almost common in both sexes from puberty to thirty. With blood or pus albumin is always present, but pus accounts for far less albumin than might at first be expected. It is frequent with heart failure, and may result from any cause of low arterial blood-pressure. Abdominal tumours, ascites, and inferior vena cava thrombosis may cause albuminuria by obstructing the outflow from the renal veins. It is not uncommon in the severe anæmias, and accompanies lardaceous disease of the kidneys. Only when casts are present as well as albumin can Bright's disease be diagnosed with certainty. Microscopical examination of the urinary deposit should never be omitted in a case of albuminuria.

2. Blood.

The Guaiacum Test.—To 1 inch of urine in a test-tube add 1 drop of tincture of guaiacum; the resin forms a white precipitate. Pour on to the surface 1 inch of ozonic ether. If blood be present, it and the ozonic ether together oxidize the guaiacum, and a blue colour appears at the junction of the fluids.

Fallacies.—(1) Pus gives a similar colour, but it is more green than blue, and appears more slowly. Pus will give the colour with the guaiacum alone.

(2) Iodides in the urine give a similar blue colour, but it appears more slowly than with blood.

(3) Adventitious organic matter due to vomit or

sputum may also give the blue colour. The fallacy may be avoided by testing a catheter specimen.

(4) It is always well to be sure that the tap water does not itself give the blue colour. In places where the tap water is not to be used for drinking purposes, enough organic matter may be present in it to give a blue ring with the ozonic ether and tincture of guaiacum test.

Heller's Test.—Phosphates have an affinity for blood pigment, and carry it down with them. Render some urine in a test-tube strongly alkaline with caustic soda, and boil it; if blood be present, the deposit of earthy phosphates is brownish-red.

Fallacies.—(1) Earthy phosphates may be deficient in the urine, and no deposit result. To obviate this add 2 drops of calcium chloride solution.

(2) Certain drugs—rhubarb, senna, santonin—give a similar reaction.

(3) The test is seldom used clinically. Its main interest lies in the fact that when phosphates are spontaneously deposited in a urine containing blood, they carry the blood pigment with them, and appear dingy brown instead of white (p. 13).

The Spectroscope Test.—Even a small pocket spectroscope is expensive, so that the test is little employed. When the microscope reveals red corpuscles, it is not needed; but in cases of hæmoglobinuria it is valuable in confirming the presence of blood pigment. Filter the urine if it be not clear; dilute it with water if it absorb too much light. With the spectroscope look at a bright light through a test-tube filled with the urine. If recent blood be present, the

two absorption bands of oxyhæmoglobin will be noted (Fig. 19), one at the junction of the yellow with the green, the other in the middle of the green rays of the spectrum. Blood that has been longer mixed with the urine gives the three bands of methæmoglobin

FIG. 19.

(Fig. 19), one in the red in addition to the two of oxyhæmoglobin.

Fallacies.—None, except that the test requires care and skill in carrying it out. Other substances have spectroscopic bands, but none exactly like the above.

The Microscope Test.—Examine the centrifugalized deposit for red corpuscles; their presence is the best proof of hæmaturia.

Fallacies.—(1) In alkaline urine or in urine of very low specific gravity the red corpuscles may swell up and disappear.

(2) In paroxysmal hæmoglobinuria and in black-water fever blood pigment is present, but no red corpuscles.

Importance of Blood in the Urine.—Except in women at the menstrual period, the presence of blood is always pathological. Laboratory tests cannot by themselves decide what part of the genito-urinary

tract the blood comes from ; but when the blood is intimately mixed with the urine, the source is usually the kidney, whilst when it comes chiefly at the beginning or the end of micturition, the source is elsewhere. Methæmoglobin in fresh urine indicates renal hæmorrhage rather than vesical. Hæmaturia does not always indicate a lesion in the urinary tract, for it may occur in the blood diseases, such as leukæmia or purpura.

3. **Pus.**—The deposit is opaque and white ; in small quantities it may be mistaken for mucus ; in larger for phosphates or for colourless urates. Urates disappear on warming, pus remains ; phosphates increase with heat, but clear up with acetic acid. Mucus and pus can be distinguished by the following tests :

The Liquor Potassæ Test.—To 1 inch of the suspected deposit in a test-tube add a few drops of liquor potassæ ; pour the mixture from one test-tube into another. Pus will have partially dissolved and become ‘ropy’ and gelatinous.

Fallacy.—The test will not detect small quantities of pus.

The Ozonic Ether Test.—To 1 inch of the deposit in a test-tube add a few drops of ozonic ether. On gently shaking, numbers of small bubbles of liberated oxygen will be seen rising through the fluid.

Fallacies.—(1) Blood also causes bubbling with ozonic ether.

(2) Normal urine causes a few bubbles to come off, so that the test will not detect small quantities of pus.

The Guaiacum Test.—This is performed as for

blood. Even without adding ozonic ether, a green-blue colour appears after a few minutes.

Fallacies.—Blood, iodides, saliva (p. 29).

The Microscope Test.—Under $\frac{1}{8}$ -inch objective pus corpuscles appear as leucocytes which have undergone granular or fatty degeneration. Their nuclei can be brought out by adding a drop of acetic acid (Fig. 15, p. 17).

Fallacy.—If it be remembered that a few leucocytes may be found in normal urines, there is no fallacy. The presence of numerous leucocytes in the urine indicates pus. The microscope is the only satisfactory test for it.

Importance of Pus.—It is always pathological. Small quantities of pus derived from the urethra or vagina may occur in acid urine. Much pus in an acid urine indicates renal suppuration, in which case the acidity is maintained by healthy urine from the sound kidney. If both kidneys be affected, or if there be cystitis, the purulent urine will be alkaline and probably ammoniacal. The condition known as coli-bacilluria (a variety of pyelonephritis) often gives rise to no macroscopic precipitate of pus, and yet pus corpuscles may be found in abundance on microscopical examination of the centrifugalized deposit. The urine is usually a little clouded, and it contains a trace of albumin. Without microscopical examination the diagnosis may be missed altogether.

4. **Sugar.**—Two sugars may occur in urine, glucose and lactose.* The following tests are for glucose :

* It is for larger books than this to discuss pentoses and other rarities.

Fehling's Test.—Fehling's solution is made by mixing equal parts of the following fluids:

Copper Sulphate Solution.

Crystallized copper sulphate	..	34.64	grammes.
Distilled water up to	..	500	c.c.

Rochelle Salt Solution.

Potassium hydrate	..	125	grammes.
Sodium potassium tartrate (Rochelle salt)	..	173	..
Distilled water up to	..	500	c.c.

The copper sulphate solution is pale green, that of Rochelle salt colourless. On mixing the two the potassium hydrate converts the copper sulphate into cupric hydrate, and the latter is kept in solution by the Rochelle salt, giving a deep blue transparent fluid. The separate fluids keep good any length of time; mixed, they slowly decompose. They should therefore be kept in separate bottles and mixed as required. There are several modifications of Fehling's solution, one of the best of which is Haines's, the object of it being to obtain a single fluid which will keep good for years. The formula is as follows:

Haines's Fluid.

Copper sulphate	..	3	grammes
Caustic potash	..	9	„
Glycerin	..	100	„
Distilled water	..	to 600	c.c.

To perform the test, boil 1 inch of Fehling's solution in a test-tube; if good, it remains clear. In another test-tube boil 1 inch of urine. Pour the hot urine into the hot Fehling's solution. If glucose be present, it

will reduce the cupric hydrate to cuprous oxide, which comes down as an orange precipitate without further boiling.

Fallicies.—(1) Lactose reduces Fehling's solution, but may be distinguished by the fermentation and the phenyl-hydrazine tests (pp. 37, 38).

(2) Slight reduction of Fehling's solution, causing first a greenish colour of the solution, and then a dark greenish-yellow precipitate, may be caused by other reducing substances in the urine, namely:

Alkapton.	Hippuric acid.
Creatinin.	Hypoxanthin.
Glycosuric acid.	Uric acid.
Glycuronic acid.	Xanthin.

And the products of certain drugs :

Acetanilide.	Helmitol.
Antipyrin.	Lead.
Amyl nitrite.	Mercury.
Camphor.	Morphine.
Carbolic acid.	Opium.
Chloral.	Salicylates.
Chloroform.	Urotropine.
Glycerin.	Volatile oils.
Glycyrrhizin.	

None of these ferment. Error from these sources is not likely to occur if the two fluids be boiled separately and then mixed. Glucose and lactose alone will give the reduction without further boiling.

(3) Albumin interferes with Fehling's test. If it be present, add acetic acid, boil, filter off the coagulated albumin, and test the filtrate.

(4) In ammoniacal urines the free ammonia may prevent precipitation of cuprous oxide. Before test-

ing, with Fehling's solution, boil well to expel the ammonia.

(5) Phosphates may be present in sufficient amount to be precipitated by the alkali in the Fehling's solution, and the flocculent deposit may have a greenish tint which may be mistaken for a positive sugar reaction if confirmatory tests are not employed.

(6) Not only may creatinin itself reduce Fehling's solution on prolonged boiling, it may also inhibit reduction by small quantities of sugar if the boiling is carried out for a short time only. This is one of the commonest causes for an ambiguous reaction with Fehling's test. If a small amount of sugar is present with an abundance of creatinin, there may be no change of colour on ordinary boiling, and yet when the test-tube is examined again after standing a little while, a greenish-yellow precipitate may be found. On the other hand, if no more is present, but much creatinin, the Fehling's solution may exhibit a partial reduction on prolonged boiling. Before it is possible to say whether sugar is present or absent in either case, it is essential to apply either the phenyl-hydrazine or the fermentation test, or both.

Trommer's Test.—Trommer's solution is a mixture of copper sulphate and potassium hydrate. It differs from Fehling's in not containing Rochelle salt. The test is performed in precisely the same way and has the same fallacies.

Böttger's Test.—The reagent used is the following:

Caustic soda	10 grammes.
Sodium potassium tartrate (Rochelle salt) ..	4 ..
Bismuth subnitrate	2 ..
Distilled water	100 c.c

To 2 inches of urine in a test-tube add 5 drops of the reagent. Boil for five minutes. If glucose be present, the bismuth is reduced, and a fine black precipitate of bismuth suboxide appears.

Fallacies.—Lactose and glycuronic acid also give the test; but uric acid, creatinin, and xanthin bases, which give some reduction with Fehling's test, give none with Böttger's.

The Picric Acid Test.—To 1 inch of urine in a test-tube add $\frac{1}{4}$ inch of saturated solution of picric acid and 5 drops of caustic potash. Boil. If glucose be present, the solution darkens in colour, becoming first dark brown and then almost black from the reduction of picric to picramic acid.

Fallacies.—(1) Lactose gives the test also.

(2) In normal urine creatinin causes some darkening, so that as a sugar test picric acid is not very delicate.

(3) Impure picric acid darkens when boiled with caustic potash. The solution should therefore be tested before it is added to the urine.

Moore's Test.—Mix $\frac{1}{2}$ inch of urine with $\frac{1}{2}$ inch of liquor potassæ in a test-tube and boil. If sugar be present a ruddy brown colour develops, going on to dark brown and then almost black.

Fallacies.—(1) The test will not detect small quantities of sugar.

(2) Carboluria, alkapton, and indican also cause darkening with potash.

The Phenyl-hydrazine Test.—To 2 inches of urine in a test-tube add as much phenyl-hydrazine chloride as will lie easily on a threepenny-piece, and

a rather larger quantity of sodium acetate. Stand the test-tube in boiling water for twenty minutes, then

FIG. 20.—Phenyl-glucosazone Crystals.

allow it to cool. If glucose be present, phenyl-glucosazone crystals are formed; these may be pipetted off and examined under the low power; they are very characteristic fans and sheaves of long bright yellow needles (Fig. 20).

Fallacies.—(1) Lactose gives yellow needles of phenyl-lactosazone, but they are smaller, and arranged less as fans and sheaves than as spheres with radiating spines (Fig. 21).

(2) If the crystals are abundant, there is no doubt about the presence of glucose; if they are in small numbers only, however, or not quite typically formed, there may be a doubt as to whether the atypical crystals may not be due to glycuronic acid.

The Fermentation Test.—Triturate a piece of good yeast, the size of a small marble, with 250 c.c. of urine. Fill a Southall's ureometer (Fig. 27, p. 58) with the mixture, and allow it to stand in a warm place for twenty-four hours. Glucose, if present, will ferment, and the CO_2 given off will collect as free gas at the top of the closed limb of the ureometer.

FIG. 21.—Phenyl-lactosazone Crystals.

Fallacies.—None. No other reducing body in the urine will ferment with yeast.

SUMMARY OF SUGAR TESTS.

For Glucose.

For Lactose.

Fehling's test serves all ordinary purposes.

Gives Fehling's test.

Phenyl-hydrazine test is a good confirmation.

Gives crystals with the phenyl-hydrazine test different to those of glucose.

Fermentation test absolute proof of the presence of glucose.

Will not ferment with yeast.

Moore's, Trommer's, Böttger's, and the picric acid tests are confirmatory.

Importance of Sugar in the Urine.—It is important to know if the sugar be glucose. Lactose is not pathological, but is closely related to pregnancy in women, occurring both before and after parturition. Glucose in appreciable quantities is pathological. Its presence may be temporary, as after alcoholic excess, an injury to the head, or a vascular lesion of the brain. It may be intermittent in the gouty or alimentary glycosuria of elderly people, or it may indicate a more advanced stage of diabetes mellitus, in which case it is often associated with acetone and diacetic acid.

5. Acetone.

The Sodium Nitroprusside Test.—To 1 inch of urine in a test-tube add two or three drops of caustic soda and then a 1 per cent. solution of sodium nitroprusside drop by drop. A ruby red colour appears. Add acetic acid. If acetone be present the colour deepens to a rich port-wine colour.

Fallacies.—(1) If potash or ammonia be used instead of soda, the test may fail.

(2) Creatinin gives a similar ruby red with the nitroprusside and caustic soda, but on adding acetic acid the colour, instead of deepening, disappears.

Rothera's Modification of the Nitroprusside Test.—To 10 c.c. of urine in a test-tube add a few drops of a fresh saturated solution of sodium nitroprusside; render the mixture definitely alkaline with ammonia, and then saturate it with crystalline ammonium sulphate. A colour similar to that of a solution of potassium permanganate, and apparently distinctive of acetone, develops gradually, reaching a maximum in about fifteen minutes, and then gradually fading. Normal urine, although liable to contain traces of acetone, gives practically no colour with the test, which is, nevertheless, much more delicate than the other methods of using sodium nitroprusside.

The Iodoform Test.—To 1 inch of urine in a test-tube add 5 drops of caustic potash; warm to body heat; add iodine solution until the liquid is yellowish-brown; warm gently. If acetone be present iodoform will form, and it may be recognized by its odour or by its yellow crystals.

Fallacy.—It is seldom possible to be sure that iodoform has been formed unless the acetone has been obtained in concentrated form by distilling the urine, collecting the first few c.c. that come over, and then applying the test to the distillate. This is scarcely possible in ordinary practice, so that although with distillation the iodoform test is a good one, for ordinary clinical work the sodium nitroprusside test is most used.

Importance of Acetone in the Urine.—Apart from sugar,

acetone has been found in lobar pneumonia, tetany, gastric ulcer, starvation, and other conditions; associated with glycosuria, acetone indicates diabetes mellitus, with acidosis and consequent liability to coma. Diabetes without acetonuria is a much more favourable stage than diabetes with acetonuria.

6. Diacetic or Aceto-acetic Acid.—To 1 inch of urine in a test-tube add liquor ferri perchlor. (B.P.) drop by drop. A white precipitate of iron phosphate forms first; but almost immediately, if aceto-acetic acid be present, the liquid becomes deep purple-red, the colour being discharged again on warming.

Fallacies.—If carbolic acid, salol, salicin, salicylates, are being taken, the urine contains phenyl compounds which give a similar reaction with ferric chloride. The colour due to these causes does not disappear on warming.

Importance.—Diacetic acid has a significance similar to that of acetone. β -oxybutyric acid is a precursor of acetone and diacetic acid, all three being progressive stages in the end-products of a pathological metabolism which is seen most characteristically in the dangerous acidosis of diabetes. β -oxybutyric acid is nearly always present along with diacetic acid and acetone, but there is no easy test for it. For clinical purposes, therefore, it is sufficient to test for the latter.

7. Glycuronic Acid.—There is no good clinical test for this acid; its main importance is that it may be mistaken for sugar because it reduces Fehling's and Böttger's solutions. But *it will not ferment*. It occurs in normal urine to a small extent, and is

increased after giving chloral, butyl chloral, nitrobenzol, camphor, salicylates, and after chloroform narcosis.

8. Bile Pigments.—The colour varies from brownish-yellow to dark green, and may be simulated by carboluria, alkaptonuria, hæmaturia, indicanuria, melanuria, methæmoglobinuria, hæmoglobinuria, hæmatoporphyrinuria; but if the urine be shaken up the colour of the froth also is greenish when bile pigments are present—not so in the other cases.

Gmelin's Test.—Upon a white porcelain slab put 1 drop of the urine, and close to it a drop of fuming nitric acid. At their point of coalescence a play of colours—yellow, red, green, blue, yellow—will occur if bile pigments be present. The test may also be performed in a similar way on filter-paper; or, if 1 inch of nitric acid be put into a test-tube, and the urine carefully poured on to its surface, the play of colours is readily seen at the junction of the two fluids.

Fallacy.—None.

The Iodine Test.—Take 1 inch of urine in a test-tube; gently pour 10 per cent. solution of iodine in alcohol on to its surface. In the presence of bile pigments a bright green ring appears between the two liquids.

Fallacy.—The test is not so delicate as Gmelin's.

Importance of Bile Pigments in Urine.—They indicate jaundice, but not its cause.

9. Bile Salts.

Pettenkofer's Test.—When to a solution of bile salts in a test-tube 1 c.c. of strong solution of cane-sugar is added, and strong sulphuric acid drop by

drop, holding the end of the test-tube in running water to keep it cool, a rich crimson colour appears.

This test, however, is not applicable to fresh urine. Bile pigments may be abundant, and yet Pettenkofer's test may fail to show bile salts. It is seldom of any clinical importance to know whether bile salts as well as bile pigments are present; but if they are to be tested for, it is essential to extract them from the urine with chloroform, and then apply Pettenkofer's test to the chloroform extract.

The Sulphur Test.—Fill a test-tube to the top with urine; sprinkle flowers of sulphur on to its surface. Whereas in healthy cold urine the sulphur floats, even minute traces of bile salts so alter the surface tension that the sulphur sinks.

Fallacies.—(1) If the urine be warm, the sulphur sinks even in the absence of bile salts.

(2) The test is almost too delicate. By it all sorts of urines are shown to contain traces of bile salts, so that the very delicacy of the test detracts from its clinical value.

Importance of Bile Salts.—Bile salts are not known to give evidence which is not equally afforded by the bile pigments.

10. **Chlorides.**—To 1 inch of urine in a test-tube add 2 drops of nitric acid and 2 drops of silver nitrate solution. Chlorides give a dense white precipitate. The nitric acid is added first to keep the phosphates and sulphates in solution.

Fallacy.—Albumin and albumose, if present, are precipitated by the nitric acid.

Importance.—Some deficiency of chlorides occurs in

many fevers, but in lobar pneumonia they may be almost or quite absent, assisting in diagnosis from empyema or simple pleurisy. Presence of chlorides does not exclude lobar pneumonia, and they may be absent in other conditions; for example, suppurative peritonitis, or in absolute starvation. In chronic nephritis there is a great delay in the elimination of chlorides, so that examination for these may afford evidence as to the degree of renal insufficiency.

II. Sulphates.—To 1 inch of urine in a test-tube add 2 drops of strong hydrochloric acid to prevent the precipitation of barium phosphate, and then barium chloride solution drop by drop. A dense white precipitate of barium sulphate comes down.

Importance.—Sulphates in the urine vary directly with the intake of food, diminishing rapidly in starvation, or in diseases in which little or no food is assimilated. Besides this, there is a very important relationship between the inorganic and the organic sulphates; the latter are spoken of as 'ethereal,' and they do not react with barium chloride as above. It is difficult to estimate them accurately, but a broad idea of the amount of ethereal sulphate present may be obtained as follows: Precipitate all the inorganic sulphates with barium as above, filter, boil the filtrate with a few drops of hydrochloric acid; this breaks up the ethereal sulphates and a corresponding precipitate of barium sulphate comes down. The greater the degree of intestinal putrefaction, the larger the proportion of ethereal to total sulphates present in the urine. Some authorities lay much stress upon the ratio which the ethereal sulphates bear to the

total nitrogen (the 'co-efficient of Amann') in this connection, but the necessary analyses are too complex to warrant further discussion here.

12. Indican.—To 2 inches of urine in a test-tube add 10 drops of strong hydrochloric acid and 2 drops of fuming nitric acid; allow to cool; add $\frac{1}{2}$ inch of chloroform, and shake up thoroughly. If indican be present, the chloroform, when it has again sunk to the bottom of the test-tube, will be tinged either blue or red. The principle of this test is to oxidize the indican. Nitro-hydrochloric acid is a convenient agent for this purpose; freshly prepared **bleaching powder** is sometimes used instead; to 20 c.c. of urine add 2 c.c. of chloroform and 20 c.c. of hydrochloric acid; then at once freshly prepared bleaching powder solution (CaCl_2O_2) drop by drop, inverting the test-tube after each addition; it is most important not to add excess of bleaching powder, or else the blue colour of the indigo is discharged again for further oxidation. The chloroform is used merely to dissolve out the oxidized indican, in order that the colour may be more easily seen in it than it can be in the urine. Whether the result is blue or red seems to depend upon the temperature the solution reaches; the less it is heated the more likely is the colour to be blue.

Fallacies.—(1) Albumin interferes with the test. It should, if present, be removed by acidifying the urine with acetic acid, boiling, and filtering off the coagulated proteid.

(2) Iodides, if present, will colour the chloroform blue.

Importance.—Indican is a mixture of potassium

indoxyl and skatoxyl sulphates, derived from the indol and skatol which arise from intestinal decomposition. Indicanuria thus points to putrefaction in the intestine from chronic constipation, from acute or ulcerative colitis or enteritis, and other similar conditions.

13. Urobilin.—Examined with the spectroscope, an absorption band between the green and blue is typical of urobilin (Fig. 22). To 1 inch of urine in a

FIG. 22.—Diagram of Urobilin Spectrum.

test-tube add $\frac{1}{2}$ inch of strong ammonia, and then zinc chloride solution drop by drop; filter. If urobilin be present the filtrate will be fluorescent, and will show the spectroscopic band more markedly than before.

Importance.—In healthy urine the chief pigment is urochrome, not urobilin. The latter is a pathological pigment. It may occur in small amounts in many febrile diseases, but it is found most particularly when the liver is diseased, especially in cases of hepatic cirrhosis and in pernicious anæmia.

14. The Diazo-Reaction.—Ammonia and the following two solutions are needed; each should be fresh:

Sulphanilic Acid Solution.

Hydrochloric acid	..	..	..	5 c.c.
Distilled water	..	..	..	100 „
Sulphanilic acid to saturation.				

Sodium Nitrite Solution.

Sodium nitrite	..	..	..	0.5 gramme.
Distilled water	..	..	..	100 c.c.

To 1 inch of urine in a test-tube add 1 inch of sulphanilic acid solution and 5 drops of the sodium nitrite. Add ammonia till alkaline, and shake thoroughly. The test is positive if the liquid becomes port-wine colour *and the froth rose red*. The rose colour of the froth is essential.

Importance.—The reaction was formerly used in the diagnosis of typhoid fever, but it has gone out of fashion as being much less valuable than Widal's test (p. 100). A positive diazo-reaction in phthisis is a bad sign; but if the reaction has at one time been found, and later has disappeared, it indicates improvement in the patient.

15. Albumose.—First remove albumin by acidifying the urine with acetic acid, boiling and filtering. On to the surface of 1 inch of nitric acid in a test-tube gently pour some of the cooled filtrate. If albumose be present, a white ring will appear at the junction of the fluids. The ring will disappear on warming, to reappear again on cooling.

Importance.—Albumosuria occurs to a slight degree in many fevers, notably pneumonia, but it has no known significance. It also occurs when there are collections of pus under pressure, as in pelvic or appendicular abscess or empyema. In large quantities it appears to be pathognomonic of endosteal sarcoma of bone, in which case the variety of albumose gives special reactions, and is known as 'Bence Jones's body.' This generally attracts attention when the urine is boiled after acidification with acetic acid; at a temperature of about 60° C.—far lower than that at which albumin coagulates—a very dense cloud forms

which does not clear up on the addition of more acetic acid as phosphates would. To confirm the presence of albumose, the urine should be saturated with ammonium sulphate crystals, filtered to remove all the proteids so precipitated, and the filtrate tested for albumose with nitric acid as above. The biuret test may also be applied to the filtrate; which, boiled with a trace of copper sulphate, will give a pink colour reaction if albumose or peptone is present.

16. **Nucleo-proteid.**—The chief interest that nucleo-proteid has for the clinician is that it may, if not recognized, lead to an erroneous diagnosis of albuminuria when the heat test alone is used (p. 27). It is therefore of great importance in examination for life insurance. Almost all urines contain a trace of nucleo-proteid. Some contain much more than a trace, in which case further examination will show that there is something in the urine which irritates the urinary passages—calcium oxalate, for example (p. 8).

Nucleo-proteid is precipitated by acetic acid. If a trace of albumin is present, together with excess of nucleo-proteid, and it is desired to find out the relative proportions of each, a good plan is as follows: Into each of two test-tubes, A and B, pour an equal quantity of the urine, and strongly acidify each with the same number of drops of acetic acid. Leave A unboiled, but boil B. When B has cooled, compare the degree of opacity in the two. In A the precipitate is due to nucleo-proteid only, in B to nucleo-proteid together with the albumin coagulated by heat. If any albumin is present, therefore, the cloud in B must be more than that in A.

17. Carboluria, Alkaptonuria, and Melanuria.

—These three conditions account for various greenish-brown, greenish-black, brownish-black, and blackish tints that are sometimes observed in urines. They may be mistaken for bile or blood pigments. The urine is frequently normal in colour when passed, but darkens when exposed to the air. In carboluria this is due to oxidation of hydroquinone and pyrocatechin; in alkaptonuria to homogentisic acid; in melanuria to melanogen, which becomes melanin on standing. The darkening is in each case accelerated by adding caustic potash. The following reactions will indicate the probable nature of a given case:

In Carboluria.—Ferric chloride gives a white precipitate of phosphates and a *purple* solution. Bromine water gives a dense pale yellow precipitate of tribromophenol.

In Alkaptonuria.—Fehling's solution gives a reduction, but the urine does not ferment with yeast. Dilute solution of ferric chloride gives a *blue* colour, which rapidly disappears, returning with the addition of each drop of the reagent.

In Melanuria.—Ferric chloride: the precipitate of phosphates carries down the melanin as a dark grey deposit, soluble in excess of ferric chloride. Bromine water gives a dense yellow precipitate, which slowly blackens. There is no reduction with Fehling's solution. Caustic potash, sodium nitroprusside, and acetic acid used in the same way as in testing for acetone (p. 39) give a deep blue colour.

Importance.—Carboluria is by no means synonymous with carbolic acid poisoning. Some patients are so

susceptible that carbolic acid lotion or dressings, carbolic acid or salicylates by the mouth, may cause the condition without there being any other symptoms. Nevertheless, it is wise to desist from the use of the particular drug or application which causes carboloria in any given case.

Alkaptonuria is congenital and the result of certain peculiarities in proteid metabolism quite compatible with good health. Few individuals present the abnormality. It is detected first by the blackening of the infant's diapers.

Melanuria may occur in patients suffering from melanotic sarcoma, but it is not constant in such cases, and it is sometimes marked in other wasting diseases.

18. **Lead.**—Add 1 gramme of ammonium oxalate to 150 c.c. of urine that has been concentrated by boiling; immerse a strip of bright magnesium wire in it for twenty-four hours. Lead, if present, will form a deposit on the magnesium. Dissolve this in dilute nitric acid. Test the solution for lead as follows:

(1) A drop of HCl added gives a dense white precipitate of $PbCl_2$.

(2) Chromate of potash gives a bright yellow precipitate of lead chromate.

(3) Ammonium sulphide in excess gives a black precipitate of lead sulphide.

(4) Potassium iodide gives a bright yellow precipitate of lead iodide.

Importance.—Lead is excreted in the urine after the administration of potassium iodide in cases of plumbism, and sometimes an otherwise doubtful diagnosis

may be cleared up by discovering lead in the urine.

19. Some Drug Reactions in the Urine :

Antifebrin.—Extract the urine with chloroform ; to the extract add nitrate of mercury solution and warm. A green colour indicates that antifebrin is being taken.

Antipyrin.—The urine may be red, suggesting blood, but the guaiacum reaction is negative. With ferric chloride, a brownish-red colour forms.

Chloral and Chloroform cause increase of glycuronic acid. With Fehling's solution there will be some reduction, but there will be no fermentation with yeast.

Copaiba, Cubebs and Balsams.—Nitric acid gives a white ring, which might be mistaken for albumin. The ring is redissolved by alcohol. Add 2 drops of HCl to urine in a test-tube. Copaiba gives a pink colour.

Gallic Acid and Tannic Acid.—Ferric chloride gives a dark inky coloration, from the formation of gallate or tannate of iron.

Iodides.—The guaiacum test gives a blue colour, which may suggest blood. Add HCl and shake up with chloroform ; the latter takes up the liberated iodine, becoming violet ; indican gives a similar reaction, but requires HNO_3 as well as HCl.

Rhubarb.—The reddish-brown urine may suggest blood. No guaiacum reaction is given. With caustic potash the red colour deepens ; on now adding HCl it becomes yellow. The pigment is precipitated by adding solution of barium chloride.

Salicylates, Salol, Creosote, Arbutin, Uva Ursi and Carbolic Acid.—With Fehling's solution there is a partial reduction on continued boiling; ferric chloride gives a purple colour which, unlike that due to diacetic acid, does not disappear on warming.

Santonin.—Bright yellow urine. Caustic potash turns it bright pink, and barium salts do not precipitate this pigment.

Senna.—Similar to Rhubarb (see above). In each case the colour of the urine is due to chrysophanic acid.

Sulphonal.—Sometimes causes hæmatoporphyrinuria, the urine being port-wine coloured, suggesting blood, and yet not giving the guaiacum test. Add 2 drops of acetic acid to some urine, and shake it up with an equal volume of ether. This extracts the

FIG. 23.—Diagram of Hæmatoporphyrin Spectra.

pigment, which, examined spectroscopically, gives the four-banded spectrum of hæmatoporphyrin, or when HCl is added, a band in the orange and another between the yellow and green (Fig. 23).

Urotropin.—The addition of caustic soda and then of solution of phloroglucin gives a deep red colour. The urine also gives some reduction with Fehling's solution.

IV. METHODS OF ESTIMATING VARIOUS CONSTITUENTS OF URINE.

1. Estimation of Albumin.

Esbach's Method.—An Esbach's albuminometer (Fig. 24) and Esbach's fluid are needed. The former is a special test-tube, graduated in its lower part by marks which correspond to parts per thousand, and with two other marks, U and R respectively.

Esbach's fluid is as follows :

Picric acid	..	..	..	10 grammes
Citric acid	..	..	..	20 "
Distilled water to	..	..	..	1,000 c.c.

Fill the tube with urine to the mark U, and add reagent to the mark R ; cork, and gently invert several times. Stand vertically for twenty-four hours. The mark then corresponding to the upper level of the precipitate gives the parts per thousand of albumin.

Fallacies.—Occasionally the albumin floats instead of sinking. In that case shake the tube up again and leave for another twenty-four hours.

The test is one of sedimentation, so that if left for more or less than twenty-four hours the reading will be proportionately less or more than it should be.

The higher graduations become closer and closer together, so that the difference for each part per 1,000 over 5 is difficult to measure accurately. If more than 5 parts per 1,000 are present, dilute the urine once, twice, thrice, as

FIG. 24.—
Esbach's
Albumino-
meter.

may be necessary, and multiply the amount then found by 1, 2, or 3 respectively.

The picric acid also precipitates pigments and aromatic compounds. The amount of these is negligible in proportion to large amounts of albumin, but with a reading less than 1 part per 1,000 the error due to pigments is great. The readings are therefore accurate only between 1 per 1,000 and 5 per 1,000, and only after standing exactly twenty-four hours.

Robert's Method.—This depends on the fact that it requires three minutes for the white ring with nitric acid to appear when there is 1 part of albumin in 30,000; no special apparatus is required. Dilute the urine ten, twenty, thirty times, and so on, and after each dilution test with nitric acid. Suppose the white ring required three minutes to appear after thirty times dilution, there would be 30 parts of albumin per 30,000, or 1 part per 1,000.

2. Estimation of Sugar.

Pavy's Method.

Pavy's Solution.

Copper sulphate	4·16 grammes
Rochelle salt	20·4 „
Caustic potash	20·4 „
Strong ammonia	300 c.c.
Water to	1,000 „
10 c.c. are reduced by 0·005 gramme of glucose.	

Pavy's solution differs from Fehling's in two respects—viz., it is ten times as dilute, and it contains ammonia; the latter is to prevent the precipitation of cuprous oxide. The fresh solution is blue; when reduced, colourless. Fill a burette with the urine diluted 1 in 20. Put 10 c.c. Pavy's solution

and 20 c.c. water into a 200 c.c. flask fitted with a cork bored with two holes. Through one hole pass the nozzle of the burette, through the other a glass tube 4 inches long, which reaches just into the flask at one end, and is open to the air at the other (Fig. 25). Heat the flask until the Pavy's solution just boils: keep it simmering; turn the stopcock of the burette so that the diluted urine drops in steadily at the rate of 3 drops per second. Do not alter this rate of dropping, but watch the colour of the fluid in the flask; it will presently fade. When the Pavy's solution is just colourless, turn the stopcock off, and read off the amount of urine added. The calculation is as follows: The 10 c.c. of Pavy's solution require 0.005 gramme of glucose for complete reduction. Suppose 3 c.c. of the twenty times diluted urine were added;

FIG. 25.—Pavy's Estimation of Sugar.

∴ in 3 c.c. of twenty times diluted urine there is 0.005 gramme of glucose.

∴ in 1,000 c.c. of twenty times diluted urine there are $\frac{0.005 \times 1,000}{3}$ grammes of glucose.

∴ in 1,000 c.c. of original urine there are $\frac{0.005 \times 1,000 \times 20}{3}$ grammes of glucose.

= 33.3 per 1,000.

The method is liable to considerable personal error; gives strictly comparable results, however, if always

performed by the same person and in the same way.

If the flask be not corked as described, the ammonia rapidly boils off, and a thick yellow deposit of cuprous oxide comes down.

The urine should always be diluted to such an extent that not less than 1 c.c. and not more than 3 c.c. are required to decolorize the 10 c.c. of Pavy's solution. (N.B.—After the estimation, the blue colour will return to the reduced Pavy's solution, owing to reoxidation of the cuprous oxide in the flask.)

Gans' Method.—This is less accurate than Pavy's, but may be employed in general practice, as it is less

complicated. It depends on the volume of CO_2 evolved on fermentation with yeast. To 10 c.c. of urine add a small quantity of finely triturated yeast; dilute to 100 c.c. with water: mix thoroughly. Transfer 10 c.c. to the bulb of a Gans' apparatus (Fig. 26), and insert the stopper. The

FIG. 26.—Gans' Apparatus.

A, before fermentation;
B, after.

graduated vertical limb is open at the top, and there is an arrangement, easily understood when the apparatus is seen, for adjusting the level of the fluid to the zero mark. Stand in a warm place till next day. The CO_2 evolved in the bulb will have forced the fluid up in the vertical limb, and the graduation corresponding to its upper level gives the parts per cent. of sugar in the original urine.

The graduations on the tube having been made experimentally with known percentages of glucose, it does not matter that a certain proportion of the CO_2 escapes from the open end.

The method avoids error from presence of other reducing bodies than sugar in the urine; but it is open to much inaccuracy owing to the variations in volume of gas in the bulb with changes in temperature and barometric pressure.

Specific Gravity Method. — This is easy, and affords approximate results. It is very useful in general practice. Mix some finely triturated yeast with a quantity of urine. Take the specific gravity of the mixture. Stand in a warm place for twenty-four hours, and take the specific gravity again. Every degree of specific gravity lost corresponds to 1 grain per ounce of glucose. For example:

Specific gravity—before fermentation ..	..	1040
“ “ after “ ..	..	1022
Grains per ounce of glucose ..		<hr/> 18

3. Estimation of Urea.

Sodium Hypobromite Method with Southall's Ureometer.

Hypobromite Solution.

Caustic soda	100 grammes
Water	250 C.C.
When cool, add bromine	25 “

Sodium hypobromite (NaBrO) is formed, and caustic soda is in excess.

Fill the bulb of the ureometer (Fig. 27, A) with this solution, and tilt the apparatus so as to displace all

the air in the closed vertical limb. Measure off 1 c.c. of urine with the pipette (Fig. 27, B). A good way of doing this is to expel *all* the air from the elastic nipple, plunge the end of the glass pipette into the urine, and release the nipple so that urine is drawn up beyond the mark. Raise the pipette out of the urine, and by gentle pressure on the nipple expel the excess down to the mark 1; then plunge the tip into clean water and

FIG. 27.—Southall's Ureometer (A), and Pipette (B).

release the nipple again. It does not matter that the measured 1 c.c. of urine will thus be slightly diluted. Now insert the end of the pipette through the bulb of the ureometer till it reaches the bottom of the vertical limb: slowly compress the nipple, expelling its contents into the hypobromite solution. As soon as the urine mixes with the hypobromite, bubbles of gas come off, the urea being decomposed into CO_2 , H_2O ,

and N_2 . The CO_2 is dissolved by the excess of soda ; and H_2O condenses ; the N_2 rises to the top of the vertical limb. Allow the froth to settle down ; read off the graduation corresponding to the lower level of the gas. There are two sets of graduations, one showing parts per cent., the other grains per ounce.

Importance. — A healthy man excretes about 400 grains, or 30 grammes, of urea in twenty-four hours ; but there are wide limits, according to the food taken. In the majority of cases a small percentage of urea means either that little nitrogenous food is being taken, or that the urine is very dilute. This is apparently as true in Bright's disease as in other illnesses. In certain fevers more urea is excreted than corresponds to the nitrogen in the food, owing to breaking down of the patient's own tissues. This may also be the case in diabetes, though no deductions can be drawn unless the nitrogen in the food is also known. In surgery much stress is laid upon the amount of urea as an indication for or against operating in cases of tuberculous kidney, stone in the kidney, and so forth. When the total urea in twenty-four hours' urine is less than 100 grains, it is held to be inadvisable to operate. This is probably the case, but it must not be thought that it is an indication that the kidneys *cannot* excrete more urea than this. It is more likely that the patients, being as ill as they are, do not give their kidneys more urea than this to excrete.

The hypobromite method is only approximate ; more nitrogen is given off than corresponds to the urea alone, less than to the total nitrogen in the urine.

For exact estimations of the urea, the complicated

Mörner-Sjöqvist process should be used (see Schäfer's 'Text-Book of Physiology,' vol. i.).

4. Estimation of Uric Acid.

Hopkins's Potassium Permanganate Method.

—To 100 c.c. of urine in a flask add 35 grammes of ammonium chloride. Allow to stand for fifteen minutes. Insoluble ammonium urate is precipitated. Filter. Wash the precipitate with hot ammonium chloride solution. Open out the filter-paper, and with hot distilled water wash the precipitate into a 100 c.c. flask. Dilute accurately to 100 c.c. with water. Pour into a beaker, add 20 c.c. strong sulphuric acid, and from a burette drop in $\frac{N}{20}$ permanganate solution made up as follows:

Potassium permanganate	..	..	7.9 grammes
Distilled water	..	..	1,000 c.c.
1 c.c. corresponds to 0.00375 gramme of uric acid.			

The crimson fluid as it drops in is at first rapidly decolorized; presently the colour disappears more slowly. When the liquid in the beaker retains a distinctly pink colour for fifteen seconds, the end reaction is reached. Suppose 8 c.c. of permanganate solution have been added;

$\therefore$ in 100 c.c. of urine there are 8×0.00375 grammes uric acid—*i.e.*, 0.03 per cent.

The average total uric acid in twenty-four hours is 0.5 gramme. (N.B.—For very accurate estimations certain 'corrections' have to be made.)

5. Estimation of Chlorides.

Volhard's Method.—This consists in adding excess of silver nitrate to precipitate all the chlorides, and

determining the excess of silver nitrate with standard solution of potassium sulphocyanide.

Standard Silver Nitrate Solution.

Fused silver nitrate	..	..	29.07 grammes
Distilled water to	..	..	1,000 c.c.

1 c.c. corresponds to 0.01 gramme of sodium chloride.

Standard Potassium Sulphocyanide (KCNS) Solution.

Potassium sulphocyanide	..	..	8.29 grammes
Distilled water to	..	..	1,000 c.c.

2 c.c. corresponds to 1 c.c. of standard silver nitrate solution.

A strong solution of iron alum is also required as indicator, and some pure nitric acid.

To 10 c.c. of urine in a 100 c.c. flask add 5 c.c. of pure nitric acid and excess of standard silver solution, say 25 c.c. measured from a burette. Dilute with water to 100 c.c. Filter off the precipitated silver chloride. Pipette 50 c.c. of the filtrate into a beaker, add 5 c.c. of iron alum solution, and then standard KCNS solution from a burette. The end reaction is the appearance of a permanent red colour, due to formation of sulphocyanide of iron when all the silver has been precipitated.

Suppose 8 c.c. of KCNS solution were thus needed—

2 c.c. of KCNS solution corresponds to 1 c.c. of standard silver solution ;

$\therefore$ 8 c.c. of KCNS solution corresponds to 4 c.c. of standard silver solution.

$\therefore$ in 50 c.c. of the filtrate there were 4 c.c. of silver solution in excess ;

$\therefore$ in 100 c.c. there were 8 c.c. of silver solution in excess.

But in the 100 c.c. there were 10 c.c. of urine ; therefore, of the 25 c.c. of silver solution added to the 10 c.c. of urine, 8 c.c. were in excess.

$\therefore$ to precipitate the chlorides in 10 c.c. of urine $25 - 8$ c.c. silver solution are required = 17 c.c. But 1 c.c. silver solution corresponds to 0.01 gramme of sodium chloride.

$\therefore$ 10 c.c. of urine contain 17×0.01 grammes of sodium chloride;

$\therefore$ 100 c.c. of urine contain $17 \times 0.01 \times 10$ grammes of sodium chloride = 1.7 per cent.

The method is exceedingly accurate, and the end reaction very definite.

Upon full diet about 15 grammes of sodium chloride are excreted per diem.

Mohr's Method.—To 10 c.c. of urine in a beaker add 100 c.c. of distilled water and 20 drops of saturated potassium chromate solution. Add standard silver nitrate from a burette; a distinct red colour appears when all the chlorides have been precipitated. Suppose 12 c.c. of silver nitrate solution have been added;

$\therefore$ in 10 c.c. urine there are 12×0.01 grammes of sodium chloride;

$\therefore$ in 100 c.c. urine there are $12 \times 0.01 \times 10$ grammes of sodium chloride = 1.2 per cent.

6. Estimation of Phosphates.

The following reagents are needed:

(1) *Standard Uranium Nitrate Solution.*

Uranium nitrate 35.5 grammes

Distilled water to 1,000 c.c.

1 c.c. corresponds to 0.005 gramme of phosphoric acid.

(2) *Sodium Acetate Solution*

Sodium acetate 10 grammes.

Glacial acetic acid 10 c.c.

Distilled water to 100 "

(3) *Potassium Ferrocyanide : Saturated Solution.*

To 50 c.c. of urine add 5 c.c. of sodium acetate solution. Warm to 80° C. Add standard uranium nitrate from a burette. The end reaction is the

appearance of a distinct brown colour when a drop of the fluid in the beaker is added to a drop of the potassium ferrocyanide upon a white slab.

Suppose 11 c.c. of uranium nitrate have been added; then in 50 c.c. of urine there are 11×0.005 grammes of phosphoric acid ;

$\therefore$ in 100 c.c. of urine there are $11 \times 0.005 \times 2$ grammes of phosphoric acid = 0.11 per cent.

On full diet the average is 3.5 grammes per diem.

7. **Estimation of Sulphates** (see p. 44).

8. **Cryoscopy.**—By this is meant the determination of the degree below zero Centigrade to which the freezing-point of the urine is reduced. It has been stated that it is of some value in determining the efficacy of the kidney functions. The process is complicated, and needs special apparatus. Full directions as to use are sold with the latter, so that they need not be given here.

9. **Estimation of the Purin Bases.**—There are many substances allied to uric acid—xanthin, hypoxanthin, guanin, and so forth—which are known collectively by the term ‘purin bases.’ There is a special apparatus for estimating these—the purinometer. It is not difficult to use. Full instructions are sold along with the apparatus.

10. **Estimation of Ammonia.**—Although much stress is often laid upon estimations of sugar in diabetic urines, far less information of clinical value can be obtained from sugar determinations than from estimations of the ammonia present at the same time. The danger of glycosuria is the associated acidosis.

The greater the degree of acidosis the larger the proportion of ammonia-nitrogen in the urine as compared either to the total-nitrogen or to the urea-nitrogen. Most methods of estimating the ammonia, however, are too elaborate for any but special laboratories. One of the simplest is Malfatti's, which depends on the fact that when a solution of an ammonium salt is treated with formaldehyde, urotropin is formed according to the following equation :

The acid thus liberated can be determined by titration as follows : Measure 25 c.c. of urine into a 250 c.c. flask ; add 50 c.c. of ammonia-free distilled water, and 4 drops of 1 per cent. alcoholic solution of phenolphthalein as indicator ; add decinormal soda-solution (p. 129) carefully from a burette until a definite pink colour is established in the flask ; now add 5 c.c. of commercial formalin that has been carefully neutralized. Acids that have up to this point been combined with ammonia are now liberated according to the formula above, and the pink colour disappears ; more decinormal soda is added till the pink just returns again. Each 1 c.c. of additional decinormal soda required for this is equal to 0.0017 gramme of NH_3 . Mathison has found that the method can be made still more accurate, and the end points of the reactions still more sharply defined, by adding 15 grammes of powdered neutral potassium oxalate to the urine, and shaking for two minutes before the titration is begun, the object being to precipitate the calcium.

CHAPTER II

EXAMINATION OF THE BLOOD

BLOOD is best obtained from the lobule of the patient's ear, which, after cleaning with soap and water and thorough drying, should be held firmly at its base between the finger and thumb of the left hand, and given one *rapid* deep puncture with the needle. An ordinary straight surgical needle answers admirably; a straight Hagedorn's needle is best of all. The blood should well up freely, and squeezing should be avoided, lest lymph be expressed with it.

Enumeration of the Red and White Corpuscles:—

The best instrument to use is the **Thoma-Zeiss Hæmocytometer**; this consists of: (1) counting chamber; (2) specially ground cover-glass; (3) red corpuscle pipette; (4) white corpuscle pipette.

The counting chamber (Fig. 28) is made of an outer flat ring of glass, R, separated by a circular trench, T, from a central platform, P, firmly cemented to a glass slide. The outer ring is $\frac{1}{10}$ mm. higher than the platform, so that the cover-glass, resting flat upon the former, is separated from the latter by $\frac{1}{10}$ mm. The centre of the platform is ruled into minute squares

(Fig. 29), each side of which measures $\frac{1}{20}$ mm.; the cubic space over each small square, between the platform below and the cover-glass above, is therefore

FIG. 28.—Blood-counting Chamber.

T=trench, P=platform, R=ring, S=slide, C=cover-glass.

(1) Surface view. (2) Side view in section.

$\frac{1}{20} \times \frac{1}{20} \times \frac{1}{10}$ cub. mm. As a rule the small squares are arranged in sets of sixteen by double rulings at

FIG. 29.—Hæmocytometer Squares and Blood Corpuscles as seen under fairly low power.

intervals. Such sets of sixteen small squares are termed 'big squares.'

The stem of the red corpuscle pipette is graduated into tenths up to a mark 1; the bulb contains a glass

bead to facilitate mixing; above the bulb is a mark 101 (Fig. 30, A).

The white corpuscle pipette is similar, but the bore is wider, the bulb smaller, and the upper graduation is 11 instead of 101 (Fig. 30, B).

The diluting fluids required are the following:

Hayem's Fluid, for dilution in counting red corpuscles:

Mercury perchloride	..	0.25	gramme
Sodium chloride	..	0.5	..
Sodium sulphate	..	5.0	grammes
Distilled water	..	100.0	c.c.

Or one may use a 0.8 per cent. solution of common salt, which is isotonic with human blood, and therefore neither lyses nor crenates the corpuscles.

Thoma's Fluid, for dilution in counting white corpuscles:

Acetic acid (B.P.)	..	..	1.0 c.c.
Gentian violet solution	..	..	0.5 ..
Distilled water	..	..	100.0 ..

In this fluid the red corpuscles are lysed; their ghosts alone are seen. The white corpuscles stand boldly out, their nuclei being just tinged with the stain.

It is convenient to reserve for blood-counts, either upon a small tray or in a case, five wide-necked stoppered bottles. In the first keep Hayem's fluid, in the second Thoma's fluid, in the third, fourth, and fifth distilled water, absolute

FIG. 30.—Hæmocytometer Pipettes.

A, for diluting blood to count the red cells; B, for diluting blood to count the white cells.

alcohol, and ether respectively. The three latter are for cleaning the pipettes. Immediately a pipette is done with blow out the contents; draw up distilled water from bottle 3 into the bulb, and blow it out again; repeat this three times. Now draw up absolute alcohol from bottle 4 in a similar way three times. Finally, draw up ether from bottle 5 three times; after blowing out the ether the last time, draw air through the bulb. The water cleans the pipette, the alcohol dehydrates it, the ether replaces the alcohol, itself rapidly evaporates, and leaves the apparatus clean and dry, ready for use. Done at once, the cleaning takes a minute; left till next day, perfect cleaning may be difficult, or, worse still, the pipette may be completely clogged.

To Count the Red Corpuscles.—Prick the ear, wipe away the first drop of blood, wait till a second the size of a hemp-seed has collected, immerse the end of the pipette in it, and draw blood up to the mark 0.5. Quickly wipe the end of the pipette, plunge it into bottle 1, draw up Hayem's fluid to the mark 101; pinch the rubber tube, hold the pipette horizontally, and rapidly roll the bulb to and fro to mix the contents thoroughly. The blood dilution is 1 in 200. The upper mark is 101, and not 100, because the Hayem's fluid in the stem up to the point 1 takes no part in the mixture. Get rid of it in the first drop that is blown from the pipette, and then put part of the next drop on to the central platform of the counting chamber. With a little practice it is easy to judge the exact amount required for this. Apply the cover-glass, and press it down. If the fluid runs off the platform into the

trench, too big a drop has been taken. Part of a drop only is required; when covered, it should just extend to the margin of the platform, but not over the edge. If it do so, or if a bubble of air get in, clean the counting chamber, and take a fresh sample. There are various modifications of the Thoma-Zeiss instrument on the market; one has a special clamp which is convenient for keeping the cover-slip accurately in position. Enumerate the corpuscles under a $\frac{1}{6}$ -inch objective, counting the number in each of the sixteen small squares which make up one big square; and count five sets of sixteen small squares.* The variations are considerable. The following is an example of a count in one big square:

3	8	10	5	} Total 100.
7	7	9	1	
6	1	13	5	
10	6	5	4	

Corpuscles often overlap dividing lines. Such are apt to be counted twice, unless a rule be made to

FIG. 31.—Counting Red Corpuscles which overlap the Dividing Lines.

* The reason for counting the red cells in five big squares rather than in more is that the subsequent calculations are thereby greatly simplified (see p. 70).

count into a square those which overlap its top and right hand, but not those over the bottom and left hand, sides (Fig. 31).

Having counted five big squares, the figures might be:

$$\left. \begin{array}{ccc} 100 & 92 & 110 \\ 96 & 102 & \end{array} \right\} \text{Total 500.}$$

The average is in this case 100 to the big square, or $\frac{100}{16}$ to the small square. But the volume of fluid over each small square is $\frac{1}{20} \times \frac{1}{20} \times \frac{1}{10}$ cub. mm. (p. 66).

$\therefore$ in $\frac{1}{20} \times \frac{1}{20} \times \frac{1}{10}$ cub. mm. of 200 times diluted blood there were $\frac{100}{16}$ red corpuscles.

$\therefore$ in 1 cub. mm. of 200 times diluted blood there were $\frac{100}{16} \times \frac{20 \times 20 \times 10}{1}$ red corpuscles.

$\therefore$ in 1 cub. mm. of original blood there were—

$$\frac{100}{16} \times \frac{20 \times 20 \times 10}{1} \times \frac{200}{1} \text{ red corpuscles.}$$

$$= 5,000,000 \text{ red corpuscles per cub. mm.}$$

It will thus be seen that, with a dilution of 1 in 200, normal blood containing 5,000,000 red corpuscles per cub. mm. gives 100 to the big square. Hence, the average number in a big square, the dilution being 1 in 200, gives the percentage of red corpuscles without further calculation; and the number in five big squares with the addition of ...0,000, is the number of red corpuscles per cub. mm. of original blood.

Importance of Counting the Red Corpuscles.—Less can be learned from the number of red corpuscles alone than from the number of red corpuscles and the amount of hæmoglobin at the same time (p. 73). A patient

may have a normal number of red corpuscles and yet may be very anæmic from deficiency of hæmoglobin. Apart from the hæmoglobin, however, a count of the red corpuscles may be by itself valuable in the following ways:

1. In congenital heart disease (*morbus cæruleus*) the red corpuscles are frequently much increased—in some cases to 7,000,000 per cub. mm.; in a few to double this. A similar but less marked increase occurs in those cyanosed from other causes, such as fibroid lung, mitral stenosis, or chronic acetanilide poisoning.

2. Those who live at high altitudes usually have more than 5,000,000 per cub. mm.

3. As an anæmic patient improves, the red corpuscles increase more rapidly than does the hæmoglobin. Hence, enumeration of the red corpuscles from week to week affords a valuable means of gauging the effect of treatment.

4. There is a peculiar disease in which the main characteristics are—(a) considerable enlargement of the spleen, (b) great increase in the number of red corpuscles (polycythæmia), up to 10,000,000 per cub. mm. or more. It is called ‘splenomegalic polycythæmia.’

To Count the White Corpuscles.—Proceed in an exactly similar manner, but with the other pipette. The bore being larger, a bigger drop of blood is required. Care is needed to prevent the fluid running out. Fill to the mark 0.5 with blood; dilute with Thoma’s fluid to the mark 11. The dilution is then 1 in 20. Use the same counting chamber, with the same precautions. In normal blood, diluted twenty times,

there are seldom more than three leucocytes in each *large* square, and often none at all in some. Count sixteen large squares; suppose there are a total of twenty leucocytes, the average in one large square would be $\frac{20}{16}$, and in one small square $\frac{20}{16 \times 16}$.

$\therefore$ in $\frac{1}{20} \times \frac{1}{20} \times \frac{1}{10}$ cub. mm. of twenty times diluted blood there are $\frac{20}{16 \times 16}$ leucocytes.

$\therefore$ in 1 cub. mm. of twenty times diluted blood there are $\frac{20}{16 \times 16} \times \frac{20 \times 20 \times 10}{1}$ leucocytes.

$\therefore$ in 1 cub. mm. of original blood there are—
 $\frac{20}{16 \times 16} \times \frac{20 \times 20 \times 10}{1} \times \frac{20}{1}$ leucocytes.
 $= 6,250$ leucocytes per cub. mm.

The percentage error in an estimation of this kind must clearly be great; nevertheless, the clinical variations in the leucocyte counts are even greater, so that the method affords data of sufficient accuracy for most clinical purposes.

The normal number of leucocytes is usually given as 5,000 per cub. mm.; but it must be remembered that there are daily variations in every person, and that it is the rule rather than the exception to find as few as 4,000 and as many as 12,000 in the same person at different times on the same day. Unless the numbers are above 15,000 or below 4,000 they cannot be called abnormal.

Importance of Counting the White Corpuscles.—Leucocytosis—*i.e.*, an increase in the total number of leucocytes to more than 18,000 per cub. mm.—often accompanies deep-seated suppuration, and may afford valu-

able help in diagnosis. Absence of leucocytosis does not exclude suppuration. A leucocytosis which increases at successive counts is strongly in favour of progressive pus formation, and therefore of operation—for instance, in case of appendicitis. Suppuration often causes a leucocytosis up to 20,000 or 25,000; it rarely reaches 60,000, and almost never exceeds this.

In lobar pneumonia there is often leucocytosis up to 30,000 or more.

In typhoid fever and malaria without complication there is no leucocytosis, but rather a leucopenia—*i.e.*, an actual diminution in the leucocytes below 4,000 per cub. mm. In many cases in which obscure pyrexia suggests typhoid fever or malaria, leucocytosis to 20,000 has led to the discovery of deep-seated pus—for example, in a pyosalpinx or hepatic or appendicular abscess.

In the diagnosis of the leukæmias, lymphatic and spleno-medullary, from Hodgkin's disease and splenic anæmia, a leucocyte count is indispensable. In the two last there is no leucocytosis. In lymphatic leukæmia the leucocytes rise to as many as 150,000 per cub. mm., or more, whilst in the spleno-medullary form 200,000 are often found, and even 600,000.

Estimation of the Hæmoglobin.

Haldane's Hæmoglobinometer or Gower's Hæmoglobinometer.—The older forms of hæmoglobinometer need not be described. Where coal-gas can be obtained no instrument is better than Haldane's. Where there is no supply of coal-gas, Gower's may be used. The two can be combined. Haldane's

apparatus (Fig. 32) consists of (1) a sealed tube, A, containing a standard* solution of carboxyhæmoglobin; (2) a graduated tube, B, for diluting the blood in; (3) an india-rubber stand, C, for these tubes; (4) a pipette, D, for measuring the blood; (5) a brass U-tube, E, for introducing the coal-gas into B; (6) a drop-bottle, F, for distilled water. The method is

FIG. 32.—Haldane's Hæmoglobinometer.

as follows: Drop distilled water into the tube B up to the mark 10. Draw blood into the pipette up to the point indicated on it. Expel it into the distilled water in B without immersing the tip of the pipette. Shake the tube B. The blood is laked at once. Now wash out the pipette as directed on p. 68. Fit the short end of the brass U-tube on to an unlighted gas-burner, and push the tube B up over the longer limb until the latter nearly touches the

* It is standardized, not empirically, but by actual gas analysis of the blood from which it was prepared.

laked blood (Fig. 33). Turn on the gas, and slowly draw away the tube B. The air in its upper part is replaced by coal-gas. On shaking up the laked blood with the gas, the carbon monoxide of the latter converts the hæmoglobin into cherry-red carboxy-hæmoglobin. Repeat the introduction of coal-gas if necessary. Now add distilled water drop by drop to the tube B, mixing it up each time, until the tint of the fluid is identical with that of the standard carboxy-hæmoglobin in tube A. The graduation now corresponding to the upper level of the fluid in tube B is the percentage of hæmoglobin in the patient's blood, that of normal blood being 100. It is well to make two readings, the first when the tint is just perceptibly darker than the standard, the other when just paler, and to take the mean of the two. With practice the error becomes very small.

FIG. 33.

Gower's apparatus is very similar in its various parts, but the standard colour-tube contains carmine solution instead of carboxy-hæmoglobin, and the laked blood is immediately diluted drop by drop until the tint of mixed oxy- and reduced hæmoglobin is as nearly as possible identical with that of the carmine. The tints can only be made to agree in good daylight, whereas in Haldane's method the solutions are both the same colouring matter, carboxy-hæmoglobin, and therefore can be compared in any light, artificial or natural. Carboxy-hæmoglobin, moreover, is a very stable compound, and the blood can be kept for

examination at any time. Oxyhæmoglobin rapidly changes in shed blood, so that, with Gower's method, comparison must be made at once.

In country districts it is well to have both a carmine and a carboxyhæmoglobin standard colour-tube for the same apparatus. In the event of coal-gas being unprocurable the former could be used; in all other cases the latter.

Tallqvist's Hæmoglobinometer.—This is not strictly accurate, but it is very simple, and for many purposes the readings are nearly enough. It consists of a book containing papers of various tints of red and some white blotting-paper. A drop of the patient's blood is received upon a piece of the latter, and as soon as it is dry its colour is compared with those of the tinted papers until one to match it is found. The percentage of hæmoglobin is thus found in a couple of minutes, but the readings are not strictly accurate, as a rule, and in the diagnosis of such a condition as pernicious anæmia the more accurate Haldane method of estimating the hæmoglobin is essential.

Importance of Estimating the Hæmoglobin.—The severity of most anæmias is better gauged by the amount of hæmoglobin than by the number of red corpuscles. In chlorotic girls the latter may be but little below normal, and yet the hæmoglobin may be very much reduced.

The **colour index** of the blood is the ratio of hæmoglobin to red corpuscles. Three types of anæmia are based upon it. In healthy blood the colour index is 1; in anæmia directly following hæmorrhage it is still 1; in the chlorotic type it is

less than 1; in the pernicious type greater than 1. These characteristics are represented by arbitrary examples in the following table:

	Percentage of Red Corpuscles.	Percentage of Hæmo- globin.	Colour Index.
Healthy person	100	100	$\frac{100}{100}$, or 1.
Anæmia from hæmorrhage	80	80	$\frac{80}{80}$, or 1.
Chlorotic type	60	30	$\frac{30}{60}$, or 0.5.
Pernicious type	20	30	$\frac{30}{20}$, or 1.5.

Thus, in anæmia immediately following hæmorrhage red corpuscles and hæmoglobin are equally diminished; in chlorotic types the hæmoglobin is diminished more than are the red corpuscles; in pernicious anæmia the red corpuscles are more diminished than is the hæmoglobin.

Most anæmias are of the chlorotic type. For example, chlorosis itself; tuberculous and cancerous cachexias; infective endocarditis and other toxæmias; the later stages of Hodgkin's disease, the leukæmias, splenic anæmia.

To the pernicious type belong pernicious anæmia and some cases of ankylostomiasis. Without estimations of both red corpuscles and hæmoglobin, and a calculation of the colour index, some cases of infective endocarditis or carcinoma of the stomach may be wrongly diagnosed as pernicious anæmia.

The Preparation and Examination of Blood Films.

The least wasteful way of storing the various stains is to use cylindrical glass jars (Fig. 34), $3\frac{3}{4}$ inches high, $1\frac{1}{2}$ inches diameter, with ground stoppers. Films can be immersed bodily in these, and the same stain used again and again.

FIG. 34.—Bottle for staining Blood Films in.

Films may be made either upon microscope slides or upon cover-glasses. If the latter are used, they should be the thinnest possible (No. 1); others are too thick for use with the oil-immersion lens. In almost all cases, however, it is more convenient to make the film upon a slide. When it is finished and stained, there is no need to cover it, nor to use Canada balsam; cedar-wood oil may be put directly on to the dried and stained film, and the latter can be examined with the oil-immersion lens without any other covering at all. The same applies to films of pus to be examined for bacteria. The slides may even be preserved for future reference uncovered; but if they are to be kept for any length of time, it is better to cover them in the usual way to prevent their getting scratched.

To make a film, a good plan is the following: Cut a cigarette-paper at right angles to its length into strips the same width as the slips. Prick the ear; wipe away the first drop of blood; when the second is as large as a big pin's head, smear the ungummed

end of a strip over its summit. Holding the other end, lay the smeared surface upon a cover-glass, and draw it steadily along (Fig. 35). The thickness of the film can be varied by altering the angle at which the paper is held to the glass. The blood

FIG. 35.—Cigarette-paper Method of making Blood Film.

should dry almost as quickly as it is spread. It is too thick if it stand in slow-drying pools. Use a fresh strip for each film; never smear the cover-glass over twice.

Another plan which is very simple and good, but which is more easily demonstrated than described, is to receive a small drop of blood on to the surface of one slide near its end, and then, laying the edge of another slide across the first through the middle of the blood-drop, push the latter steadily along the surface of the first slide. A perfectly even film is thus obtained.

Unless carefully fixed, the corpuscles wash off the glass in the process of staining.

Methods of Fixing Films.—Four of the chief methods of fixing films are as follows:

1. **In Absolute Alcohol and Ether.**—This requires no special apparatus. Fill three-quarters of one of the glass jars with absolute alcohol and sulphuric ether, equal parts. Drop in the films when dry, and leave them till next day. An hour is said to be enough, but twenty-four are better.

2. **By Heat.**—A hot-air oven with a good gas regulator is required. Fix for two hours at a temperature

of 120° C., or for half an hour at 140° C. For certain stains fixing by heat is essential, but it is inconvenient in private practice.

3. **By Jenner's Method** (see below), which fixes any stains at the same time.

4. **By Formalin.**—Immerse the films in a 1 per cent. solution of formalin for one minute.

Methods of Staining Films.—Many different ways of staining blood films have been recommended, but it will serve no useful purpose to describe in detail any but those which are widely acknowledged to be the simplest and best—namely :

For ordinary purposes	{	1. Jenner's method.
		2. The hæmatoxylin and eosin method.
For malaria and other hæmatozoa	{	3. Leishman's method.

1. **Jenner's Method.**—This has a paramount advantage, in that it fixes and stains at the same time, and therefore is very rapid. The films may be stained and ready for examination ten minutes after the patient's ear has been pricked. The method is a little uncertain, but practice makes it less and less so. There is considerable variation in the efficacy of different bottles of the stain. The main point, however, upon which success or otherwise depends, is the condition of the wash water.

Formula for Jenner's Stain.

Crystalline methylene-blue eosin	..	0.5 gramme
Methyl alcohol		100 C.C.

It is well to keep the wash water in a special jar, and to use the same again and again. It is essential that distilled water be used ; but even distilled water may be too alkaline, from the presence of ammonia, in which case too much of the blue stain will be removed. It is possible to correct this by accurate titration with very dilute acetic acid ; but a simpler plan is to pour a few drops of the Jenner's stain into the jar of wash water, till the latter is tinged blue, and then, dirty though it may appear to be, to keep this carefully for the washing of future films. To stain, drop the film, as soon as it has dried after making, into the Jenner's stain for five minutes ; then lift it out, allow as much stain as possible to run off, and transfer the film to the wash water ; move it gently to and fro for about half a minute ; when it looks quite pink, remove it from the wash water ; dry with filter-paper and examine.

The red corpuscles are bright pink ; their nuclei, if present, blue. The large lymphocytes have pale blue nuclei and paler blue clear protoplasm. Small lymphocytes have deep blue nuclei and a rim of pale blue protoplasm. Polymorphonuclear cells have blue nuclei, and pink protoplasm containing fine pink granules. Eosinophile corpuscles have blue nuclei and pink protoplasm full of very big red granules. Myelocytes have dark blue nuclei and pale blue protoplasm, the latter containing either blue streaks and granules or, in the case of eosinophile myelocytes, red granules. The granules in any basophile cells that may be present are deep violet. Bacteria, malaria, or filaria parasites are pale blue if stained at all.

2. **Delafield's Hæmatoxylin and Aqueous Eosin Method.**—The disadvantage of this method is that it is slow; its advantages are—first, that it is very certain; secondly, that no distilled water is necessary, for the washing not only may be, but must be, done with ordinary tap water.

Delafield's Hæmatoxylin.—To 400 c.c. of a saturated aqueous solution of ammonia alum add 4 grammes of hæmatoxylin dissolved in 25 c.c. of absolute alcohol. Leave the solution exposed to the light and air in an unstoppered bottle for three or four days; filter, and add to the filtrate 100 c.c. of glycerin and 100 c.c. of methyl alcohol. Allow the solution to stand in the light until it is a dark colour, refilter, and preserve in a stoppered bottle.

Aqueous Eosin Solution.

Eosin (aqueous)	..	..	..	1 gramme
Distilled water	..	..	..	100 c.c.

The films should be fixed for twenty-four hours in absolute alcohol and ether (p. 79); the procedure is then as follows:

Wash with tap water to remove alcohol and ether; blot with filter-paper; immerse in hæmatoxylin for five minutes; wash thoroughly in running tap water; the slight alkalinity of the latter serves to 'blue' the hæmatoxylin; blot; immerse in eosin for thirty seconds; wash again in running tap water; dry with filter-paper, and examine.

The red corpuscles are stained pink; their nuclei, if present, blue. The small and large lymphocytes have deep purplish-blue nuclei and pale protoplasm of the

same colour. Myelocytes stain similarly to large lymphocytes, but are distinguished by the granules in their protoplasm. Polymorphonuclear cells have bluish-purple nuclei and bright pink protoplasm, the latter showing fine granules far less distinctly than with Jenner's stain. Eosinophile cells have bluish-purple nuclei and bright pink protoplasm crowded with large pink granules. Basophile cells and hæmatozoa do not stain well by this method.

3. **Leishman's Method.**—This is the stain, *par excellence*, for malaria parasites. It is also a very good way of staining ordinary blood films; it is similar to Jenner's, and, like the latter, fixes and stains at the same time. The stain consists of a modified eosinate of methylene blue dissolved in methyl alcohol; it is the latter which fixes the films. The formula is—

Eosinate of methylene blue (Grübler)	0·15	gramme
Methyl alcohol (Merck)	100	c.c.

The dye dissolves with difficulty in the methyl alcohol. A good plan is to put the eosinate into a mortar and pour on 20 c.c. of the alcohol at a time, rubbing well with the pestle till the whole 100 c.c. has been used. Place the slide, film uppermost, upon a perfectly horizontal table; filter just enough of the stain on to the film to cover the latter evenly; leave for one minute, covering with a large watch-glass to prevent evaporation and precipitation of the stain. Now pour on distilled water drop by drop till as much has been added as there was stain originally; leave the slide for another five minutes, and then wash thoroughly with distilled water, to which some Leish-

man's stain has been added as described for Jenner's stain (p. 81); blot; dry, and examine.

The red corpuscles are bright pink; the nuclei of the leucocytes are dark reddish-purple; malaria parasites are bright blue with almost black granules; filaria parasites also stain blue.

It is probable that anyone who has tried the above methods will not feel any need for others. The following have all at one time or other been much in vogue, but they are now less in fashion.

4. **Methylene Blue and Eosin.**—(a) Alcoholic solutions for films fixed by heat:

Alcoholic Eosin.

Eosin	..	..	..	0.5 gramme
Distilled water	..	..	..	30.0 c.c.
Absolute alcohol	..	..	..	70.0 „

Alcoholic Methylene Blue.

Methylene blue	..	..	..	1.5 gramme
Rectified spirit	..	..	..	100.0 c.c.

Allow to stand for twenty-four hours, shaking vigorously from time to time. Filter. Stain in the eosin first, for one minute; wash in distilled water; blot; stain in methylene blue for five minutes; wash in distilled water, and dry.

(b) Aqueous solutions, for films fixed in alcohol and ether:

Watery Eosin (see p. 82).

Watery Methylene Blue.

Methylene blue	..	..	..	1.5 gramme
Distilled water	..	..	..	100.0 c.c.

Shake up daily for two weeks. Filter

Stain the film in eosin for one minute ; wash in distilled water ; blot ; stain in methylene blue for five minutes ; wash in distilled water, and dry.

These two methods are uncertain, though when successful they are pretty. If the methylene blue stains faintly, try warming it gently:

5. Ehrlich's Triple Stain :

Saturated solution of orange G. ..	..	13·14 c.c.
Saturated solution of acid fuchsin ..	..	6·7 "
Distilled water	..	15·0 "
Absolute alcohol	..	15·0 "
Saturated solution of methyl green ..	..	12·5 "
Absolute alcohol	..	10·0 "
Glycerin	..	10·0 "

Measure off the ingredients in the order named. The mixture may be used at once ; but to make each of the saturated solutions it contains requires several days. The stain keeps well, but whereas one batch may stain excellently, the next may fail entirely.

Films must be fixed by heat. Immerse in the stain for one to five minutes ; the time varies with different bottles of stain and with the length of time the films have been fixed by heating. Wash with *distilled* water, and dry.

6. Romanowski's Method for Malarial Parasites.—This is inferior to Leishman's stain in that the films have to be fixed, either by alcohol and ether or by heat, before staining. The stain consists of a mixture of 1 part of saturated aqueous methylene blue and 2 parts of 1 per cent. aqueous eosin. The slides, when fixed, are immersed in this for two hours, washed in distilled water, and dried.

7. Joseph's Modification of the Romanowski

Satin yields even better results than Leishman's stain according to some authorities. It is made up as follows :

Solution I. : Polychrome Methylene Blue.—Dissolve 1 gramme of Grübler's ordinary methylene blue in 100 c.c. of distilled water ; after twenty-four hours decant the supernatant fluid, and to 90 c.c. add 10 c.c. of a 1 per cent. aqueous solution of caustic soda, and heat on a water-bath to 60° C. for twenty-four hours.

Solution II. : Eosin.—Dissolve 1 gramme of Grübler's yellow-shade water-soluble eosin in 5 c.c. of absolute alcohol, and add 95 c.c. of distilled water.

To make up the stain, take 35 c.c. of Solution I., add 50 c.c. of methyl alcohol (Merck), shake well, add 30 c.c. of Solution II., and make up to 350 c.c. with methyl alcohol. The stain needs neither filtration nor evaporation, may be used at once, and will keep for several months if the bottles are well stoppered. The method of using the stain is precisely like Leishman's (p. 83), and the appearances obtained are similar.

Examination of the Film.—This should be made with $\frac{1}{12}$ -inch oil-immersion lens, a substage condenser, and a mechanical stage. Select that part of the film in which the corpuscles do not overlap. A differential leucocyte count may be made at the same time. Look for the following points :

Changes in the Red Corpuscles.—In a well-fixed film of normal blood these are smooth in outline, quite round, of approximately equal sizes, and stained rather less at the centre than at the periphery. If the cells are crenated or run together into batches so as to

compress one another into polygonal shapes (Fig. 37), the fault is in the film-making.

In severe anæmias red corpuscles may vary—

1. **In Size.**—A normal red corpuscle, 7μ in diameter, is termed a **normocyte**; a smaller is a **microcyte**; a larger a **megalocyte** (Figs. 36 and 38).

2. **In Shape.**—Instead of round, they may be pear-shaped, oval, fiddle-shaped, or ovoid; such corpuscles are termed **poikilocytes**, and the condition **poikilocytosis** (Fig. 39).

3. **In Staining Capacity.**—A normocyte should stain throughout. When hæmoglobin is very deficient, the corpuscles may resemble quoits, a thin rim of red surrounding a colourless interior. In severe anæmias, and particularly in pernicious anæmia, some of the red cells lose their affinity for acid dyes, and stain bluish-red or violet instead of pink. The condition is termed **polychromatophilia** or **polychromasia**. It has little import except that it is never found in normal blood. Sometimes an otherwise red-stained corpuscle exhibits fine dots of blue in its substance; this condition is a prominent feature of certain cases of plumbism, and it is known as **punctate basophilia** (Fig. 41).

4. **In the Presence of Nuclei.**—Nucleated red corpuscles have a deep blue nucleus and a rim of pink protoplasm, which readily distinguishes them from the lymphocytes, in which both nuclei and protoplasm are blue. A nucleated normal-sized red corpuscle is termed a **normoblast**; a nucleated small corpuscle a **microblast**; a large one a **megaloblast**. Occasionally

more than one nucleus is present in a still larger cell, termed a **gigantoblast** (Fig. 40).

Importance of the Changes in the Red Corpuscles.—Poikilocytosis, alterations in size, and presence of nucleated red corpuscles, may occur in *any* severe anæmia. Though seen most typically in pernicious anæmia, these changes are not pathognomonic. If there is a *considerable preponderance of megalocytes*, together with poikilocytosis, and no leucocytosis, there is great likelihood that the condition is pernicious anæmia.

A high proportion of nucleated red corpuscles indicates very severe anæmia, and a less good prognosis; conversely, if the proportion of nucleated forms diminishes, the patient is improving.

The Differential Leucocyte Count.—A mechanical stage is essential. Omit the thicker parts of the film; in the remainder examine the leucocytes consecutively, allotting each to its appropriate class. Begin at one end of the film, say the top right-hand corner, and screw it continuously past the eye until the left edge of the thin part is reached; then, watching through the microscope, screw it down one field's breadth, and then count the leucocytes continuously back again until the right edge is reached; turn the film down another field's breadth, and count it transversely across to the left edge again; and so on, backwards and forwards, until the required number of corpuscles, say 250, has been enumerated. In normal blood these will fall into four main groups, to which various names have been given. They are—

1. **Small Lymphocytes** (Fig. 42).—These are

DESCRIPTION OF PLATE I

FIG. 36.—Normocytes.

- „ 37.—Badly Fixed and Crenated Red Cells.
- „ 38.—Megalocytes and Microcytes.
- „ 39.—Poikilocytes.
- „ 40.—Nucleated Red Corpuscles.
- „ 41.—Punctate Basophilia.
- „ 42.—Small Lymphocyte.
- „ 43.—Large (Hyaline) Lymphocyte.
- „ 44.—Polymorphonuclear Leucocyte.
- „ 45.—Coarsely Granular Eosinophile Leucocyte.
- „ 46.—Myelocyte.
- „ 47.—Eosinophile Myelocyte.
- „ 48.—Basophile Leucocyte.
- „ 49.—Malaria Parasite : Ring Form.
- „ 50.—Filaria Embryo and Red Cells.
- „ 51.—Malaria Parasite : Crescent Form.
- „ 52.—Spirilla of Relapsing Fever and Red Cells.

The various cells, except those of Fig. 50, are drawn approximately to scale as they are seen under an oil-immersion lens.

PLATE I.

Fig. 36.

Fig. 37.

Fig. 38.

Fig. 39.

Fig. 40.

Fig. 41.

Fig. 42.

Fig. 43.

Fig. 44.

Fig. 45.

Fig. 46.

Fig. 47.

Fig. 48.

Fig. 49.

Fig. 50.

Fig. 51.

Fig. 52.

slightly larger than red corpuscles, variable in size, characterized by a comparatively large, deep-staining, spherical nucleus and a rim of protoplasm which is sometimes very narrow, sometimes considerable. These constitute from 20 to 36 per cent. of leucocytes in health.

2. **Large Lymphocytes** (Fig. 43).—Also termed 'hyaline'; these are much larger cells, with notched or kidney-shaped nucleus, and much pale-staining clear protoplasm. These constitute from 2 to 10 per cent. in health. Some cells are intermediate between typical small and typical large lymphocytes, classed with the former by some, by others with the latter. Most class all such intermediate cells with the small lymphocytes, reserving the term 'large lymphocytes' for the typical large cells only.

3. **Polymorphonuclear Cells** (Fig. 44).—These are somewhat smaller than large lymphocytes; usually circular, with a well-defined nucleus variously lobed and twisted. The name of this cell is derived from this characteristic of the nucleus, which may be horseshoe-shaped, or twisted like the letters Z, S, E. Other names used are 'neutrophile' and 'finely granular oxyphile.' The protoplasm is abundant, and shows minute red granules; these cannot be mistaken for the coarse granules of the next variety when once the latter have been seen. The polymorphonuclear cells constitute about 65 per cent. of all leucocytes in health.

4. **Coarsely Granular Eosinophile Cells, or 'Eosinophiles'** (Fig. 45).—These are slightly larger than the polymorphonuclear cells, with a similar twisted nucleus, and protoplasm, which is crowded with unmis-

takable coarse red granules. They vary from 0·5 to 5 per cent. in health.

Two other forms of leucocytes are found under certain pathological conditions, namely :

5. **Myelocytes** (Fig. 46).—These are large cells with round or oval pale-staining nuclei, and much protoplasm, in which there are many granules and streaks, by which these cells are distinguished from large lymphocytes, whose protoplasm is hyaline. In certain myelocytes the granules stain bright red ; these **eosinophile myelocytes** (Fig. 47) might be mistaken for coarsely granular eosinophile cells were it not that the nuclei of the latter are lobed, whilst that of each myelocyte is round.

6. **Basophile Cells** (Fig. 48).—These are about the size of small lymphocytes, which they resemble, except in that their protoplasm contains half a dozen or more very big granules, which are stained deep violet by Jenner's stain.

Importance of the Differential Leucocyte Count.—So variable are the proportions of the white corpuscles in health that too much stress must not be laid upon comparatively slight increases in any particular variety : nevertheless, the following broad facts about them are clinically important :

The Small Lymphocytes are increased :

Up to 35 per cent. or more in many healthy infants ; up to 40, 50, or 60 per cent. in children suffering from congenital syphilis, whooping-cough, and many non-suppurative infantile complaints.

In typhoid fever, especially in mild attacks ; in cases of obscure pyrexia, without leucocytosis, an increase

of small lymphocytes up to 35 or 40 per cent. favours a diagnosis of enterica, and in the tropics may be a valuable means of excluding malaria.

In lymphatic leukæmia, in which there is leucocytosis to 50,000 or 150,000 or more, the small lymphocytes often reach as high as 95 per cent., a point which is pathognomonic of the disease. There is a much slighter relative increase without leucocytosis in Hodgkin's disease and in pseudo-leukæmia infantum (Von Jaksch's disease).

In many debilitating diseases in adults, such as syphilitic, malignant, or tuberculous cachexia, cirrhosis of the liver, and so forth, provided there be no associated abscess formation, the differential leucocyte count very often shows considerable relative increase of small lymphocytes and diminution of polymorphonuclear cells; no great clinical importance can be attached to the fact, except in excluding suppuration.

The Large Lymphocytes are increased :

In malaria, in which the total number of leucocytes is, as a rule, under 5,000 per cub. mm., and the large lymphocytes may be 15 to 30 per cent.; this affords a valuable means of diagnosis between malaria and enterica, and between malaria and hepatic abscess. The relative increase in large lymphocytes does not disappear immediately on giving quinine, as do the malaria parasites, so that it can be utilized as a point in the diagnosis of malaria, when no parasites can be found owing to quinine having been administered.

The Polymorphonuclear Cells are increased :

In lobar pneumonia and in cases of deep-seated suppuration, such as empyema, appendicular abscess,

pyosalpinx. Leucocytosis and a rise of polymorpho-nuclears to over 80 per cent. may often help in the diagnosis of deep-seated pus.

The Coarsely Granular Eosinophile Cells are increased :

In many cases of true asthma, when they may reach anything between 6 and 60 per cent.

In certain parasitic affections, notably in ankylostomiasis; in *Bilharzia hæmatobia*; in trichinosis; in filariasis; and in patients suffering from tapeworms. The eosinophilia may be slight—up to 10 per cent., for example. It is often between 10 per cent. and 20 per cent., and occasionally reaches 50 per cent. Nothing under 5 per cent. can be termed eosinophilia. There is, as a rule, no eosinophilia with *Oxyuris vermicularis* or *Trichocephalus dispar*; nor is there any with the parasitic skin diseases — tinea tonsurans, and so forth.

In certain skin diseases, particularly true pemphigus, erythema bullosum, herpes gestationis, and hydroa herpetiformis, where from 6 per cent. to 20 per cent. may be found. In other skin diseases, such as psoriasis or eczema, the majority of patients show no eosinophilia, though cases with increase to 5 per cent. or more occur occasionally.

The Myelocytes are present in large numbers in one disease only—namely, spleno-medullary leukæmia, where they may constitute 45 per cent. of all the white corpuscles. They are not, however, distinctive of this condition; 1 or 2 per cent. may be found in splenic anæmia, Hodgkin's disease, pernicious anæmia, or lymphatic leukæmia.

The Basophile Cells are not pathognomonic of any disease, though they are practically absent in health. They are most numerous in spleno-medullary leukaemia, reaching to 2 to 5 per cent. Smaller numbers are found in other anæmias, but what importance attaches to this is not yet known.

The following are some specimens of characteristic differential leucocyte counts.

S.=small lymphocytes ; L.=large lymphocytes ; P.=polymorphonuclear cells ; E.=coarsely granular eosinophile cells ; M.=myelocytes ; B.=basophile cells.

Normal Adult Blood.

			Per Cent.
S.	..	..	25
L.	..	..	8
P.	..	..	65
E.	..	..	2
M.	..	..	absent
B.	..	..	absent

Normal Child's Blood.

			Per Cent.
S.	..	..	38
L.	..	..	8
P.	..	..	53
E.	..	..	1
M.	..	..	absent
B.	..	..	absent

Spleno-medullary Leukæmia.

			Per Cent.
S.	..	..	12
L.	..	..	1
P.	..	..	37
E.	..	..	2
M.	..	..	45
B.	..	..	3

Lymphatic Leukæmia.

			Per Cent.
S.	..	..	91
L.	..	..	2.7
P.	..	..	6
E.	..	..	often absent
M.	..	..	0.2
B.	..	..	0.1

Malaria.

			Per Cent.
S.	..	..	20
L.	..	..	18
P.	..	..	60
E.	..	..	2
M.	..	..	absent
R.	..	..	absent

Eosinophilia.

			Per Cent.
S.	..	..	25
L.	..	..	8
P.	..	..	53
E.	..	..	14
M.	..	..	absent
B.	..	..	absent

Abscess Formation.				Various Cachexias without Pus Formation.			
Per Cent.				Per Cent.			
S.	..	..	12	S.	..	..	40
L.	..	..	2'5	L.	..	..	6'2
P.	..	..	85	P.	..	..	53
E.	..	..	0'5	E.	..	..	0'5
M.	..	..	absent	M.	..	..	absent
B.	..	..	absent	B.	..	..	0'3

It should be borne in mind that, apart from malaria and other parasitic conditions, the only diseases in which the blood examination affords pathognomonic evidence of the diagnosis are—

Spleno-medullary leukæmia.

Lymphatic leukæmia.

Pernicious anæmia.

In many other conditions a blood examination has to be made in order to exclude one or other of the above. In Hodgkin's disease, for example, the evidence afforded by the blood is negative. One cannot diagnose Hodgkin's disease by examining the blood; but having examined it and found a simple anæmia, leukæmia is excluded, and Hodgkin's disease remains as a possibility.

The following notes on the main features of the blood in different conditions may be useful:

Spleno-medullary Leukæmia.—Hæmoglobin index 1 or less; leucocytosis, possibly to 400,000 or more; myelocytes up to 45 per cent.

Lymphatic Leukæmia.—Hæmoglobin index 1 or less; leucocytosis up to 150,000 or more; small lymphocytes may be 95 per cent.

Hodgkin's Disease, Banti's Disease, and Splenic Anæmia.—Hæmoglobin index 1 or less;

no leucocytosis; leucocytes often in normal proportions—*i.e.*, there are no positive blood changes.

Pernicious Anæmia.—Hæmoglobin index greater than 1; no leucocytosis; white corpuscles in normal proportions or else lymphocytes relatively increased; in blood films preponderance of megalocytes.

(N.B.—In all the above the red corpuscles may vary in size, show poikilocytosis, and present varying numbers of nucleated forms.)

Chlorosis.—Hæmoglobin index less than 1; white corpuscles in normal proportions. Not infrequently the anæmia consists entirely in diminution of the hæmoglobin, the red corpuscles being little, if any, below normal.

Typhoid Fever.—Leucopenia and relative increase in the small lymphocytes.

Malaria.—Leucopenia and relative increase in the large lymphocytes. Parasites present at certain times.

Abscess Formation and Lobar Pneumonia.—Leucocytosis and relative increase in the polymorphonuclear cells. The pus must be under pressure; there is no leucocytosis when the abscess is draining; nor in superficial suppuration, such as impetigo; nor in cases where the organisms are not confined, as in fungating endocarditis; nor must the suppuration be very virulent—in cases of general suppurative peritonitis, for example, there is usually neither leucocytosis nor increase in the polymorphonuclear cells.

Cachexias without Suppuration.—Anæmia of the chlorotic type, with or without slight leucocytosis; relative increase of the small lymphocytes.

Morbus Cæruleus (Congenital Pulmonary Stenosis,

or Patent Interventricular Septum).—The red cells are remarkably increased—up to 10,000,000 per cub. mm., for example. The hæmoglobin is also increased, but to a less extent. The leucocytes remain normal.

Mitral Stenosis, Chronic Bronchitis, and Emphysema, and other lesions associated with more or less cyanosis: the blood shows changes similar to those of morbus cæruleus, but to a less extent.

Altitude.—Those who live on mountains have simple polycythæmia, similar to that of mitral stenosis, but with out cyanosis.

Tænia solium	}	Chlorotic anæmia, with moderate eosinophilia.
Tænia mediocanellata		
Hydatid disease		
Bothriocephalus latus		

Trichinosis	}	Considerable eosinophilia, often up to 15 per cent., and sometimes 50 per cent. or more.
Ankylostomiasis		
Bilharzia hæmatobia		
Filariasis		
Asthma during the attacks		
Pemphigus		
Erythema bullosum		
Hydroa herpetiformis		
Herpes gestationis		

Filariasis: Parasites present at certain times.

Trypanosomiasis: Parasites present at certain times.

The Presence of Parasites—Malaria.—Films should be stained by Leishman's method (p. 83), though it is occasionally possible to see the parasites in the fresh blood under the microscope. The corpuscles should be searched as in making a differential leucocyte count, and hæmatozoa cannot be reported absent unless none have been found after half an hour. The best time for finding them is when the films are

made at the beginning of an actual rigor. After quinine has been given none will be present.

The appearance of the parasites will vary with the stage of their development. Between the tertian and the quartan varieties there are but slight microscopic differences. Each begins as a small pale spherule situated excentrically within a red corpuscle. As it grows it expands at the expense of the protoplasm of the red cell, and produces dark pigment granules, which collect near its centre. This is the ring stage, at which it is commonly detected. In the stained film what remains of the affected red corpuscle is pink, and within it there is a bright blue parasite surrounded by an aggregation of almost black pigment granules (Fig. 49). The next stage, that of sporulation, is seldom seen, but not infrequently the parasite, instead of being entirely enclosed within the red cell, is elongated into a crescentic shape (Fig. 51), with pigment granules at its centre, and the ghost of a red corpuscle attached to its concave side. It appears that these crescents are not distinctive of any particular variety of malaria, but that either tertian or quartan parasites may occur in the crescentic form. A flagellate variety is also described, but it is only seen in fresh undried blood that has been allowed to stand outside the body. It is possible that both the flagellate and the crescent forms are really stages in the sexual cycle which the parasite passes through in the mosquito.

Filaria Sanguinis Hominis.—This is only found in persons who have been in the tropics. The mature worm occupies the pelvic lymphatics. Embryos find their way to the blood, and in the common form,

Filaria nocturna, are present in the peripheral circulation at night only. In a rarer form, *Filaria diurna*, they occur in the peripheral blood only in the daytime. In rarer cases still, *Filaria perstans*, they are found both day and night. The embryo (Fig. 50), if present, is readily seen in films stained by Leishman's method.

The Spirillum of Relapsing Fever (Fig. 52).—This occurs in the blood during the febrile attacks, and may be found in films stained by Leishman's method. Each is from 10 μ to 40 μ long, but only 1 μ thick; in fresh blood it is highly motile.

Estimation of the Specific Gravity of the Blood.—Chloroform (specific gravity 1480), benzol (specific gravity 888), a sensitive urinometer graduated from 1020 to 1080, and a glass cylinder deep enough to take the urinometer, are required. Mix chloroform and benzol until the specific gravity of the mixture is 1058, which is that of normal blood. Prick the lobule of the ear; hold a small vessel containing some of the chloroform-benzol mixture under the ear, and allow a drop of blood to fall into it. Transfer this to the main bulk of mixed chloroform and benzol. Should the blood-drop float, the specific gravity of the mixture is higher than that of the blood. Reduce it by adding more benzol. Should the blood-drop sink, raise the specific gravity of the mixture by adding chloroform. When the blood-drop neither sinks nor floats, the specific gravity of the mixture is the same as that of the blood, and may be read off on the urinometer. A special specific gravity apparatus is required if very accurate readings are desired, because the urinometer

graduations are for urines whose surface tension is different to that of a chloroform-benzol mixture.

Importance of Estimating the Specific Gravity of Blood.—The specific gravity rises when the blood becomes concentrated, and therefore affords a means of diagnosis between collapse and shock. Collapse follows great loss of fluid, the blood becomes concentrated, the specific gravity rises; treatment by transfusion is indicated. In shock there has been no loss of fluid, the specific gravity is not raised, transfusion is not indicated.

The specific gravity also affords a valuable check upon hæmoglobin estimations. Specific gravity and hæmoglobin rise and fall together. Approximately—

A sp. gr. of 1035 corresponds to 30 per cent. of hæmoglobin.

„	1040	„	45	„	„
„	1050	„	65	„	„
„	1055	„	75	„	„
„	1060	„	100	„	„

Lipæmia.—In certain cases of diabetes mellitus the blood becomes pale and ‘milky,’ a condition which is in some cases due to fine globules of fat, in others to fine particles of a precipitated proteid circulating in the blood. A yellow colour of the retinal bloodvessels may suggest the onset of ‘lipæmia.’ Microscopic examination of a drop of blood may confirm it. The best method of staining fat droplets is to mix a fresh drop of blood with a drop of Sudan III., a stain which is prepared as follows:

Alcohol (96 per cent.), 100 c.c.
 Sudan III. (Grübler), to saturation.
 Filter and add alcohol (50 per cent.), 75 c.c.
 Filter again.

The fat particles become bright red, whilst nothing else present takes the stain. They indicate the imminence of coma, but in certain cases early recognition has led to the administration of large quantities of carbonate of soda in time to avert the fatal issue. Lipæmia has also been noted in severe alcoholism, in phosphorus-poisoning, and after fractures.

Widal's Reaction for Typhoid Fever.—The 'clumping reaction' is not so much a vital as a physical phenomenon. The test may be made with dead cultures, although, where possible, it is best to send the patient's blood-serum to a bacteriological laboratory for testing with living cultures.

FIG. 53.—Widal Tube.
(Half the natural size.)

To obtain the serum, a glass pipette (Fig. 53), drawn into a capillary tube at either end, is needed. The patient's ear is pricked as in making a blood film; into the little pool of blood which collects one end of the pipette is put, and blood immediately runs into it by capillary attraction. No suction is needed. If necessary, the rate of flow of blood into the pipette may be accelerated by keeping the free end below the level of the ear, but on no account should the blood be allowed to flow farther than half-way along the tube. Now, holding the pipette horizontal, seal off the clear capillary end in a spirit flame; as this end cools the blood will be drawn into the body of the pipette, making it easy to seal off the other end in the same way. In about half an hour, when the blood has

clotted, the tube may be turned vertically with its clear end downwards, so that the serum may run down into it comparatively free from red corpuscles. The tube is in most cases sent to a bacteriologist for the completion of the test, the usual process of which is briefly as follows :

The end of the pipette is broken off, and the serum received into a clean watch-glass. A platinum wire loopful is transferred to a cover-slip, and on to the same cover-slip are put nine separate loopfuls of normal saline. The drops are then mixed, so that the serum dilution is 1 in 10. A loopful of this diluted serum is transferred to another cover-slip, and to it nine loopfuls of salt solution are added. The second serum dilution is 1 in 100. A loopful of broth culture of typhoid bacilli is now put on to each of two clean cover-slips; to the first a loopful of the serum diluted 1 in 10 is added, making a dilution of 1 in 20; to the second, a loopful of the serum diluted 1 in 100, making a dilution of 1 in 200. Each cover-slip is inverted over a hanging-drop chamber, the edge having been vaselined to prevent evaporation, and the bacilli are looked at in the hanging drop under high power. If the test be positive, the bacilli will adhere to one another in batches or 'clumps,' instead of being evenly distributed. The test gives positive proof of typhoid fever if the clumping occur within half an hour with a serum dilution of 1 in 200. If the clumping take place with a dilution of 1 in 20, and not with 1 in 200, typhoid fever may, or may not, be present. The reaction is seldom fully positive until after the end of the first week of the fever.

Examination for Bacteria in Blood.—Circulating blood seldom contains enough bacteria to be detected in films. In most cases it is necessary to obtain enough blood to send to special laboratories for cultivation. The best way to obtain it is direct from a vein in the arm by means of a syringe. The skin must be made surgically clean, but all antiseptic must be removed with boiled water, or else the organisms will be prevented from growing. The arm is prepared as for venesection, but, instead of incising the skin, the needle of the syringe is plunged obliquely into the distended vein. The syringe and needle must be sterile, but free from antiseptic, and to prevent the blood from clotting a few c.c. of 1 per cent. solution of citrate of potash should be present in it to start with. Another method is to collect blood in a series of sterilized Widal tubes from the lobule of the ear (p. 100), after rendering the latter aseptic, but this does not yield so much blood, and contamination is more likely to occur. The more blood obtained, the more likely are organisms in it to grow; about 10 c.c. is a usual amount to take. The kind of cases in which the procedure is required are those of septic infection or suspected fungating endocarditis, when identification of the organism will lead to the use of the proper vaccine or serum in treatment. In typhoid fever it is said that the *Bacillus typhosus* may be found in the blood early in the illness, thus affording a possible means of diagnosis before the Widal reaction is positive.

The Opsonic Index of the Blood.—Opsonins are the substances in the blood-serum which control

the phagocytic power of the leucocytes. There appear to be different opsonins for different bacteria, so that each person has one opsonic index to tubercle bacilli, another to staphylococci, and so on. To determine the opsonic index against any particular organism it is necessary to obtain blood-serum from the patient by the same process as for Widal's test (p. 100), strict asepsis being observed. The specimen is then sent to a bacteriological laboratory, where they prepare—

1. Serum from a normal person.

2. Fresh leucocytes, obtained as follows: Drops of blood from a finger are received into normal saline containing a little citrate of soda. The mixture is centrifugalized, and the supernatant fluid poured off. The mixed red and white corpuscles at the bottom are then shaken up with fresh normal saline, and re-centrifugalized. The supernatant fluid is poured off; the deposit furnishes the leucocytes required.

3. An emulsion of the particular organism against which the opsonic index is to be taken.

The procedure is briefly as follows: Upon each of two capillary glass tubes, fitted with a rubber teat at one end, a mark is made about an inch from the other end. The rubber teat is compressed, and the end of one of the pipettes is put into the washed corpuscles; by slowly relaxing the pressure on the teat, the latter are drawn up until they reach the mark on the tube. A bubble of air is then drawn in, and the end of the pipette dipped into the patient's serum. The latter is drawn up till it reaches the mark, another bubble of air is drawn in, and then emulsion of bacteria is drawn up to the mark in the same way. The tube now con-

tains equal quantities of — (a) washed corpuscles ; (b) patient's serum ; (c) emulsion. The contents are mixed by expelling them on to a slide, reaspirated high up into the capillary tube, and the latter is then sealed above and below. With the other pipette a precisely similar operation is carried out, substituting normal serum for that of the patient. Both sealed tubes are now put into the incubator for fifteen minutes. Blood films are then made from the contents of each tube, and appropriately stained. The leucocytes will be found to have taken into themselves a certain number of bacteria. The numbers so taken in by 50 polymorphonuclear cells in each slide are counted. Suppose in the preparation in which the patient's serum was used 50 leucocytes had taken in 85 bacteria, and that in the other 50 leucocytes had taken in 170, the patient's opsonic index for the organism in question would be $\frac{85}{170}$, or 0.5.

Wassermann's Serum Reaction for Syphilis is of considerable diagnostic value, but the technique is complex, and special laboratory skill is required in carrying it out. There are many modifications of the original method ; special publications on the subject should be consulted. In practice blood is obtained from the patient with aseptic precautions, as described on p. 102, and sent to a bacteriological laboratory for the completion of the test, which is one depending upon what is known as 'fixation of complement.'

Estimation of Calcium Salts in Blood.—Calcium salts have a very important rôle in the coagulability of blood. The possible importance of

calcium in hæmophilia, menorrhagia, scurvy, blood diseases, and so forth, is therefore great. The following is a clinical method of estimating calcium in the blood:

Bell's Method.—The small capsule (Fig. 54 [1]) contains 200 cub. mm. of 1 in 30 aqueous solution of oxalic acid; when about to be used it is broken at A. The patient's ear is pricked, 100 cub. mm. of blood are drawn into the pipette (Fig. 54 [2]), immediately expelled into the oxalic acid, and the capsule sealed. Its contents become laked; ten minutes later they are ready for further dilution, though the next steps may be postponed for hours if necessary. The bulb is broken at B, and 250 cub. mm. of the following reagent added to it:

Acetic acid (1 per cent.) ..	95 parts
Glycerin	5 ..

After thorough shaking, 100 cub. mm. are transferred to a vessel containing 500 cub. mm. of *distilled* water. This makes the final

mixture, a drop of which is transferred to a Thoma-Zeiss hæmocytometer (p. 65), and the number of calcium oxalate crystals counted precisely as in making a blood examination. A large number of squares should be counted. One crystal per square is equiva-

FIG. 54.—Bell's Apparatus for Estimating Calcium in Blood.

lent to 1 in 6,000 CaO. Healthy adults have between 1 in 4,000 and 1 in 5,000 CaO in their blood.

To Estimate the Coagulability of the Blood.—
This requires a special apparatus, termed a 'coagulometer.' Full directions are issued with the instrument.

CHAPTER III

EXAMINATION OF THE SPUTUM

I. NAKED-EYE CHARACTERS.

CERTAIN macroscopic appearances of the sputum are typical.

In Bronchitis it is at first small in amount, frothy, and contains small, clear, sticky pellets of mucus. Later the expectoration, though still frothy upon the surface, has a thick, opaque, yellowish deposit of muco-pus.

In Phthisis with Cavitation nummular sputum is distinctive—heavy airless lumps of muco-pus, each of which, if expectorated on to a dry surface, flattens out into a circular meniscus, which does not coalesce with its neighbours ; hence called ‘nummular,’ or coin-like. If expectorated into carbolic lotion or other fluid, each more or less spherical, airless mass sinks to the bottom, but remains distinct from its neighbours. Streaks of red blood occur in the earlier stages ; later, blood-clots resembling in shape the lung cavities. It must not be forgotten that streaks and clots of blood result also from adenoids, mitral stenosis, fibroid lung, whooping-cough, and severe bronchitis.

In Fibroid Lung with Bronchiectasis the amount of sputum may be great, up to as much as 1 pint or more in the twenty-four hours. It often smells very

foul. Expectoration usually occurs at comparatively long intervals, when a large quantity of sputum is coughed up at a time. Received into a specimen-glass, it settles into layers: upon the surface, frothy serum; at the bottom, a thick sediment of muco-pus; between, clear serous fluid.

In cases where an Empyema has ruptured into a Bronchus the sputum is very similar to that of bronchiectasis.

In Lobar Pneumonia the sputum is termed 'rusty.' This name applies to its usual colour, a dull red-brown. The colour is not constant, for it is due to blood, and may be any hue that a bruise may be, from bright red to dull red, brown, bluish, green, or yellow. The chief characteristic of pneumonic sputum is its viscosity. So glairy is it that, if expectorated into a dry porringer, the latter may be turned upside-down and yet the sputum will only very slowly slide along its surface.

In Plastic Bronchitis, a rare condition, casts of the bronchioles occur. Similar casts may be found in cases of diphtheria in which membranous exudate has extended in the tubes.

Anchovy-sauce-coloured Expectoration is rare, but is almost pathognomonic of hepatic abscess ruptured into the lung.

No other types of sputum can be called characteristic. In infarction of the lung it is said to resemble 'prune-juice,' in growth of the lung 'red-currant jelly,' but neither is constant. Morning expectoration of thick phlegm stained black by smoke and dust occurs in healthy persons who live in towns or smoky atmospheres.

II. MICROSCOPICAL CHARACTERS.

The Sputum in Phthisis.

It is in cases of phthisis, suspected or certain, that investigations of the sputa are so important. The following points should be attended to:

1. The amount. Large quantities indicate cavitation or bronchiectasis.

2. The character. Nummuli indicate cavitation.

3. The presence of blood. Streaks in early phthisis; clots when there is cavitation.

4. The presence of elastic fibres.

5. The presence of tubercle bacilli, and whether few, many, or very many.

6. The presence of other bacteria, particularly staphylococci, streptococci, pneumococci, influenza bacilli, and tetrads.

Elastic Fibres. — To the sputum add excess of 10 per cent. caustic soda, and boil for five minutes.

All the organized elements except the elastic fibres, adventitious hairs, and the more highly resistant organisms, such as tubercle bacilli, are destroyed. Pour the swollen gelatinous residue into 3 inches of water in a specimen-glass, stir well, and allow to settle. If necessary, centrifugalize the deposit. Transfer some of the deposit

FIG. 55.—Elastic Fibres from Lung, the lower less magnified and showing Alveolar Arrangement.

to a slide, cover, and examine under a low power. Elastic fibres (Fig. 55) are sharply defined double-contoured threads, sometimes single, sometimes branched, and occasionally arranged in a meshwork corresponding in outline to a pulmonary air vesicle.

Importance.—They indicate a serious condition of active lung disintegration, such as occurs in advancing phthisis, gangrene, or pulmonary abscess.

Tubercle Bacilli.—Make a film. It is essential that the sample taken for this purpose should be from the interior of a nummulus. It is best picked out by a fine pair of forceps, and it should be smeared fairly thickly over a clean slide and allowed to dry in the air. Disinfect the forceps with heat. In many cases it is sufficient to make a film in this way without previous treatment of the sputum, but where tubercle bacilli are few, they may not be found without first breaking up the nummuli by the caustic-soda method of detecting elastic fibres (p. 109). There are various other methods of concentrating the bacilli when the numbers present are so few that it is difficult to detect them; perhaps one of the best of these is the *Antiformin process*, which is as follows:

To a quantity of sputum in a beaker add sufficient antiformin to make the result a 20 per cent. solution of the latter. Stir thoroughly to break up the pellets of sputum, and then leave to stand for an hour or more until the liquid appears of one consistence throughout. At first there is generally some bubbling from the liberation of gas by the interaction of organic material and the antiformin; the latter destroys the mucus and pus, leaving only the more

resistant substances, including tubercle bacilli, hairs, keratin particles, and cellulose. The resultant fluid has a high density, however, and to reduce this and render it easier to centrifugalize it, an equal quantity of 95 per cent. alcohol is added, the whole being well stirred, and then allowed to sediment until next day; or, better still, centrifugalized to hasten the sedimentation. Films are then prepared in the ordinary way from the deposit so obtained. After such treatment bacilli will adhere better if the slides be albuminized (see p. 21).

Having made and dried the film, fix it by passing it slowly through a Bunsen flame five times, film upwards, then stain by the **Ziehl-Neelsen method**, for which the following reagents are required:

1. *Carbol Fuchsin Stain.*

Saturated alcoholic solution of fuchsin	10 c.c.
Carbolic acid crystals	5 grammes
Distilled water to	100 c.c.
Add absolute alcohol	10 "

2. *Sulphuric Acid* (25 per cent. solution).

3. *Carbol Methylene Blue Stain.*

Methylene blue	1.5 grammes
Carbolic acid crystals	5.0 "
Distilled water to	100.0 c.c.
Add absolute alcohol	10.0 "

Warm some carbol fuchsin in an evaporating basin until it begins to steam freely, and immerse the film in it for five minutes; then wash the slide in tap water, and immerse in 25 per cent. sulphuric acid; the latter changes the red colour to orange yellow, and much of the stain is discharged. In one minute wash

off the acid with tap water. The red colour will reappear. If it be a faint pink, enough stain has been removed; if a darker red, return the film to the sulphuric acid until, on washing with tap water, only a faint pink remains. The tendency is to remove too little stain rather than too much. Now blot with filter-paper and transfer to carbol methylene blue for three minutes; wash in tap water; blot; allow to dry in the air; examine under $\frac{1}{12}$ -inch oil-immersion lens with a good Abbé condenser.

The principle of the above method is as follows: Tubercle bacilli take up stain with difficulty, but once they are stained it is difficult to unstain them. Cold carbol fuchsin will not stain them, warm will; but at the same time all the other structures are greatly overstained. The sulphuric acid corrects this by removing the fuchsin from everything except the tubercle bacilli. Leprosy bacilli and smegma bacilli are the only other important organisms in man which retain the stain in the same way. Carbol methylene blue counterstains the remaining structures, so that tubercle bacilli stand out red in a blue field. Each measures about $3\ \mu$ in length, or less than half the diameter of a red corpuscle, and they are exceedingly thin. They frequently have a beaded appearance, though they are said not to form spores. They may occur singly or in pairs joined end to end (see Plate II., Fig. 59, facing p. 120), or in clusters of half a dozen or more.

Importance.—Tubercle bacilli are never adventitious; their presence means tuberculous trouble. It is not possible, however, to say that tuberculosis is absent

because no organisms have been found. Careful and repeated search may be needed before such negative evidence can be relied on. In acute miliary tuberculosis of the lung they may not be found at all. On the other hand, presence of large numbers is not a hopeless sign. The prognosis may be indicated more certainly by the presence or absence of elastic fibres, or of the organisms of secondary infection, than by the mere numbers of tubercle bacilli; nevertheless, the most certain proof of the diagnosis of phthisis is the discovery of tubercle bacilli in the sputum.

Other Pathogenic Bacteria in Phthisical Sputum—Staphylococci.—Each is about $1\ \mu$ in diameter, and they are arranged in clusters and irregular clumps (Plate II., Fig. 60).

Streptococci.—Each is about $1\ \mu$ in diameter, and they are arranged in short or long strings (Plate II., Fig. 61).

Diplococci.—Each is about $1\ \mu$ in diameter, and they are arranged in pairs, with an unstained capsule round each pair (Plate II., Fig. 63).

Importance.—As a general rule, when many tubercle bacilli are present, there are few other organisms, and *vice versa*. Many of the severe symptoms of phthisis are due to staphylococci and streptococci. When such secondary infection has occurred, cachexia and toxic symptoms increase, pyrexia is more constant, and the prognosis, as a rule, worse. In this respect the presence of streptococci is more serious than that of staphylococci only. Diplococci have no particular significance in such cases; they may, indeed, be found in perfectly healthy persons.

Non-Pathogenic Bacteria in Phthisical Sputum.—Bacteria of various kinds occur ; their names are not important.

Tetrads, or Sarcinæ, consisting of cocci arranged in fours, or multiples of four (Fig. 56). These indicate putrefaction in the cavities in the lung, and are therefore a bad sign, though in themselves harmless. They occur in bronchiectasis as well as in phthisis.

FIG. 56.—Tetrads and Pus Cells.

(High power of microscope.)

Pus Cells and Mucus in Phthisical Sputum—The former are chiefly polymorphonuclear leucocytes, the latter structureless streaks and wisps. Both are stained blue.

Importance.—In the earlier stages of phthisis mucus predominates ; later pus cells may be extremely numerous, and indicate suppurating cavities.

In addition to the above constituents of a film, large squamous cells may occur ; these are derived from the buccal mucosa, and have no pathological significance.

The Sputum in Lobar Pneumonia.

The viscid character of the sputum is almost pathognomonic ; microscopical examination is confirmatory, but not essential. Lobar pneumonia is caused by more organisms than one. Films stained either with carbol methylene blue (p. 111), carbol fuchsin (p. 111), or by Gram's method (p. 117), will show many pus cells, some red corpuscles, streaks of mucus, probably groups of staphylococci and strings of streptococci, and numbers of one or other of the following bacteria :

Diplococcus Pneumoniæ of Fraenkel (Plate II., Fig. 63).—This is the causative agent in most cases. Each organism is a pointed coccus about $1\ \mu$ in longest diameter; they occur in pairs, within a capsule which remains unstained by ordinary methods. They stain with carbol fuchsin and with carbol methylene blue, and retain the stain by Gram's method (p. 119). Their capsule may be stained by the following process:

Capsule Stain.

Dahlia	..	..	..	..	0.5 gramme
Methyl green (100 crystals)	..	..	..	..	1.5 grammes
Saturated alcoholic solution of fuchsin	10				c.c.
Distilled water to	..	..	..	200	,,

Immerse a film in the stain for five minutes; wash gently in distilled water, and dry.

Pneumobacilli of Friedländer.—These are short rods occurring in pairs within a capsule. They stain with carbol fuchsin and with carbol methylene blue, but not by Gram's method. They are rare, and difficult to identify by the microscope alone.

Influenza Bacilli.—These are exceedingly minute rods, $1.5\ \mu$ long by $0.3\ \mu$ broad. They are not arranged in pairs, and have no capsule. They stain with carbol fuchsin and with carbol methylene blue, but not by Gram's method. Cultural methods are needed for their certain identification.

Importance.—Each of these organisms may occur in the sputum of patients who are not suffering from pneumonia; more stress is to be laid upon the viscosity of the sputum than upon the micro-organisms it contains, unless possibly in differentiating influenzal pneumonia from the ordinary form.

Charcot-Leyden Crystals. — These consist of phosphate of spermin, and are visible in fresh sputum

FIG. 57.—Charcot-Leyden Crystals and Pus Cells.

(High power of microscope.)

under the high power as colourless crystals of elongated diamond shape (Fig. 57). They are not pathognomonic of any particular affection, nor are they common. They occur in some cases of asthma, but not in all. They have been found in bronchiectasis and in

chronic phthisis. Since they have also been obtained from seminal fluid, and from leucocythæmic blood, they are regarded as products of cell disintegration.

Curschmann's Spirals.—These may be seen in fresh sputum under the low power, and may even be visible to the naked eye. Each consists of a central core, round which is woven a thick spiral meshwork of delicate fibres (Fig. 58). Apparently they are mucoid casts of the finest bronchioles. They have been found in broncho-pneumonia, in bronchiectasis, in capillary bronchitis, in asthma.

If any significance can be attached to them at all, other than that they indicate affection of the bronchioles, it is in asthma, where their presence may prove that the patient has bronchial asthma as distinct from other forms, such as uræmic or cardiac.

FIG. 58.—Curschmann's Spiral.

(Low power of microscope.)

CHAPTER IV

EXAMINATION OF PUS

THE chief point to be determined is the variety of organism present. Smear out some of the pus in a thin layer upon a clean slide; allow the film to dry spontaneously; when quite dry, fix it by passing it five times through a Bunsen or spirit flame. Stain by one or other of the following methods:

1. **With Carbol Methylene Blue** (p. 111).—Immerse the film in the stain for five minutes; wash gently in water for a few seconds; blot, and dry in the air.

2. **With Carbol Fuchsin** (p. 111).—Immerse the film in the stain for three minutes; wash vigorously in water for thirty seconds; blot, and dry in the air.

This and the preceding method will stain all the organisms described below except the tubercle bacillus. Carbol fuchsin readily overstains; the reverse is true of carbol methylene blue; in each case the carbolic acid acts as a mordant, and is essential.

3. **By Gram's Method.**—The following solutions are required:

(a) *Aniline Water*:

Aniline oil	..	..	..	..	5 c.c.
Water	..	..	..	..	100 "

Shake up together in a bottle ; the water becomes impregnated with the aniline oil, but globules of the latter remain undissolved. The water must therefore be filtered when used.

(b) *Gentian Violet Stain :*

Saturated solution of gentian violet in absolute alcohol.

(c) *Gram's Iodine Solution :*

Iodine	..	..	..	..	1 gramme
Potassium iodide	..	..	..	..	2 grammes
Distilled water	..	..	..	..	300 c.c.

Fill a suitable small vessel three-parts full of filtered aniline water ; add 5 to 10 drops of gentian violet solution ; immerse the film in the mixture for five minutes. Wash off the free stain with water ; transfer the film to Gram's iodine solution ; previously violet, it here becomes a deep slate colour. After one minute remove it from the iodine, and wash with alcohol, which removes much of the colour and makes the film pale grey. Wash in water again ; the film should now become a quite pale violet colour. If the tint be at all deep, insufficient stain has been removed, in which case repeat the iodine process ; then blot off the water with filter-paper, and allow the film to dry in the air.

Some organisms are Gram - positive — that is to say, they retain the gentian violet in spite of treatment with iodine ; others are Gram-negative. The method is therefore of considerable assistance in diagnosis.

Gram-Positive Organisms.

Staphylococci.
 Streptococci.
 Pneumococci.
 Anthrax bacilli.
 Diphtheria bacilli.
 Tetanus bacilli.
 Actinomyces (parts).

Gram-Negative Organisms.

Gonococci.
 Pneumobacilli.
 Bacillus coli communis.
 Bacillus typhosus.
 Bacillus mallei.
 Cholera vibrios.
 Plague bacilli.
 Diplococcus intracellularis.

4. **By the Ziehl-Neelsen Method** for tubercle bacilli (see p. 111).

5. **By the Capsule Stain** for pneumococci (see p. 115).

Examine the stained film with $\frac{1}{12}$ -inch oil-immersion lens, using an Abbé condenser; without the latter, bacteria are difficult to see. There is no need to use Canada balsam and a cover-glass unless the film is to be kept. Occasionally, and particularly in tuberculous pus, no organisms may be present. More often bacteria will be found. Identification can seldom be made with absolute certainty without the cultural methods of a special laboratory. The following, however, are the microscopical characters of some organisms that may be met with* :

Staphylococcus Pyogenes.—Each is spherical, about $1\ \mu$ in diameter; it occurs in irregular clusters or masses. Three varieties are described: (a) **Albus**; (b) **aureus**; (c) **citreus**, according to the colour of the culture on gelatin. Under the microscope all look alike (Plate II., Fig. 60). It is of frequent occurrence in subcutaneous abscesses, sputum from

* It often happens that pus contains non-pathogenic as well as pathogenic bacteria; much experience is needed in distinguishing these. For descriptions of them a larger book upon the subject should be consulted.

lung cavities, acne, boils, and, indeed, most forms of suppuration. It is usually less virulent than other pyogenic organisms, but is also able to produce very serious infections such as acute suppurative periostitis, osteomyelitis, or fatal septicæmia.

Streptococcus Pyogenes.—Each is spherical, about $1\ \mu$ in diameter; it forms chains, which may be long or short, the organism being called in the one case **Streptococcus longus**, in the other **Streptococcus brevis** (Plate II., Fig. 61).

It is a virulent organism in many cases, not so often found as staphylococcus, but generally a more serious infection. It may occur in simple abscesses, phthisical sputum, pyorrhœa alveolaris, empyemata, suppurative peritonitis, erysipelas, fungating endocarditis, osteomyelitis, septic conditions of the uterus, pyæmia, septicæmia.

Gonococcus.—Each is rather larger than a staphylococcus, and is kidney-shaped rather than spherical; they occur in pairs, but have no capsule, and they are found *inside* the pus corpuscles as well as outside (Plate II., Fig. 62).

Its presence in urethral pus may elucidate a doubtful case of urethritis, though it dies out in chronic cases. It may also occur in gonorrhœal ophthalmia, synovitis, abscesses, and infective endocarditis. It is often of great medico-legal importance, in connection with stains on clothes, vaginal discharges in children or young girls, and so forth.

Pneumococcus of Fraenkel.—Each is $1\ \mu$ in diameter, not quite spherical, owing to a sharp, lancet-shaped projection. They are arranged in pairs, with

DESCRIPTION OF PLATE II

Fig. 59.—Tubercle Bacilli and Pus Cells.

„ 60.—Staphylococci.

„ 61.—Streptococci.

„ 62.—Gonococci.

„ 63.—Pneumococci.

„ 64.—Diphtheria Bacilli.

„ 65.—Anthrax Bacilli.

„ 66.—Tetanus Bacilli.

„ 67.—Actinomyces.

„ 68.—Cholera Vibrio.

„ 69.—Plague Bacilli.

„ 70.—Vincent's Angina Organisms.

The magnification is approximately that of the $\frac{1}{12}$ -inch oil-immersion lens.

PLATE II.

Fig. 59.

Fig. 60.

Fig. 61.

Fig. 62.

Fig. 63.

Fig. 64.

Fig. 65.

Fig. 66.

Fig. 67.

Fig. 68.

Fig. 69.

Fig. 70.

the points in opposite directions, and each pair is surrounded by a capsule (Plate II., Fig. 63).

It is the chief cause of lobar pneumonia, in the sputum of which it may be found in abundance. It is, however, sometimes found in healthy persons or in phthisical sputum where there is no pneumonia. It occurs in many cases of empyemata, and more rarely causes peritonitis, arthritis, meningitis, pericarditis, otitis media, laryngitis, infective endocarditis or chronic or acute septicæmia.

Pneumobacillus of Friedländer (see p. 115).

Diplococcus Intracellularis of Weichselbaum, or Meningococcus.—Each is $1\ \mu$ in diameter. They are arranged in pairs, with no capsule, and occur both inside and outside the pus cells. They do not retain the stain by Gram's method, in this respect differing from pneumococci, but resembling gonococci; they differ from the latter in not being kidney-shaped; they give rise to posterior basal and to epidemic cerebro-spinal meningitis, and are present in the pus beneath the pia mater. They may sometimes be found in the cerebro-spinal fluid obtained by lumbar puncture.

Bacillus Coli Communis.—Each is a short rod, measuring $3\ \mu$ by $0.5\ \mu$, with blunt, rounded ends. In the fresh state it is motile by means of flagella, which require special and somewhat difficult methods of staining. It does not form spores. Occurring abundantly in the large intestine, it is there harmless. Escaping to other tissues, it may become very virulent, and is found in many cases of appendicular abscess, general suppurative peritonitis, vulvitis, pyelitis, pyelonephritis and cystitis.

Bacillus Typhosus, or Eberth's Bacillus.—This is indistinguishable from the *Bacillus coli communis* except by special methods of culture. Microscopically the two are identical.

Diphtheria Bacilli, or Klebs-Loeffler Bacilli.—These are slender rods about $3\ \mu$ by $0.4\ \mu$, but varying considerably. They are non-motile, and do not form spores, though they often show beads of lighter and darker staining which resemble spores (Plate II., Fig. 64). They often show a tendency to form short chains, and one end is frequently swollen more than the other, so that a clublike appearance is presented. It is very important in a doubtful case of sore throat to examine for the bacilli. Depress the patient's tongue with a spatula whilst he breathes through his mouth; collect a little of the suspected matter from the tonsil upon a pledget of cotton-wool; make a film from this; fix it by heat, and stain either by Gram's method (p. 117) or, better still, by Neisser's stain :

Neisser's Stain for Diphtheria Bacilli.

Methylene blue (Grübler's)	..	..	1 gramme
Rectified spirit	..	..	20 c.c.
Distilled water	..	..	950 "
Glacial acetic acid	..	..	50 "

Contrast Stain.

Vesuvium	..	..	..	2 grammes
Distilled water	..	..	..	1,000 c.c.

Boil; allow to stand; filter; and store for use.

Immerse the film in Neisser's stain for four seconds; rinse rapidly in distilled water; immerse in the contrast stain for four seconds; rinse in distilled water; blot and dry. Diphtheria bacilli will be

stained brown, with an occasional bright blue granule in their interior. Their presence may thus be established in a few minutes, and under conditions when there may be no facilities for employing cultural methods. It is, nevertheless, always advisable to have a cultivation made at the same time, in order that the diagnosis may be confirmed or otherwise by examination of the culture after twenty-four hours' incubation.

Many cases of diphtheria need no such examination to assist the diagnosis. Many others cannot be diagnosed from septic sore throat or follicular tonsillitis without it. On the other hand, diphtheria bacilli sometimes occur in the throats of people who are seemingly healthy, so that, without clinical signs as well, the mere presence of the organisms does not constitute diphtheria.

Anthrax Bacillus.—Each is a characteristic and comparatively large organism, $7\ \mu$ long and $1.2\ \mu$ broad, non-motile, spore-forming, and many may be joined end to end in a threadlike chain (Plate II., Fig. 65).

Anthrax infections occur mainly in those who work with raw foreign hides or wool. The common lesion is a local 'malignant pustule' on the face or hand. Early detection of the organisms is all-important. Whilst they are local to the pustule, serum treatment, with or without excision, saves the patient. After infection of the blood-stream the disease is beyond control.

Tetanus Bacillus.—Each is a slender rod, $4\ \mu$ long by $0.4\ \mu$ broad. They do not form chains as a

rule, and, when fresh, are very slightly motile. Their chief characteristic is the formation of a large bulging spore at one end, which gives them a 'drum-stick' shape (Plate II., Fig. 66). It is by the clinical signs that the disease is diagnosed. The discovery of the organism has its chief value in determining where the inoculation has been, and, consequently, the part requiring surgical attention. The bacillus is anaërobic, and occurs locally in some punctured or septic wound. Along with it numbers of other organisms are usually present.

Actinomyces.—This is rare in man. It affects least rarely the lower jaw, and very rarely the cæcum, the liver, or the lung. To the naked eye the pus may contain minute granules, not bigger than a small pin's head, usually semi-translucent and greenish-grey. Quite rarely their colour is an orange yellow. Under the high power each granule is seen to have three parts, namely: (a) A central *mycelium* of fine interlacing threads; (b) *cocci*, or small spore-like bodies, intermingled with the mycelium; (c) *clubs*, the swollen, pear-shaped ends of filaments which project radially outwards from the surface, and give the 'rayed' form to the mass (Plate II., Fig. 67). The mycelium and cocci stain both with the ordinary aniline dyes and by Gram's method; the clubs stain with the former, but not by Gram's process.

Notwithstanding the rarity of the affection, its diagnosis from other forms of suppuration is all-important, because potassium iodide may cure it without need for operative measures.

Glanders.—The *Bacillus mallei* is a minute rod,

3 μ by 0.4 μ , with a tendency to form short chains. It is non-motile, does not form spores, and does not stain by Gram's method.

It occurs sometimes in those who have to do with horses. The lesions are in the skin or in the lung, and resemble those of tubercle or those of suppuration. Discovery of a bacillus in the sputum or pus should be followed by cultural identification of the organism.

Cholera Vibrio.—This is found in the rice-water stools. It is a bacillus curved like a comma (Plate II., Fig. 68)—broader at one end than the other, with a flagellum at the narrow end, by means of which it is motile when alive. It is very small, 1.5 μ by 0.5 μ , and there is a tendency for two to join end to end, with their curves opposite ways, like the letter S.

Plague Bacilli.—These are small oval rods with rounded ends which stain deeply, whilst the central part remains unstained, a characteristic to which the term 'pole-staining' has been given (Plate II., Fig. 69). They do not form spores, are non-motile, and occur singly as a rule, though they may form chains. They may be looked for in the affected glands or buboes, or, in pulmonary cases, in the sputum.

For **Tubercle Bacilli**, see p. 110.

For the **Amœba of Dysentery**, see p. 145. This gives rise to hepatic abscess, the pus from which has a characteristic chocolate colour. When obtained from the interior of the abscess, the pus is often sterile; but if scrapings from the abscess wall be taken, the amœba is usually demonstrable. Dysentery itself has more than one causal factor; the variety

due to Shiga's bacillus does not tend to be followed by hepatic abscess.

The Spirochæta Pallida.—This is the causal organism in syphilis. There are many known spirochætæ, but the *S. pallida* is distinct from the others. It is a motile spirillum, about $7\ \mu$ long, and extremely thin, being less than $0.25\ \mu$ across: hence the difficulty in seeing it. It has from six to fourteen abrupt, deep, regular, narrow corkscrew twists. It stains with Leishman's stain (p. 83), or, better, with a modification of this, sold as **Giemsa's stain**, the formula for which is as follows:

Azur II. eosin	..	..	..	3 grammes
Azur II.	..	..	..	8 "
Glycerin (Merck's chemically pure)	..	..	..	250 "
Methyl alcohol	..	..	..	250 "

The film is fixed in absolute alcohol for twenty minutes, and then covered with Giemsa's stain, to 15 c.c. of which 10 c.c. of distilled water have been added; it is left for at least an hour, and may be left all night, then washed well in distilled water and dried. *S. pallida* will now be rose-pink, whereas other spirochætæ would be bluish or purple. It has been found most often in chancres and in secondary lesions, such as condylomata and mucous ulcers; it is less common in tertiary lesions, but has been detected in them and in congenital cases. The surface of the lesion should be cleaned and scraped with a sharp spoon, films being made from the blood-stained scrapings obtained from the deeper parts.

Vincent's Angina.—This is a condition which resembles diphtheria in the appearance of the lesions

and the epidemic character of the complaint, but differs from it in not having its serious sequelæ, and in not requiring the serum treatment. When examining for diphtheria bacilli in these cases, Vincent found two characteristic organisms (Plate II., Fig. 70) —(a) a spirillum, and (b) a fusiform bacillus. The former by itself is not unlike some non-pathogenic spirilla, but the fusiform bacillus is readily recognized; it is a slender, slightly curved bacillus, tapering off to a point at each end. It is stained slowly by carbol methylene blue, so that the film should be immersed for fifteen minutes instead of five. It is also stained by Leishman's method.

CHAPTER V

EXAMINATION OF THE GASTRIC CONTENTS

A. CHEMICAL EXAMINATION.

THE **apparatus** required is—

One glass funnel, 6 inches in diameter ; one glass funnel, 2 inches in diameter ; two glass beakers of 300 c.c. capacity ; one glass flask of 1,000 c.c. capacity ; one burette for decinormal soda solution ; one pipette, 100 c.c. ; one pipette, 10 c.c. ; four small evaporating basins of 50 c.c. capacity ; one small flat porcelain slab ; test-tubes ; fine muslin for straining ; filter-papers for the smaller funnel ; litmus-papers, blue and red ; a Bunsen burner, or spirit lamp ; and iron tripod ; reagent bottles, conveniently 2 or 4 ounce capacity ; one big bottle for standard soda solution ; and one flat saucepan, about 6 inches in diameter and about 2 inches deep, for use as a water-bath.

The following **reagents** are necessary :

Günsburg's Reagent.—A test for free hydrochloric acid :

Phloroglucin	..	..	..	2 grammes
Vanillin	..	..	..	1 gramme
Rectified spirit	..	..	..	100 c.c.

Keep the bottle stoppered and not exposed to bright

light. The solution need not be freshly prepared, but it loses its delicacy as a test in time, so that fresh should be made occasionally.

Congo-red Test-Papers.—A test for free acid, organic and inorganic :

Congo red ..	..	..	..	0.1 gramme
Water ..	..	..	..	100.0 c.c.

Into the solution dip filter-paper, allow the latter to dry, cut it into strips, and store ready for use.

Uffelmann's Reagent.—A test for lactic acid :

Carbolic acid ..	..	..	..	3 grammes
Distilled water ..	..	..	..	100 c.c.

To a little of this solution, about 1 inch in a test-tube, add 1 or 2 drops of liquor ferri perchlor. (B.P.) ; an amethyst blue colour results. This is the complete reagent, but it does not keep, so that the ferric chloride and carbolic acid solutions should be kept in separate bottles, and mixed when required.

Decinormal Soda Solutions for the Estimation of Acidity.—A decinormal solution is such as contains one-tenth part of the molecular weight, in grammes, of a univalent acid or base, dissolved up to 1,000 c.c., usually in water. The cheapest method in the case of soda is to use sticks of caustic soda, but they are difficult to weigh accurately, because they absorb water from the air. The molecular weight of NaHO is $23 + 1 + 16 = 40$, so that 4 grammes of solid caustic soda, dissolved up to 1,000 c.c. in water, makes a decinormal solution.

The best plan is to make the solution as accurately as possible ; then to titrate it with a known solution of

sulphuric acid, and thus determine its actual strength. Provided the strength of the soda solution be accurately known, it matters little whether it be absolutely decinormal or not.

The following is an accurate but more expensive method: Weigh a small test-tube, dried and corked; with a clean knife cut the outside crust from one or more pieces of pure sodium; transfer the bright metal to the test-tube; cork, and weigh again. Deduct the weight of the test-tube and cork to find that of the sodium; transfer the latter to a clean flask containing about 200 c.c. of absolute alcohol. It dissolves with evolution of heat, but with less violence than in water; it forms sodium ethylate, which can then be diluted with distilled water to the calculated volume and kept for use. Every 2·3 grammes of sodium must be diluted up to 1,000 c.c. to make a decinormal solution, of which 1 c.c. corresponds to 0·0023 gramme of sodium, or to 0·004 gramme of NaHO, or to 0·00365 gramme of HCl.

Phenolphthaleïn Solution. — The indicator in estimating acidity:

Phenolphthaleïn	..	..	..	0·2 gramme
Rectified spirit	..	..	..	60·0 c.c.
Distilled water to	..	..	..	100·0 „

The solution is colourless in the presence of even weak acids, and is turned rose-red by alkali.

Discs of Hard-boiled White of Egg for Testing for Pepsin.—Boil an egg for twenty minutes; let it get cold; remove the shell; peel the white from the yolk, cut the former into strips, and punch out discs

with a cork-borer. Put the discs into glycerin in a small bottle ready for use.

The following reagents are not necessary, but may be employed as additional or alternative tests for free hydrochloric acid :

Boas's Reagent.

Resorcin	5 grammes
White sugar	3 "
Rectified spirit	100 c.c.

Tropæolin OO.

Methyl orange	0.2 gramme
Rectified spirit	25.0 c.c.
Distilled water to	100.0 "

Methyl Violet Solution.

Methyl violet	0.05 gramme
Distilled water	100.0 c.c.

The Method of Procedure.—First, *strain the vomit* through fine muslin, using the large funnel and flask. Do not use filter-papers, as these alter the acidity of the filtrate by allowing certain of the acid substances to pass through with greater readiness than others.

Next, *test the filtrate with blue litmus-paper*. The vomit may be alkaline. On the other hand, if blue litmus be turned red, the 'acidity' may be due to one or more of four causes :

(a) Acid salts (acid sodium phosphate).

(b) 'Combined' hydrochloric acid: that is, HCl molecularly combined with the proteid. In normal digestion after a proteid meal, the first part of the HCl secreted combines loosely with the proteid molecules ; during this time salivary digestion of carbohydrates continues, lactic acid is usually found, and it

is not until about an hour later that enough HCl has been secreted to leave a surplus 'free.' From this time onwards more 'free' HCl is present and less lactic, until the HCl reaches 0·2 per cent. and lactic acid is absent.

(c) Organic acids, chiefly lactic, with traces of acetic and butyric.

(d) 'Free' hydrochloric acid.

To Determine the Causal Factors of the Acidity :

Test with a Congo-red Paper.—If the colour of the test-paper be not altered, neither organic acid nor free HCl is present; the acidity is due to acid salts and combined HCl. If the paper be turned blue, both free HCl and lactic acid may be present. If the red be changed to a dirty brown, lactic acid is probably present without free HCl.

To confirm the presence of free HCl, **test with Günsburg's reagent** as follows: Boil some water in the saucepan, as a water-bath. Put 3 drops of Günsburg's fluid and 3 drops of the vomit filtrate into a small evaporating dish; float the latter on the boiling water. Free HCl is indicated by a beautiful rose-red colour appearing, first at the drying margins, and later all over the evaporated residue. The test is easy, but is liable to fail if the vomit be not strained, if more than a few drops of either vomit or reagent be used, or if the water-bath be dispensed with and direct heat employed.

Günsburg's test gives a similar rose-red colour with any free inorganic acid, such as nitric or sulphuric, also with *strong* solutions of organic acids. There is never sufficient organic acid in the vomit to do this, nor are

nitric or sulphuric acids present : hence Günsburg's is one of the best for free HCl in the stomach contents.

Next use **Uffelmann's Test for Lactic Acid**. Fill 1 inch of a test-tube with the reagent, add strained vomit drop by drop until the amethyst blue colour is discharged. Free HCl makes the solution colourless, lactic acid a pale canary yellow. It is the *solution* that is yellow. Frequently, on adding acid vomit, a yellowish cloud of phosphates is formed ; an alkaline vomit gives a pale orange colour due to a precipitate ; in all cases *filter* the mixture. If the *filtrate* is clear pale yellow, lactic acid is present ; if a faint yellow, there is a trace ; if colourless, none.

The additional tests mentioned above are performed as follows :

Boas's reagent is used exactly as Günsburg's. A similar, but fainter, rose-red indicates free HCl. Longer heating is necessary, the rose colour sometimes not appearing for ten minutes after evaporation is complete. Allowing for this, Boas's test is almost as satisfactory as Günsburg's, is cheaper, and the solution keeps good longer.

The Tropæolin Test.—Place a drop of the Tropæolin OO upon a porcelain slab, spread it out in a thin layer, and dry it at 40° C. To do this place the slab close to the foot of the Bunsen burner. When dry, put a single drop of the strained vomit upon the centre of the yellow stain, and continue warming at the foot of the burner as before. A beautiful violet colour appears if free HCl be present. A similar violet is given by *strong* lactic acid, but not with the amount present in a vomit. The ease of this test strongly recommends itself in private practice.

The **Methyl Violet Test** is also very simple. Put 1 drop of methyl violet solution upon a white dish close to 1 drop of the strained vomit. A change from violet to bright greenish-blue at their point of coalescence is regarded by many as proof that free HCl is present. There are, however, conditions in which sufficient lactic acid may be present to produce a similar change.

Summary.—With blue litmus-paper, a red reaction indicates one or more of the following: Acid salts, combined HCl, lactic acid, free HCl.

With Congo red, a blue reaction indicates lactic acid or free HCl. A brown reaction indicates lactic acid, and probably no free HCl. With acid salts or combined HCl, the red does not change.

With Günsburg's reagent, a rose-red colour indicates free HCl.

With Uffelmann's reagent, a yellow colour of the *filtrate* indicates lactic acid.

Boas's reagent (rose-red)	} May be used as
Tropæolin OO (violet)	
Methyl violet (to greenish-blue)	
	alternative to
	Günsburg's test.

Importance of the Tests.—The absence of free HCl from vomit within an hour of proteid food is physiological. Its absence later indicates deficient acid secretion, resulting in interference with peptic activity and a liability to the growth of organisms. Knowledge of the deficiency is an important indication for treatment, the result of which may be gauged by subsequent testings.

In gastric carcinoma free HCl is often absent, a fact to which diagnostic value has been attached; but it may be absent in many other conditions—for example,

in fevers, chronic heart disease, Bright's disease, phthisis, the cachexia of malignant disease elsewhere than in the stomach, cirrhosis of the liver, and, indeed, in almost any debilitating malady. If, however, the patient's malady is known to be gastric, and possibly carcinoma, then the absence of free HCl at the right time after a test-meal is a point in favour of the diagnosis of carcinoma of the stomach.

Presence of lactic acid is not pathological for the first hour after food; longer than this its presence in quantity indicates undue fermentation.

To Estimate the Acidity of the Vomit.

The Total Acidity.—Fill the burette with decinormal soda solution. Pipette 100 c.c. of strained vomit into a beaker. Add 10 drops of phenolphthaleïn solution. Place the beaker upon a white slab under the burette. Add the soda, a few drops at a time, until a definite pink colour appears and persists. It is well to have a second beaker containing colourless acid solution for comparison. The volume of soda solution added gives the amount of caustic soda needed to neutralize the total acids in the 100 c.c. of vomit.

The Acidity due to Free Hydrochloric Acid.—Most methods are too complicated for practical use. The following is simple, and gives approximate results: Prepare a series of small evaporating basins, and into each put 2 drops of Günsburg's reagent. Pipette 100 c.c. of strained vomit into a beaker. Add decinormal caustic soda from the burette; after the addition of 5 c.c., put 1 drop of the beaker contents into the first evaporating basin, and dry the latter on the

water-bath. Add 5 c.c. more soda to the beaker, put 1 drop of the beaker contents into the second basin, and dry it on the water-bath. Repeat this operation with successive evaporating basins after each addition of 5 c.c. decinormal soda. Suppose a deep rose-red results in the first basin, a less deep with the second, a faint with the third, and none at all with the fourth, it follows that between 15 c.c. and 20 c.c. of decinormal soda are needed to neutralize the free HCl in 100 c.c. of vomit. Now pipette 100 c.c. of strained vomit into a fresh beaker; prepare the basins with Günsburg's reagent as before; add 16 c.c. of soda to the beaker, and take 1 drop from the latter for the first basin, and similarly 1 drop for successive basins after 17 c.c., 18 c.c., and 19 c.c. of soda have been added respectively. If the rose-red reaction be faint in the first basin, fainter in the second, just perceptible in the third, and absent in the fourth, it follows that between 18 c.c. and 19 c.c., or 18.5 c.c., of decinormal soda solution just neutralize the free HCl in 100 c.c. of vomit.

If the total acidity be already estimated, the difference between the two gives the acidity due to lactic acid, combined HCl, and acid salts.

Importance of the Estimation.—It is unusual in health to find more than 0.2 per cent. of free HCl. More indicates hyperacidity, which may be corrected by prescribing more proteid food to combine with the HCl, or by giving alkalies to neutralize the acid at that period after a meal when it is present in excess.

When doubt exists in diagnosis between gastric carcinoma and a simple ulcer, hyperacidity favours the latter.

To Test for Pepsin.

Into each of four test-tubes put 5 c.c. of strained vomit and 1 disc of egg-white. To the second add 2 grains of dried pepsin, to the third 2 drops of dilute hydrochloric acid (B.P.), to the fourth 2 drops of dilute hydrochloric acid and 2 grains of pepsin. Warm the saucepan water-bath to 40° C., stand all the test-tubes in it, and leave them at this temperature for two hours. The egg in the fourth tube will have swollen up and been partly dissolved. If it be dissolving in the third, and not in the first, acid is deficient in the vomit, but pepsin is present. If it be dissolving in the second, and not in the first, pepsin is deficient. If it dissolve in the fourth, but not in the first three, both pepsin and acid are deficient.

To Test for Rennin.

Neutralize 10 c.c. of strained vomit by adding 2 drops of phenolphthaleïn as an indicator, and dilute caustic soda until a faint pink colour appears. Add 10 c.c. of fresh cow's milk previously boiled and allowed to cool. If rennin be present, the caseinogen of the milk will be precipitated in flakes.

B. MICROSCOPICAL EXAMINATION.

Mix a little of the deposit from the muslin filter with a drop of salt solution upon a slide, put on a cover-slip, and examine it with a low power, shutting off a good deal of the light with the diaphragm.

Particles of food will be seen. **Fat droplets**, highly refractile, structureless globules, large and

small, staining black with a 1 per cent. solution of osmic acid or bright red with Sudan III. **Starch grains**, round or oval, concentrically striated, staining blue with dilute iodine. **Elastic fibres**, double contoured, sometimes branched and twisted spirally at their free ends. **Muscle fibres**, transversely striated, unmistakable. **Vegetable fibres**, also unmistakable.

Importance.—Starch grains, elastic fibres, muscle fibres, may be found when no food containing any of these has been eaten the same day. They then indicate delayed emptying of the stomach.

Vegetable Parasites.—For these the $\frac{1}{6}$ -inch power is needed. Two sorts may be looked for:

Sarcinæ ventriculi, or 'packet cocci,' each resembling a bale of wool, tightly corded so as to bulge

FIG. 71.—*Sarcinæ ventriculi*.
(Medium power of microscope.)

FIG. 72.—Yeast Cells.

between the cords. There may be four cocci in each packet, or any multiple of four (Fig. 71). Each coccus is about the size of a leucocyte, has a yellow tinge, and with iodine stains mahogany-brown.

Yeast fungi, or *torulæ*, round or oval cells in branching chains or clusters (Fig. 72).

Importance.—Neither is present in health. They indicate chronic delayed emptying of the stomach,

particularly in cases of dilated stomach, whatever the cause.

Test-Meals.

Carbohydrate stimulates the flow of gastric juice less than does proteid, but leads to an earlier appearance of free hydrochloric acid. The more proteid given, the less early is free hydrochloric acid to be expected (see p. 131). Before interpreting the results of an examination, it is essential to know what food has been taken, and at what interval before the stomach contents were obtained. In such cases as require particular investigation, therefore, it is usual to give a 'test-meal' and to recover it by means of an emetic or by the stomach-tube. The test-meals recommended are—

Ewald's Breakfast.

White bread	..	..	..	70 grammes
Weak tea	..	..	..	300 c.c.

Empty the stomach in one hour.

Klemperer's Breakfast.

White bread	..	..	..	70 grammes
Milk	..	..	..	500 c.c.

Empty the stomach in two hours.

Germain Sée's Lunch.

White bread	..	..	..	150 grammes
Minced meat	..	..	..	80 ..
Cold water	..	..	..	300 c.c.

Empty the stomach in two hours.

Riegel's Dinner.

Soup	..	..	..	100 c.c.
Beef-steak	..	..	..	60 grammes
White bread	..	..	..	50 ..

Empty the stomach in five hours.

In practice, Ewald's breakfast is the most convenient.

CHAPTER VI

EXAMINATION OF THE FÆCES

A. GENERAL EXAMINATION.

THE fæces are extremely variable in amount, form, colour, and odour ; but the following types are never healthy :

The 'Pea-soup' Stool.—Copious, unformed, dirty yellow, very ill-smelling. Chiefly seen in typhoid fever, but by no means always present in this disease.

The 'Rice-water' Stool.—Colourless, odourless, alkaline, watery, with numbers of small white flocculi of epithelium and mucus floating in it. It is seen in cholera, but very similar stools may occur in severe arsenical poisoning and in the very severe diarrhœa that occasionally results from cathartic drugs.

The Blood and Mucus Stool.—Small in amount, without fæcal colour, often odourless, consisting of slimy mucus mixed with bright red blood. It occurs in several conditions, notably intussusception, acute colitis, ulcerative colitis, dysentery.

The Clay-coloured Stool.—May be well or ill formed ; usually offensive, but not always so. The colour resembles that of putty, and is due to absence of bile pigments. It indicates obstruction of the large bile-ducts, as by catarrh, gall-stone, or growth. It is

therefore associated with jaundice ; but jaundice may occur without clay-coloured stools when the large bile ducts are free, as in cirrhosis of the liver.

The Fatty, Clay-coloured Stool.—When bile is absent fat digestion is imperfect, but when neither bile nor pancreatic juice reaches the bowel it almost ceases. A copious iridescent scum of fat upon a clay-coloured motion suggests occlusion of both bile and pancreatic ducts, such as results from carcinoma of the head of the pancreas, or from chronic pancreatitis.

The Infantile Zymotic Enteritis Stool.—Very scanty, drying upon the diaper as a grass-green stain of a most offensive odour ; occasionally streaked with blood.

The Purulent Stool.—Obvious pus occurs sometimes, coating the motion, or coming from the bowel as a separate evacuation. It may occur in dysentery or ulcerative colitis, but more often follows rupture of an appendicular or other abscess into the colon or rectum.

The Mucous Colitis Stool.—Consists of mucus in the form of white gelatinous casts of the interior of the bowel. Under the microscope the casts are transparent and structureless ; embedded in them are fragments of undigested food, leucocytes, scattered epithelial cells, blood-corpuscles, micro-organisms, and crystals of phosphates and of cholesterin. They have every variety of length, from an inch up to some feet ; are often narrow and riband-like, suggesting tape-worms, but distinguished at once by their not being segmented. They are diagnostic of mucous

colitis, a most troublesome disorder, occurring in women of nervous temperament. It is common for such patients to pass *intestinal sand* also from time to time. The 'sand' may amount to a drachm or more at an evacuation. It consists of small granules of inorganic salts, especially phosphate, carbonate, and sulphate of calcium, together with a certain amount of organic matter. The sand often has a reddish-brown colour.

The Stool with Blood in it.—The character of blood in a motion varies with the site of the bleeding-point. Bright streaks of blood upon the outside of a stool indicate hæmorrhage near the rectum—*e.g.*, from piles, or sigmoid or rectal carcinoma or other stricture. Bright blood mixed with the fæces comes from higher up—*e.g.*, from tuberculous, typhoid, or other ulceration of the ileum or cæcum. Blood and mucus occur with intussusception or inflammation or ulceration of the colon. When bleeding is high up—*e.g.*, from a gastric or duodenal ulcer, or from ruptured varicose veins at the lower end of the œsophagus in cases of cirrhosis of the liver—or when blood has been swallowed and passed on, the motions are black and tarry, a condition called 'melæna.' Black motions may also be due to—

1. Certain drugs.

Iron.	} Converted by the sulphuretted
Manganese.	
Bismuth.	
	hydrogen in the bowel into
	sulphides.

2. Certain foods, notably bilberries and charcoal biscuits.

To confirm the presence of blood, a little of the motion may be smeared upon a piece of lint, a drop of tincture of guaiacum placed by the side of it, and ozonic ether poured over both. If blood be present, a bright blue colour appears. The guaiacum reaction, however, has many fallacies (p. 129), and a more conclusive test is with the spectroscope. Shake up a little of the fæces in a test-tube with water acidulated with acetic acid, allow the solid particles to settle, and examine the supernatant fluid for the absorption bands of hæmatin. In some cases, especially of malignant disease of the alimentary canal, the amount of blood present may be so small as to require special extraction before it can be detected. When testing for this 'occult blood,' a quantity of fæces is thoroughly stirred up and mixed with about one-third its volume of glacial acetic acid; an equal volume of ether is then added, and the whole shaken up repeatedly to dissolve the acid hæmatin in the ether; on standing a little while the ether with the dissolved pigment floats to the top; it may be pipetted off and tested for blood either by the guaiacum test or spectroscopically.

The Stool with Undigested Milk in it.—The curds may be obvious to the naked eye. Whenever patients who are upon milk diet suffer from diarrhœa—*e.g.*, in typhoid fever—it is important to examine the fæces before ordering astringents. Should curds be present, modification of the milk with sodium citrate, lime, or barley water may cure the diarrhœa at once.

B. SPECIAL EXAMINATIONS.

Gall-Stones.—Mix the fæces with water, and pour them on to a fine-meshed sieve beneath a running tap; triturate well with a stick.

FIG. 73.—Cholesterin Crystals.
(High power.)

A gall-stone, if present, will become visible after the softer fæcal matter has passed through. It may be an obvious gall-stone to the naked eye; if there is any doubt, test it as follows:

1. **For Cholesterin.**—

Dissolve a portion of the stone in chloroform; add sulphuric acid. If cholesterin be present, the solution assumes a blood-red colour, going on to purple.

Or, extract a part of the stone with ether; evaporate the solution; dissolve the residue in absolute alcohol, and allow to crystallize. Examine for the characteristic crystals of cholesterin (Fig. 73) under the microscope.

2. **For Bile Pigments.**—Dissolve a portion of the stone in weak hydrochloric acid. Apply Gmelin's test (p. 42) to the solution.

Importance.—Discovery of a stone in the fæces after an attack of biliary colic is the best proof that the stone has been got rid of. Should the stone be faceted, other gall-stones are, or have been, present.

Typhoid Sloughs.—These may be looked for on a sieve in the same way as gall-stones.

Importance.—Less now than formerly. Previous to

the discovery of the Widal test, the diagnosis of typhoid fever often depended on finding sloughs from Peyer's patches in the stools.

Bacteria.—These swarm in fæces, and are for the most part unimportant. To examine them, pick out a very small portion of the excretion with forceps, make and fix a film, and stain by the methods described on p. 117, *et seq.* The organisms of importance are—

The Cholera Vibrio (p. 125).

The Amœbæ of Dysentery.—These can only be found in stools less than twelve hours old ; later they disintegrate. They are best obtained from the smaller

FIG. 74.—Amœbæ of Dysentery.
(High power of microscope.)

pellets of mucus. Each is a rounded or irregular protoplasmic mass about $30\ \mu$ in diameter, finely granular, refractile, with several vacuoles, and ingested red corpuscles or bacteria (Fig. 74). In warm 0.6 per cent. salt solution they may be watched sending out blunt pseudopodia, and gradually crawling along the slide. In the fresh state they show no nucleus, but stained specimens show one about the size of a red corpuscle. The amœbæ disintegrate with heat, so that films cannot be fixed in the ordinary way ; they

should be immersed for five minutes in the following solution :

Gulland's Fixing Solution.

Absolute alcohol	..	..	..	25	c.c.
Pure ether	..	..	..	25	„
Solution of corrosive sublimate in absolute alcohol (strength, 2 grammes in 10 c.c.)	..			0.5	„

and then stained with carbol methylene blue or with carbol fuchsin (p. 117). Another method of demonstrating them is to stain with carbol thionin blue for ten minutes ; wash the excess of stain off with distilled water ; rapidly rinse in distilled water containing 1 per cent. of acetic acid ; wash again in distilled water ; blot, dry, and examine. The amœbæ stain a deeper purple than other cells that may be present.

The amœbic form of dysentery is only one of several ; it is the kind most liable to be followed by hepatic abscess.

The **Typhoid Bacillus** is so similar to the *Bacillus coli communis* (p. 121) that the two cannot be distinguished without elaborate cultural methods. The same applies to Gaertner's bacillus, to ptomaine poisoning, to Shiga's dysentery bacillus in paratyphoid bacilli, and to the bacteria of zymotic diarrhœa.

Intestinal Parasites.

These are of two classes : I. Cestodes, or tapeworms. II. Nematodes, or threadworms.

I. **Cestodes.** — These have two life - cycles : a sexual, the hermaphrodite tapeworm, in one host ; and an asexual, cystic, in another. There are three

whose tapeworm stage occurs in the human small intestine—namely :

Tænia Solium.—Its intermediate host is the pig ; the head of the worm (Fig. 75) has four round sucking-discs and a single row of hooklets, about twenty-six in number.

Tænia Mediocanellata.—Its intermediate host is the ox ; the head of the worm (Fig. 76) has four round sucking-discs, and no hooklets.

Bothriocephalus Latus.—Its intermediate host is

FIG. 75.—Head of
Tænia Solium.

FIG. 76.—Head of
Tænia Mediocanellata.

FIG. 77.—Head
of *Bothriocephalus Latus*.

(Semi-diagrammatic ; medium power of microscope.)

a fish, such as the pike ; its head (Fig. 77) has two longitudinal sucking-discs, and no hooklets.

These worms vary from 10 to 20 feet in length, and are composed of hundreds of opaque white segments joined end to end. Each segment is a complete hermaphrodite organism, containing an ovary, a testis, and a water-channel system. At the upper end of the worm is a narrow neck, and a head not much larger than that of an ordinary pin. After the administration of anthelmintics, diligent search should be made

for this head, for there is no certainty of the eradication of the worm until the head itself has been passed. It is by the characteristics of the head that the variety of tapeworm is diagnosed.

The head of *Tænia echinococcus* has four round sucking-discs and a ring of hooklets like *Tænia solium* ; but the worm is only $\frac{1}{4}$ inch long, consisting of a head, a neck, an immature and a mature segment ; it inhabits the intestine of the dog, and it is only the cystic stage which affects man, causing hydatids.

II. **Nematodes.**—These have the sexes distinct.

Ascaris Lumbricoides.—This is 6 to 8 inches long ; is not unlike an earthworm, but is distinguished by the following points : (1) There is no imbrication of segments, so that the finger can be drawn along it either way without catching ; (2) the mouth is at the extremity instead of under a blunt prostoma, as in the earthworm ; (3) it tapers towards each end ; (4) it has no cingulum or fleshy band, such as surrounds the earthworm at the junction of the anterior and middle thirds.

Oxyuris Vermicularis.—This is $\frac{1}{3}$ -inch long, blunt at the head end, tapering to a pointed tail (Fig. 78), and appearing in the fæces like white, slowly moving

FIG. 78.—*Oxyuris Vermicularis*.
(Low power of microscope.)

short threads. It inhabits the large intestine, and is common in children. Enormous numbers may be passed by the same patient.

Ankylostomum Duodenale.—This inhabits the duodenum. Not uncommon in the East, it is rare in England, though occasionally there are epidemics like that amongst the Cornish miners. Should ankylostomiasis be suspected, thymol should be administered,

FIG. 79.—*Ankylostomum Duodenale*.

A. Much-magnified head.

B. Much-magnified tail of male.

C. Entire parasite, less magnified.

and the fæces examined for the worms. Each is $\frac{1}{4}$ inch long, greyish-white in colour, cylindrical; the posterior end is broad, and in the male is provided with an umbrella-like expansion with eleven ribs. The worm tapers forward to a narrow neck, which ends in a bulging mouth, the margin of which is furnished with

four characteristic clawlike hooks and two blunt teeth (Fig. 79), which enable the parasite to adhere firmly to the duodenal wall. The symptoms of ankylostomiasis are so similar to those of pernicious anæmia that cases are easily overlooked. Eosinophilia may suggest the diagnosis, which is confirmed by giving thymol and finding the parasite in the fæces.

Trichocephalus Dispar.—This is about $1\frac{1}{2}$ inches long, and consists of an opaque white body often wound up like a watch-spring, and a long threadlike anterior extremity (Fig. 80). It inhabits the cæcum.

FIG. 80.—*Trichocephalus Dispar*.
(Natural size.)

Though not uncommon, it causes no symptoms. It is acquired by ingestion of food

containing ova derived from the dejecta of other persons who have the worm.

Various vegetable substances may readily simulate intestinal parasites; after bananas in particular the patient may think he is passing tapeworm segments. Pips of oranges, lemons, melons, or vegetable marrows have frequently caused unnecessary alarm. Celery fibres may simulate *Oxyuris vermicularis*, and so may fragments of onions, dates, oatmeal, rhubarb, or pineapple. The error can be avoided at once by testing the pseudo-parasite for cellulose which, after weak sulphuric acid has acted on it for a little while, gives the blue starch reaction with iodine.

The Ova of Intestinal Parasites.—These, when abundant, may be found in simple smears of the

fæces. More often it is necessary to concentrate them. The following is an easy way of doing this: Put as much fæces as would lie on a shilling into a test-tube; fill the latter three-parts full of normal salt solution (0·6 per cent.). Cork it, and shake the contents violently for some seconds. Now allow the tube to stand for a quarter of an hour. The heavier portions of the fæces, including the ova, will have subsided to the bottom. Pour off the dirty supernatant liquid, and replace it with fresh normal saline. Shake violently again, to disintegrate as many of the larger fæcal fragments as possible. Allow to stand, and repeat the process until the normal saline becomes practically clear after standing for a quarter of an hour. Now pour off the surface fluid, transfer a drop of the sediment to a slide with a pipette, cover the drop, and examine the preparation under a medium power, using a mechanical stage so as to insure every part being looked at.

A kaleidoscopic conglomeration of all sorts of fragments and cells will be seen, amongst which striated muscle fibres, connective tissue, epithelial cells, leucocytes, spiral vessels from plants, vegetable cells, carbonaceous particles, phosphatic crystals, hair stumps, pieces of keratin, bacteria, and débris will occur. Ova, if present, attract attention at once. Before experience is gained all sorts of things may be taken for them, but once seen they cannot be mistaken. They are all of comparatively large size, of perfectly even outline, with a definite capsule surrounding an indefinite embryo, and they are mostly stained a deep brown by the fæcal pigment stercobilin. They can easily be

seen with the low power, and they are quite large under the $\frac{1}{6}$ -inch objective.

The most characteristic of all the ova is that of *Trichocephalus dispar* (Fig. 81), each being an elongated

FIG. 81.—Ovum of *Trichocephalus Dispar*.

FIG. 82.—Ovum of *Tænia Solium*.

(Both are magnified to the size as seen under $\frac{1}{8}$ -inch objective.)

oval with clear rounded bulges at the ends. The eggs of the *Tænia solium* and *T. mediocanellata* are much alike, each being almost round, with a very thick envelope marked with highly characteristic radiating

FIG. 83.—Ovum of *Ascaris Lumbricoides*.

(Slightly more magnified than are Figs. 81 and 82.)

lines (Fig. 82). *Oxyuris vermicularis* ova seldom occur in the stools; the worm leaves the body to lay them. The egg of *Ascaris lumbricoides* (Fig. 82) is a broad oval, and consists of unsegmented protoplasm with a clear, thick capsule, and a characteristic brown uneven

envelope outside this. *Bothriocephalus latus* ova have no radiating lines in the capsule like those of *Tæniæ*. If the eggs of *Ascaris lumbricoides* had the outermost envelope removed, and the internal protoplasm coarsely segmented into a mulberry-like mass, it would be like the egg of *Bothriocephalus latus*.

CHAPTER VII

MICROSCOPICAL EXAMINATION IN AFFECTIONS OF THE SKIN

Ringworm.—Though many varieties have been described, differing from one another in the cultural characters of the fungi, it is sufficient for most practical purposes to make the following two classes :

1. **Tinea tonsurans**, ringworm of the scalp, due to a vegetable parasite, which forms a mycelium inside the hairs, produces small spores, and is called the **Trichophyton microsporon endothrix**.

2. **Tinea circinata**, ringworm of the body, due to a vegetable parasite, which forms a mycelium outside the hairs, produces large spores, and is called the **Trichophyton megalosporon ectothrix**.

There is a third parasite described, with large spores and its mycelium inside the hair—the **Trichophyton megalosporon endothrix**. Apparently, this is much rarer than the other two.

The essential things to look for are the mycelium and the spores. With forceps pick out one of the stubbly hairs from the suspected patch, and proceed by one or other of the following methods :

(a) **To Detect the Mycelium.**—Immerse the hair

in ether for fifteen minutes, or for twenty-four hours if there be time, in order to remove the grease; transfer to liquor potassæ for fifteen minutes, wash in water, and place in a drop of glycerin upon a slide; apply a cover-slip; press it down; blot up the excess of glycerin; ring the specimen with Canada balsam, and examine under the $\frac{2}{3}$ -inch objective. The mycelium may be seen within or around the hair.

(b) **To Stain the Spores.**—Immerse the hair in ether to remove the fat as before; then stain by Gram's method (p. 117), and mount in glycerin; or dehydrate with absolute alcohol, clear with xylol, and mount in Canada balsam. Examine under the $\frac{1}{6}$ -inch

FIG. 84.—Spores on Hair
in *Tinea Tonsurans*.

FIG. 85.—Spores on Hair
in *Tinea Circinata*.

power. The spores will be deep violet; in the case of *tinea tonsurans* they form a closely packed casing to the hair, resembling a mosaic (Fig. 84); in *tinea circinata*, on the other hand, the large spores are less densely packed, and form rows which have been compared to rosaries (Fig. 85).

Tinea Versicolor.—With the back of a scalpel scrape the surface of one of the brown patches of affected skin; transfer the fine scrapings to a drop of

liquor potassæ on a slide; apply a cover-slip, and examine under $\frac{1}{6}$ -inch objective. Amongst clumps of epithelial débris will be seen the **Microsporon furfur**, interlacing jointed threads of colourless mycelium with bunches of spores between them.

Favus.—Pick out a hair and as small a portion of the surrounding crust as possible. Transfer it to a drop of liquor potassæ on a slide, apply a cover-glass, press it down, and examine with a $\frac{2}{3}$ -inch objective. A fungus, the **Achorion Schoenleinii**, will be seen as a dense branching mycelium, completely infiltrating the hair, and forming a dense coating of large spores upon its surface. It closely resembles a trichophyton. The affection is rare in England. A typical case is easy of recognition, but if the primary affection be obscured by impetigo, diagnosis without the microscope may be difficult.

Scabies. — The parasite is called **Sarcoptes hominis**, or **Acarus scabiei**, and it is related to the spider. There are two sexes. The male lives upon the surface of the skin, is the smaller, and is very difficult to find. The female, when impregnated, penetrates the stratum corneum, and burrows along in the rete Malpighii, depositing eggs as she goes. Her point of entry is marked by a vesicle, and her path by a typical black 'run.' To obtain the parasite, lay a pin flat upon the surface of the skin, and push it with a rotatory movement into the epidermis at that end of the burrow which is away from the vesicle, taking care not to draw blood. The acarus appears as a small pearly object at the end of the pin. Under

the microscope it is oval, with eight legs, two in front and two behind on each side, with a mouth between

FIG. 86.—Ventral View of Female *Sarcoptes Hominis*.

(Low power of microscope.)

the front legs, and a few short hairs at the tail end (Fig. 86).

CHAPTER VIII

EXAMINATION OF SEROUS EXUDATIONS, CEREBRO-SPINAL AND CYSTIC FLUIDS

Serous Fluids from the Peritoneal or Pleural Cavities.—These are alkaline in reaction, contain 1 per cent. of salts, are usually of a straw-yellow colour, and are either quite clear or slightly turbid. The points to examine are—

The Specific Gravity.—This may be taken in a specimen-glass with an ordinary urinometer. It varies from 1005 to 1035.

The Amount of Albumin.—This may be estimated by Esbach's albuminometer, the fluid being suitably diluted with water if necessary (p. 2). It varies from 2 to 40, but more commonly lies between 10 and 30, parts per 1,000.

Importance.—From the specific gravity of the fluid and the amount of albumin it contains it is sometimes possible to decide whether an effusion is inflammatory, or whether it is a passive dropsy such as occurs with cardiac failure. The specific gravity and the amount of albumin are higher in the former than in the latter. Definite conclusions, however, can only be drawn when the figures are extreme: that is to say, a specific

gravity of 1020 or more, or 30 parts per 1,000 of albumin, would indicate an inflammatory origin; a specific gravity of 1010 or less, or 5 parts per 1,000 of albumin, would favour a diagnosis of passive dropsy; figures between these would be compatible with either.

The specific gravity is no measure of the amount of albumin present. Ascitic fluid of low specific gravity may contain much albumin.

The Presence or Absence of Blood.—This may be looked for by the guaiacum test (p. 29), or by the microscope or spectroscope (p. 31).

Importance.—It is common to find blood in the later portions of a tapping, or in a serous effusion that has been tapped before; but if the fluid contain blood at the first tapping, it either indicates a very acute inflammation, or rouses a suspicion of malignant disease.

Cell Elements. — These are best examined by allowing the fluid to stand an hour or more in a specimen-glass, pipetting off and centrifugalizing the lowest part, preparing and staining a film of the deposit, and examining it under the high power. Instead of films being made, it is often more convenient to examine a drop of the centrifugalized deposit, tinged with methylene blue, in precisely the same way as in the case of a urinary deposit (p. 5). There may be no cells at all, or there may be leucocytes, squamous pleural or peritoneal cells, red corpuscles, or even particles of malignant growth. It is rare to find bacteria, owing to the great bulk of fluid in which they are suspended.

Importance.—The presence of many cells, particularly leucocytes, is almost proof of the inflammatory origin of the effusion. It is said that preponderance of lymphocytes indicates tubercle. The detection of a particle of new growth clinches the diagnosis of the cause of the ascites.

Spontaneous Coagulation.—This may occur in both inflammatory and in passive effusions on standing. Should the floating jelly-like coagulum be large and appear quickly, the fluid is probably inflammatory, but otherwise no deduction can be drawn.

The Presence of Fat Globules.—In rare cases the fluid is almost like milk, and the microscope reveals droplets which stain black with osmic acid, or red with Sudan III. (p. 99); the condition is termed *chylous ascites*, of which one chief cause is infection with filaria. It may also result from rupture of the receptaculum chyli after abdominal injury or obstruction to the thoracic duct; or in association with subacute nephritis. Chylous effusion in the chest is very rare, but has occurred after injury to the thoracic duct in the neck.

Cerebro-Spinal Fluid.—This is clear, colourless, alkaline, with a specific gravity which may be as low as 1002. It contains 1 per cent. of salts, a mere trace of albumin, and a body—pyrocatechin—which reduces Fehling's solution on boiling, but does not ferment with yeast, nor yield glucosazone crystals with phenylhydrazine. After head injury, escape from the ear or nose of a clear fluid which reduces Fehling's solution is strong evidence that the base of the skull has been fractured.

The fluid may also be obtained by lumbar puncture for diagnostic purposes, the following being some of its commoner variations from the normal :

In **acute meningitis** it drips out faster than the normal drop per second rate ; it is more or less opalescent or turbid ; it contains more than a trace of coagulable proteid ; its centrifugalized deposit, instead of being nearly free from cells, shows numerous leucocytes, those of the small lymphocyte type preponderating in the tuberculous cases ; whilst polymorphonuclear cells are most numerous in suppurative meningitis. Meningococci may be found on direct staining of films made from the centrifugalized deposit ; whilst cultural or inoculation methods may serve to detect tubercle bacilli, streptococci, staphylococci, pneumococci, or influenza bacilli, according to the type of case.

In **chronic meningitis** the changes are less in degree than they are in acute cases ; in association with syphilis of the nervous system—general paralysis or tabes dorsalis—the cerebro-spinal fluid may exhibit considerable numbers of small lymphocytes in the centrifugalized deposit, even though the fluid itself may appear to be quite clear ; whilst Wassermann's reaction (p. 104) is applicable to it in the same way that it is to blood serum.

Uræmia is often associated with considerable excess of urea in the cerebro-spinal fluid ; it may be tested for by the sodium hypobromite method (p. 57).

Pancreatic Cyst Fluid.—This may be either clear or turbid. It is of watery consistence, alkaline, with 1 per cent. of salts, a specific gravity of about 1005, and little or no coagulable proteid. The essential

characteristic is that it contains proteolytic and amylolytic ferments. These may be tested for as follows:

1. **Proteolytic Ferment.**—Put a disc of hard-boiled white of egg into a test-tube, add some of the fluid, and stand in a water-bath at 37° C. for an hour. Should pancreatic ferment be present, the egg-white will have become partially digested.

2. **Amylolytic Ferment.**—Boil some water in a test-tube, and carefully stir in some finely powdered starch until a thin mucilage is obtained. Cool to 37° C., and add some of the fluid. From time to time take a drop of the liquid and test it with a drop of iodine. If amylolytic ferment be present, the starch will rapidly cease to go blue with iodine, but will presently go red instead, and, finally, will remain colourless, from the formation, in the first instance, of erythrodextrin, and later of achroödextrin. The ferment will ultimately convert the latter into dextrose, which may be detected by its reduction of Fehling's solution.

Importance.—The exact nature of a cyst in the upper part of the abdomen may be obscure even after laparotomy. Its pancreatic origin is proved beyond doubt if the above ferments are detected in the fluid from it.

FIG. 87.—Hydatid Hooklets.

(High power of microscope.)

Hydatid Cyst Fluid.—This is colourless, clear or opalescent, alkaline, and has a specific gravity of about 1010. It contains but a trace of albumin, and by this may be distinguished from ascitic fluid. The main diagnostic point, however, is the presence in it of the 'hooklets' (Fig. 87)

of dead scolices. It must be borne in mind that a hydatid cyst may grow to a great size, with production of daughter and granddaughter cysts, and yet may never have produced scolices. The fluid from such a 'barren' cyst will contain no hooklets.

Hydrocele Fluid.—Its characters are similar to those of ascitic fluid, and almost as variable. A spermatocoele is distinguished from a hydrocele by the presence of spermatozoa in the former. They may be detected in a drop of the centrifugalized deposit from the fluid (Fig. 17, p. 20). It is not uncommon to find cholesterin crystals (Fig. 73, p. 144) in hydrocele and spermatocoele fluids.

Ovarian Cyst Fluids.—There are two types :

1. Ovarian cysts contain a viscous, brownish-green fluid, which pours almost like treacle. It has a high specific gravity (*e.g.*, 1030), alkaline reaction, and 1 per cent. of salts. Albumin is present in variable quantity ; the viscosity is due to a substance called **pseudomucin** or **metalbumin**. It differs from mucin in not being precipitated by acetic acid, and from albumin in not being coagulable by heat. On the addition of alcohol it forms a dense stringy precipitate, and after boiling with HCl it yields a substance which readily reduces Fehling's solution. Microscopical examination of the deposit usually shows much detritus, along with red corpuscles and leucocytes, large epithelial cells, which are sometimes in clumps, cholesterin crystals, and large circular, highly refractile colloid granules, which are said to be characteristic.

2. Parovarian cysts contain watery, colourless fluid, alkaline, of low specific gravity (*e.g.*, 1006), with 1 per

cent. of salts. Albumin is scanty, pseudo-mucin is absent, and cells are very few.

Importance.—The nature of the fluid may be the only means of determining in what structure the cyst arose. The prognosis is more serious in the case of an ovarian than in that of a parovarian cyst.

CHAPTER IX

THE PRESERVATION AND STAINING OF MORBID ANATOMY SPECIMENS

MORBID anatomy specimens may be preserved in one or other of three ways, namely :

- I. In bulk, without regard to keeping the natural colour.
- II. In bulk, with retention of the natural colour.
- III. For microscopical purposes.

I. Preservation in Bulk Without Colour.—The simplest method is to use either (*a*) rectified spirit, or (*b*) 5 per cent. solution of formalin in ordinary water. Care should be taken that the shape of the tissue is correct before it is immersed in the fixing fluid ; cavities should be gently stuffed out with cotton-wool, and there should be wool in the bottom of the vessel to prevent deformity from the weight of the tissue. Much fluid should be used to start with, and it should be changed once or twice subsequently. The formalin penetrates more rapidly than the spirit. Both cause shrinkage of cells, so that if microscopical sections are to be cut it is well to fix a small portion specially for this purpose (p. 166). Hardening will be complete in a week or two. The specimen can then be kept in

just as much of the fluid as will cover it, sealed up in a suitable wide-necked jar.

II. Preservation of Tissues for Colour.—The colour of most tissues is due to hæmoglobin or its derivatives, to preserve which special methods are required. One of the modifications of **Kaiserling's Method** is best. The three following solutions are needed :

1. Pure formalin	..	..	1,600 c.c.
Potassium acetate	..	..	240 grammes
Potassium nitrate	..	..	120 ,,
Water	..	..	8,000 c.c.
2. Alcohol	..	..	80 per cent. solution
3. Distilled water	..	..	4,000 c.c.
Glycerin	..	..	4,000 ,,

The tissue should not be exposed either to the air or to any moisture longer than can be avoided before it is immersed in a large bulk of the first fluid, which should be changed next day, and the tissue left in it for about a week, or, in the case of large specimens, such as livers or brains, for two or even three weeks. It is then of a dull dark colour. Transfer it to the 80 per cent. alcohol, and leave it until the natural colour of the tissue reappears—this should be in from one to six hours. Transfer it to the third fluid, in which, after one or more changes, as may be necessary, it is finally mounted. It is well to keep the preparation unexposed to any bright light.

III. Hardening and Fixing for Microscopical Purposes.—Spirit causes so much shrinking and deformity of the cells that it is unwise to use it. For most ordinary purposes the best fixing fluid is a saturated solution of **corrosive sublimate** :

Mercuric chloride ..	..	..	6.5 grammes
Water ..	..	..	100 C.C.

If normal saline (0.6 per cent.) be used instead of plain water, rather more mercuric chloride is required for saturation, and the results are slightly better. A piece of tissue, about $\frac{1}{2}$ inch in diameter and $\frac{1}{3}$ inch thick, should be immersed in this for twenty-four hours; if left longer it becomes too hard and brittle. It should now be soaked in water for twelve hours, and afterwards carried through a series of bottles containing successive strengths of alcohol—25 per cent., 50 per cent., 75 per cent., 90 per cent., and absolute—being left for twelve hours in each before transference to the next. After twelve hours in the last, it should be put into a second bottle of absolute alcohol for another twelve hours, to insure absolute dehydration. If it is intended to keep the tissue for a time before cutting sections, it is best to stop at the 75 per cent. alcohol stage, at which the tissue can be kept indefinitely without getting too hard.

Embedding in Paraffin.—For actual embedding and section the tissue will usually be sent to a special laboratory. The process of embedding in paraffin is, briefly, as follows :

From absolute alcohol the tissue is put into three changes of xylol, in each of which it stays twelve hours. If still opaque, it is not quite dehydrated, and should be returned to absolute alcohol. If quite clear, it is taken from the xylol, the excess of which is blotted off, and put into paraffin wax kept just melted on a water bath. The wax should be such as has a melting point of 48° C. In twelve hours the tissue is transferred to

fresh paraffin, and again into fresh for another twelve hours. The xylol by this time is all evaporated, and its place in the tissue taken by paraffin. A special mould is now got ready, filled with melted paraffin, and into it, with that surface downwards which it is desired to cut, the tissue is put with a pair of forceps warmed sufficiently to prevent solidification of the paraffin on them. As it cools, the paraffin in the mould sets, a process which is completed by immersing the whole in cold water. The block is then taken from the mould, the excess of paraffin removed with a sharp knife, and the embedded tissue is ready for fixing to the microtome and cutting. One of the best instruments for the purpose is the Cambridge Improved Rocking Microtome.

Mounting and Staining the Sections.—The paraffin sections should be floated out on water which is just warm enough to make them go flat. A microscope slide is partly immersed in the water, and held obliquely under a section, which is then drawn into position by means of a needle. The slide is removed from the water with the section on it, and after the excess of water has drained off, it is placed to dry on a water-bath whose temperature is just less than the melting-point of the paraffin. In about twelve hours it is ready for staining as follows: Pour on a few drops of xylol to dissolve away the paraffin; remove the xylol with absolute alcohol; wash in water to remove the alcohol; pour on a few drops of Gram's iodine solution (p. 118), and leave for a minute or two. The object of this is to remove the last traces of corrosive sublimate by converting it into the more soluble iodide.

If the iodine is not used, the final preparation is very apt to contain numbers of nail-like crystals of sublimate; wash off the iodine with spirit, and immerse the section in **Ehrlich's acid hæmatoxylin**, the formula of which is as follows:

Hæmatoxylin	..	..	..	2 grammes
Glacial acetic acid	..	..	..	10 C.C.
Absolute alcohol	..	..	..	100 „
Glycerin	..	..	..	100 „
Water	..	..	..	100 „
Alum to saturation.				

The stain is best when it has been made up a month or more. It should be filtered from time to time; and, as it stains very deeply, it is wise to dilute it with twice its bulk of fluid made up as above, but without the hæmatoxylin. After staining for five minutes, wash the preparation well in gently running tap-water. The section, previously red, will be turned blue by the alkali in the latter. When quite blue, immerse for not more than thirty seconds in aqueous eosin (p. 82); remove excess of the latter quickly with spirit; remove the spirit with absolute alcohol to dehydrate the section completely; immerse in xylol, and, if the section does not become transparent throughout, return to absolute alcohol; drain off the excess of xylol, and mount in Canada balsam. Cell nuclei are stained blue, cell protoplasm and connective tissue red.

The Freezing Microtome.—This has the advantage of producing sections quickly, and is used when immediate examination is required—for example, during operations, when it is not certain whether a tumour is malignant or not. The principle is

to freeze the tissue in gum by means of an ether spray. Full directions are issued with the instrument.

Embedding in Celloidin.—Some tissues, notably eyes and brain matter, are unsuitable for embedding in paraffin; celloidin is used instead. The tissue is carried through to the absolute alcohol stage, as described on p. 167. Then, instead of xylol, it is put into a mixture of equal parts of absolute alcohol and ether for six hours, and transferred first to a thin and then to a thick solution of celloidin, in each of which it remains a week. It is then mounted in thick celloidin upon the microtome block-carrier, and placed in a desiccator until firm enough not to pit on pressure. The block is further hardened in 80 per cent. spirit for a couple of days, after which sections are cut upon a special microtome.

Müller-Formol Fixing Fluid.—This is used in preference to corrosive sublimate for brain and spinal cord tissues. Its formula is as follows :

Müller's fluid	{	Sodium sulphate ..	25 grammes
		Potassium bichromate ..	25 "
		Water	1,000 c.c.
		Formalin	50 "

The formalin should be added to the Müller's fluid immediately before use. Large pieces of tissue require to be in it for a week or more, during which the fluid should be changed at intervals. Small pieces are fixed in about a day.

Decalcification of Bony Tissues.—It is not possible to cut sections of bony tissues until they have been decalcified. They should first be fixed

like other tissues, after which the mineral salts are gradually removed by immersion in repeated changes of weak nitric or hydrochloric acid. The following is a simple **decalcifying fluid** :

Alcohol, 75 per cent. solution	..	..	100 c.c.
Non-fuming strong nitric acid	..	..	2 „

The process takes a month or more, after which the tissue is dehydrated and embedded in paraffin.

The Stain for Amyloid (Lardaceous) Tissues.
—The paraffin section, after the water stage following the iodine has been reached (p. 168), is immersed for ten minutes in a saturated solution of methyl violet in distilled water. It is then well washed, first in water, then in 1 per cent. acetic acid, and then in water again. It cannot be dehydrated, because spirit removes all the stain. It should therefore be mounted in glycerin, the cover-slip being ringed with Canada balsam. The lardaceous material stains bright pink, the other tissues violet.

CHAPTER X

EXAMINATION OF SEMINAL STAINS

It is sometimes a matter of the very greatest moment, especially from a medico-legal point of view, to be able to determine whether a given stain is seminal or not. The only positive proof is the discovery of spermatozoa (Fig. 17, p. 20) in the stain. In the case of fresh material this is easy, but when the cloth has dried it is sometimes a difficult matter, because the heads of the spermatozoa so readily break off from the tails. A good plan is to excise the suspected spot and to fix both cloth and stain together in Müller's fluid (p. 170) for twenty-four hours, and thus lessen the liability of the spermatozoa to break into fragments.

After fixing, the cloth can be washed well with clean normal saline solution to remove superficial dirt; a suitable portion should then be excised, laid flat on a microscope slide, and firmly and steadily scraped from end to end with the edge of a scalpel, first on one surface, then on the other; the scrapings can be washed off the scalpel on to the slide by raising the cloth in the fingers, squeezing it steadily until a drop exudes, this drop being allowed to run down the edge of the

scalpel on to the slide. A film can be made with this in the ordinary way, fixed in the flame, stained for a minute with aqueous eosin (p. 82), washed gently with distilled water, dried by blotting, and examined under the oil-immersion lens.

CHAPTER XI

TESTS FOR THE COMMONER POISONS

Alkaloids.—Before testing for the following alkaloids they must be extracted from the gastric contents. One of the simpler processes for this is as follows :

Otto's Method.—Acidify the vomit with weak sulphuric acid to convert the alkaloid into the sulphate. Strain through muslin. Agitate the filtrate with ether to remove all fat. Pour off the ether; render the residual watery solution alkaline with potassium carbonate to liberate the alkaloid. Shake up with ether again; the sulphate of the alkaloid was insoluble in ether, but the alkaloid itself is soluble. Concentrate the ethereal solution, and test it as follows :

Belladonna.—Add a little fuming nitric acid; dry upon a water-bath; cool, and moisten with a drop of potash dissolved in absolute alcohol. Atropin will give a *violet* colour changing to red.

Opium.—Evaporate the ethereal solution to dryness; dissolve in cold strong sulphuric acid, and add a drop of strong solution of bichromate of potash. Morphia gives a bright *green* colour.

Strychnine.—Evaporate the ethereal solution to dryness; to the residue add 2 drops of sulphuric acid, and then 2 drops of strong bichromate of potash solution. Strychnine will give a beautiful *purple-blue* colour changing to crimson, and finally to a light red.

Arsenic—I. *Marsh's Test.*—Put a piece of pure zinc into a flask (Fig. 88, B), cork, and through the thistle funnel C pour a little distilled water. Add pure

FIG. 88.—Apparatus required in performing Marsh's Test for Arsenic.

sulphuric acid until there is brisk evolution of hydrogen. Collect a little of the gas escaping from the side-tube A in a test-tube, and ignite it. If it burn in the test-tube without explosion, all the air has been expelled from B. Ignite the hydrogen jet A. Hold a cold porcelain dish in the flame. No stain should be deposited on the white surface if the hydrogen be pure. Now pour in some of the suspected stomach

contents through C, and presently hold the cold porcelain dish in the flame A again. If arsenic be present it will be coming off as arseniuretted hydrogen, and will be deposited as a shining black metallic stain on the white dish.

Antimony will give a similar black stain, but it lacks the metallic lustre.

2. *Reinsch's Test*.—Strongly acidify the stomach contents with hydrochloric acid, and allow to stand one hour. Filter. Introduce into the filtrate a brightly polished strip of copper foil and boil. If arsenic be present, a steel-grey coating will appear upon the copper. This may be tested further by dissolving in nitric acid and evaporating to dryness. On adding silver nitrate to the residue a brick-red precipitate of arsenate of silver will appear.

The difficulty of the test is to insure both the acid and the copper being free from arsenic to begin with.

When arsenic has been taken by a patient over a long period, the fact can be determined by applying the tests for arsenic to the hair that grew during that time. The arsenic becomes stored in the hair and in the nails, which ordinarily contain only minute quantities of the metal.

Carbolic Acid.—Just acidify the gastric contents with hydrochloric acid. Filter. To the filtrate add bromine water. Carbolic acid gives a dense yellow precipitate of tribromophenol.

Chloral.—This is not easy of detection, but may be tested for as follows: Shake the suspected fluid up with pure ether, which will dissolve the chloral hydrate; separate off the ethereal solution, add silver

nitrate and ammonia, and boil. If chloral is present in any quantity, a mirror of metallic silver should form.

Coal-Gas.—The patient's blood will be a bright cherry-red colour. The spectroscope will give the bands of carboxyhæmoglobin identical with those of oxyhæmoglobin (p. 31), but not altering to that of reduced hæmoglobin on adding ammonium sulphide and shaking up well.

Copper Salts.—With a platinum wire hold a little of the suspected fluid in a Bunsen burner or spirit flame. Copper gives a bright green colour to the flame. Add HCl, and evaporate the stomach contents to small bulk. Filter. Add potassium ferrocyanide to the filtrate. Copper will give a characteristic chocolate precipitate.

Corrosive Sublimate.—Mix the gastric contents with an equal bulk of ether, and shake up well. The ether will dissolve the corrosive sublimate. Pour off the ethereal solution; concentrate it by evaporation; add iodide of potassium. Mercury will give a scarlet precipitate of iodide of mercury, soluble in excess of potassium iodide.

Hydrochloric Acid.—Filter the acid stomach contents. Add nitric acid to the filtrate, and then silver nitrate solution. A dense white precipitate of silver chloride results. Normal gastric juice contains hydrochloric acid, which will give the same test. A control must be made with a 0.2 per cent. hydrochloric acid solution, and the density of the two precipitates compared.

Lead Salts (see p. 50).

Nitric Acid.—Neutralize the gastric contents with carbonate of potash. Filter. Evaporate the filtrate to small bulk. Mix with an equal bulk of strong sulphuric acid; allow to cool *completely*. Pour into a test-tube, and carefully add a solution of ferrous sulphate. The latter fluid floats upon the former, and the appearance of a black ring at the junction of the two indicates nitric acid.

Oxalic Acid.—Boil the gastric contents with caustic potash. Even if lime has already been given to the patient, potassium oxalate will be formed. Filter. To the filtrate add solution of calcium sulphate. Oxalic acid will give a dense white precipitate, insoluble in acetic, but soluble in hydrochloric, acid.

Phosphorus.—Dilute the vomit with twice its bulk of water, acidify with sulphuric acid, transfer to a glass retort with a long condensing tube, and distil in the dark. Minute traces of phosphorus will render the condensing vapour luminous.

Prussic Acid.—This may be evident by the smell. To test for it, exactly neutralize the gastric contents, whether they be acid or alkaline, transfer them to a flask, and distil slowly upon a water-bath. To the distillate add silver nitrate; prussic acid will give a curdy white precipitate, insoluble in cold nitric acid, but soluble in hot. To confirm, add liquor potassæ and hydrochloric acid to the precipitate, and then a few drops of sulphate of iron solution. If prussic acid be present, Prussian blue will be formed.

Sulphuric Acid.—Filter the acid stomach contents. Add a solution of barium chloride to the

filtrate. Sulphuric acid gives a dense white precipitate of barium sulphate, insoluble in hydrochloric or nitric acids. The test would naturally be valueless had any sulphate, such as that of zinc, been given by the mouth.

Veronal.—This may be recovered from the urine. The latter is shaken up with several changes of ether; the ethereal extracts evaporated to dryness on a water-bath; the residue dissolved in boiling water, the resulting solution decolourized by filtering through animal charcoal, and concentrated on the water-bath; on cooling veronal crystallizes out.

INDEX

- ABDOMINAL tumours, albumin-
uria with, 29
- Abscess, appendicular, 47, 91,
141
formation, differential leuco-
cyte count with, 94
leucocytosis in, 95
polymorphonuclear cells
in, 91, 95
hepatic. See Hepatic abscess
pulmonary, sputum from, 110
- Abscesses, albuminuria with, 110
micro-organisms in, 119, 120,
121
- Acarus scabiei, 156
- Acetanilide, Fehling's solution,
reduced after, 35
polycythæmia from, 71
- Acetic acid, effect on carbonates
in urine, 27
effects on mucus, 13
on phosphates, 13,
27, 32
on pus, 13
on urates, 13
in heat test for albumin,
27
nuclei brought out by,
17, 33
nucleo - proteid precipi-
tated by, 48
- Aceto-acetic acid. See Diacetic
acid
- Acetone in urine, significance of,
39, 40, 41
tests for, 39, 40
- Achorion Schœnleinii, 156
- Acidity of vomit, estimation of, 135
- Acidity of urine, cause of, 3
- Acidosis, acetonuria in, 41
ammonia in urine, a measure
of, 64
in diabetes, 63
- Acne, staphylococci in, 120
- Acroödextrin, 162
- Actinomyces, Gram-positive, 119,
124
- Actinomycosis, parts affected by,
124
potassium iodide in, 124
- Acute rheumatism, urine highly
acid in, 3
yellow atrophy, leucin and
tyrosin in, 11
- Adenoids, hæmoptysis from, 107
- Albumin distinguished from
nucleo-proteid, 27
fallacy in Fehling's test, 35
in testing for indican, 45
heat test for, 26, 27
effervescence with, 27
Heller's nitric acid test for,
25
magnesium sulphate and
nitric acid test for, 28
methods of estimating, 53, 54
nitric acid test for, fallacies
of, 26
picric acid test for, 28
potassium - ferrocyanide test
for, 28
precipitated by nitric acid,
13
salicyl-sulphonic acid test for,
28
- Albuminized slides, 21

- Albuminometer, Esbach's, 53
 Albuminuria in bacilluria, 17
 causes and significance of, 28, 29
 in pneumonia, 47
 with low blood-pressure, 29
 with pyuria, 29, 33
 Albumose, reaction to salicyl-sulphonic acid, 28
 ring with nitric acid from, 26
 Albumosuria, Bence-Jones's type, 27, 47
 significance of, 47, 48
 test for, 47, 48
 with sarcoma, 27
 Alcohol and ether, films fixed by, 79
 dehydration by, 167
 Alcoholism, glycosuria in, 39
 lipæmia in, 100
 Alkaline tide, 3
 effect on urine, 1
 Alkaloids, tests for, 174
 Alkapton, Fehling's solution reduced by, 35, 49
 ferric-chloride reaction with, 49
 greenish-black urine from, 4
 Alkaptonuria, darkening with potash, 37
 significance of, 49, 50
 simulating bile in urine, 42, 49
 test for, 49
 Altitudes, polycythæmia at high, 71, 96
 Amann, co-efficient of, 45
 Ammonia, estimation of, 63, 64
 in estimating acidosis, 64
 Ammonio-magnesian phosphate, 9, 11
 Ammonium oxalate test for lead, 50
 sulphate test for lead, 50
 urate, 10, 11
 Amœba of dysentery, 125, 145
 fixing fluid for, 145
 in hepatic abscess, 125
 Amphoteric reaction, 3
 Amyl nitrite, Fehling's solution reduced after, 35
 Amyloid tissue, staining for, 171
 Amyolytic ferment, test for, 162
 Anæmia, albuminuria in, 29
 pernicious. See Pernicious anæmia
 Anasarca, polyuria after, 1
 Aniline dye, green urine from, 4
 water, 117
 Ankylostomiasis, colour index in, 77
 eosinophilia with, 92, 96, 150
 simulating pernicious anæmia, 150
 Ankylostomum duodenale, 149
 Anthrax from hides, 123
 bacilli, characters of, 123
 Gram-positive, 119
 serum treatment, 123
 Antifebrin in urine, 51
 Antiformin process of concentrating tubercle bacilli, 110
 Antimony, test for, 176
 Antipyrin, Fehling's solution reduced after, 35
 in urine, ferric chloride test for, 51
 red urine from, 51
 Anuria, causes of, 1
 Appendicitis, albumosuria with, 47
 bacillus coli in, 121
 empyema from, 47
 leucocytosis with, 73
 simulated by coli bacilluria, 22
 Appendicular abscess, polymorphonuclear cells increased in, 91
 purulent stools from, 141
 Arbutin, Fehling's solution reduced after, 52
 ferric chloride, reaction with, 52
 greenish-black urine from, 4
 Arsenic, rice-water stools from, 140
 storage in hair and nails, 176
 tests for, 175, 176
 Arthritis, pneumococci in, 121
 Ascaris lumbricoides, 148
 ovum of, 152
 Ascites, albuminuria with, 29

- Ascites, chylous, 24, 160
 lymphocytes in tuberculous, 160
- Ascitic fluid, characters of, 158
- Asparagus, urine smell after, 4
- Asthma, Charcot-Leyden crystals in, 116
 Curschmann's spirals in, 116
 eosinophilia with, 92, 96
- Atropin, test for, 174
- Bacilluria, albuminuria in (and see Coli-bacilluria), 17
- Bacillus anthracis, 119, 123
 coli communis, 146
 characters of, 121
 Gram-positive, 119
 in urine, 4, 17, 21, 22
 lesions due to, 121
- diphtheriæ. See Diphtheria
- bacilli
- dysenteriæ of Shiga, 126, 146
- fusiformis, 127
- influenzæ, 109, 115, 160
- lepræ, 112
- mallei, 124
 Gram-positive, 119
- paratyphosus, 146
- pestis, 125
- tetani, 123
- tuberculosis. See Tubercle
- bacilli
- typhosus, 119, 122, 145
 Gram-positive, 119
 in urine, 21, 22
- Bacteria in blood, 102
 in stools, 145
- Balsams, nitric acid ring from, 51
- Bananas, intestinal parasites simulated after, 150
- Banti's disease, blood changes in, 94
- Barium chloride, precipitation of rhubarb pigment in urine by, 51
 test for sulphates in urine, 44
 for sulphuric acid, 178
- Basophile cells, 90, 93
- Belladonna, test for, 174
- Bell's method of estimating calcium, 105
- Bence-Jones's albumosuria, 27, 47
- Benzol and chloroform, specific gravity method, 98
- Beta-oxybutyric acid, 41
- Bilberries, black motions after, 142
- Bile pigments in urine, colour of, 4
 other conditions simulating, 42, 49
 tests for, 42
 salts, tests for, 42, 43
- Bilharzia eggs in urine, 24
 embryos, 24
 eosinophilia from, 92
 geographical distribution, 24
 hæmaturia from, 24
- Biliary catarrh, stools with, 140
- Bismuth, black motions from, 142
 subnitrate in Böttger's test, 36
- Biuret test for albumose, 48
- Black motions, 142
- Blackwater fever, 31
- Bladder cells in cystitis, 15
 simulated by renal cells, 16
 growth, particles in urine, 16
 irritability from oxalates, 8
 from uric acid, 9
 tuberculosis, pyuria with, 22
- Bleaching powder, test for indican, 45
- Blood and mucus in stools, 140
 bacteria in, 102
 changes in Banti's disease, 94
 in cachexia, 77
 in chlorosis, 95
 in congenital heart disease, 96
 in cyanosis, 71, 96
 in emphysema, 96
 in fungating endocarditis, 77, 95
 in Hodgkin's disease, 93
 in leukæmia, 91, 92, 93, 94
 in malaria, 73, 91, 95, 96, 97
 in mitral stenosis, 96
 in pernicious anæmia, 87, 88, 95
 in splenic anæmia, 94
 cherry-red, 177

- Blood, coagulability of, 106
 cultures, 102
 diseases, importance of calcium in, 105
 renal tube-casts in, 19
 estimation of calcium in, 105
 examinations, 65
 filaria parasites in, 96
 films, 78
 staining of, 80, 86
 in serous effusions, 159
 in stools, 142
 tests for, 143
 malaria parasites in, 96, 97
 occult, 143
 precipitated in urine, 5
 pressure, low, albuminuria with, 29
 specific gravity of, 98
 tests for, 29, 30
 trypanosomes in, 96
- Boas's reagent, 131
 test, 133
- Boils, staphylococci in, 120
- Bone, decalcification of, 170
- Boric acid in preserving urine, 21
- Bothriocephalus latus, 147
 ovum of, 153
 eosinophilia with, 96
- Böttger's solution, reduced by glycuronic acid, 37, 41
 test for glucose, 36
 fallacies of, 37
- Brain, embedding in celloidin, 170
- Bright's disease. See Nephritis
- Bromine test for carbolic acid, 176
 for carboloria, 49
 for melanuria, 49
- Bromoform, greenish-black urine from, 4
- Bronchiectasis, Curschmann's spirals in, 116
 Charcot-Leyden crystals in, 116
 sarcinæ in, 114
 sputum in, 107
 tetrads in, 114
- Bronchitis, chronic, blood changes in, 96
 Curschmann's spirals in, 116
- Bronchitis, hæmoptysis in, 107
 sarcinæ in, 114
 sputum in, 107
 in plastic, 108
- Broncho - pneumonia, Curschmann's spirals in, 116
- Buckthorn, greenish-black urine from, 4
- Butyl chloral, glycuronic acid from, 42
- Cachexia, colour index in, 77
 differential leucocyte count with, 94
 leucocytes in, 93, 95
 lymphocytes in, 91, 95
- Cæcum, actinomycosis of, 124
 trichocephalus in, 150
- Calcium carbonate in intestinal sand, 142
 in blood, estimation by Bell's method, 104, 105
 oxalate, envelope crystals, 7
 phosphate, 10, 13
 sulphate in intestinal sand, 142
 test for oxalic acid, 178
- Calculus from cystin, 12
 from uric acid, 9
 mulberry, 8
 oxalate crystals with, 8
 red corpuscles with, 8
 renal cells in urine in, 15
 tailed renal pelvic cells in, 16
- Camphor, glycuronic acid from, 42
 reduction of Fehling's solution after, 35
- Capsule stain, 115
- Carbol fuchsin stain, 111, 117
 methylene blue stain, 111, 117
- Carbolic acid, test for, 176
- Carboloria, darkening with potash, 37
 Fehling's solution reduced in, 52
 ferric chloride reaction in, 41
 52
 greenish-black urine from, 4
 reduction of Fehling's solution in, 35, 52
 significance of, 49 50

- Carboluria, simulating bile in urine, 42, 49
 hæmaturia, 49
 tests for, 49
 Carbonates, cloud with heat test, 27
 effect of acetic acid on, 27
 Carboxyhæmoglobin, 177
 Carcinoma of bowel, blood in stools from, 142
 of pancreas, stools with, 141
 of stomach, gastric HCl in, 134
 simulated by pernicious anæmia, 77
 Cascara sagrada, greenish-black urine from, 4
 Casts in sputum, 108
 intestinal, 141
 in urine, 19
 Catheterization, anuria after, 1
 Cauliflower, oxaluria after, 7
 Caustic soda in detecting elastic fibres, 109
 Cayenne-pepper deposit of uric acid, 5, 8
 Celery fibres, simulating oxyuris, 150
 Celloidin, embedding in, 170
 Cells, renal and vesical, 15, 16
 Cellulose, test for, 150
 Centrifugalizing machine, 5
 Cerebro-spinal fluid, alterations in disease, 161
 Fehling's solution reduced by, 161
 from ear, 161
 micro-organisms in, 160, 161
 normal character of, 160
 urea in, 161
 Wassermann reaction in, 161
 Cestodes, 146
 Chancre, spirochæta in, 126
 Charcoal, black stools after, 142
 Charcot-Leyden crystals, 116
 Children, gonococci in, 120
 leucocytes in, 93
 oxyuris in, 148
 Chloral, glycuronic acid from, 42, 51
 Chloral, reduction of Fehling's solution by, 35, 51
 test for, 176
 Chlorides, estimation of, 60, 62
 in urine, reduced in various conditions, 44
 test for, 43
 normal amounts excreted, 62
 Chloroform and benzol, specific gravity method, 98
 glycuronic acid from, 42, 51
 reduction of Fehling's solution by, 35, 51
 Chlorosis, blood changes in, 95
 colour index in, 77
 hæmoglobin in, 76
 Cholera, rice-water stools in, 125, 140
 vibrio, 125
 Gram-negative, 119
 Cholesterol crystals, 144
 in gall-stones, 144
 in hydrocele fluid, 163
 in ovarian cysts, 163
 in spermatocele fluid, 163
 Chromate test for lead, 50
 Chrysophanic acid in urine, 3, 52
 Chylous ascites, 24, 160
 Chyluria, causes of, 24
 Cigarette-paper method of making blood-film, 78, 79
 Cirrhosis of the liver, blood in stools from, 142
 gastric HCl in, 135
 lymphocytes in, 91
 stools in, 141
 urobilin in, 46
 Citrate of potash in preventing blood-clot, 102
 Clay-coloured stools, 140
 Coagulability of blood, estimation of, 106
 Coagulometer, 106
 Coal-gas poisoning, 177
 Coefficient of Amann, 45
 Coffin-lid crystals, 9
 Coli bacilluria, characters of urine in, 17
 nocturnal enuresis from, 22
 pyuria with, 17, 33

- Coli bacilluria, simulating appendicitis, 22
 Colitis, blood and mucus stools with, 140, 142
 indicanuria from, 46
 purulent stools with, 141
 Collapse, specific gravity of blood in, 99
 Colour index, 76, 77
 preservation in tissues, 166
 Condylomata, spirochæta in, 126
 Congenital heart disease, polycythæmia in, 71, 96
 syphilis, lymphocytes in, 90
 spirochæta in, 126
 Congo red test-papers, 129, 132
 Constipation, indicanuria from, 46
 Copaiba, nitric acid ring from, 26, 51
 Copper sulphate in Fehling's solution, 34
 in Haines's fluid, 34
 in Pavy's solution, 54
 in Trommer's test, 36
 test for, 177
 for arsenic, 176
 Cornish miners, ankylostomiasis in, 149
 Corrosive sublimate crystals, 169
 for fixing tissues, 166
 test for, 177
 Creatinin, inhibition of Fehling's test by, 36
 in sodium nitro-prusside test, 40
 no reduction of Böttger's test by, 37
 picric acid test and, 37
 reduction of Fehling's solution by, 35
 Creosote, Fehling's solution reduced after, 52
 ferric chloride reaction with, 52
 greenish-black urine from, 4
 Cryoscopy, 63
 Crystals, ammonio-magnesian phosphate, 9
 Charcot-Leyden, 116
 coffin-lid, 9
 dumb-bell, 8
 envelope, 7
 knife-rester, 9
 of calcium oxalate, 7
 of cholesterin, 144
 of corrosive sublimate, 169
 of cystin, 11, 12
 of uric acid, 8
 phenyl-glucosazone, 38
 phenyl-lactosazone, 38
 thorn-apple, 10, 11
 tyrosin, 11
 Cubebs, nitric acid ring from, 51
 Curschmann's spirals, 116
 Cultures from blood, 102
 Cyanosis, blood changes in, 96
 polycythæmia from, 71
 Cylindroids, mucous, 25
 Cystin, calculi from, 12
 crystals, 11, 12
 familial incidence, 12
 Cystitis, ammoniacal urine in, 4
 ammonio-magnesium phosphate in, 9
 bacillus coli in, 121
 bladder-cells in, 15
 pyuria from, 17, 22, 23
 thorn-apple crystals with, 11
 Dates, oxyuris simulated after, 150
 Decalcification of bone, 170
 Decinormal soda solution, 129
 Dehydration by alcohol, 167
 Delafield's hæmatoxylin and eosin stain, 82
 Diabetes insipidus, low specific gravity of urine in, 2
 pallor of urine in, 3
 polyuria in, 1
 mellitus, acetonuria in, 39, 41, 63
 glycosuria in, 39
 high specific gravity of urine in, 2
 lipæmia in, 99
 pallor of urine in, 3
 polyuria in, 1
 smell of urine in, 4
 yeast fungi in urine in, 23
 Diacetic acid associated in glycosuria, 39

- Diacetic acid in urine, significance of, 41
 test for, 41
 Diarrhoea, milk in stools in, 143
 urate deposit from, 13
 Diazo reaction, 46, 47
 Differential leucocyte count, 88
 93, 94
 Dilated stomach, sarcinae in, 139
 Diphtheria bacilli, 122
 Gram-positive, 119, 122
 Neisser's stain for, 122
 casts in sputum in, 108
 Vincent's angina simulating, 126
 Diplococcus intracellularis, 119, 121, 161
 Dog, taenia echinococcus in, 148
 Drinking, effect on urine, 1, 2, 3
 Drug reactions in urine, 51, 52
 Drugs, black motions from, 142
 as fallacies in nitric-acid albumin test, 26
 causing reduction of Fehling's solution, 35
 chrysophanic acid from, 52
 glycuronic acid derived from, 42
 rice-water stool from, 140
 Drum-stick bacilli, 124
 Dumb-bell crystals of uric acid, 8
 Duodenal ulcer, blood in stools from, 142
 Dusting powder, starch in urine from, 5, 14
 Dysentery, amœba of, 125, 145
 blood and mucus stool with, 140
 purulent stools in, 141
 Shiga's bacillus in, 126
 Dyspepsia, oxalates from, 7

 Ear, cerebro-spinal fluid from, 161
 Earthy phosphates, 10, 13
 Eberth's bacillus. See *Bacillus typhosus*
 Eczema, no eosinophilia with, 92
 Effervescence in heat test for albumin, 27
 Egg-white in pepsin testing, 130
 Egypt, bilharzia in, 24
 Ehrlich's diazo-reaction, 46
 Ehrlich's hæmatoxylin, 169
 triple stain, 85
 Elastic fibres in sputum, 109, 110, 113
 Embedding of tissues, 167, 170
 Embryos of bilharzia, 24
 Empyema, appendicular abscess with, 47
 pneumococci in, 121
 polymorphonuclear cells increased in, 91
 ruptured into bronchus, sputum from, 108
 streptococci in, 120
 Emphysema, blood changes in, 96
 Endocarditis, fungating. See Fungating endocarditis
 Enteritis, indicanuria from, 46
 zymotic, stools in, 141, 146
 Enuresis, from coli bacilluria, 22
 Envelope crystals, 7
 Eosin, alcohol solution, 84
 aqueous solution, 82
 in urine, 3
 Eosinophile cells, 89
 myelocytes, 90
 Eosinophilia in ankylostomiasis, 92, 96, 150
 various causes of, 92, 96
 Epilepsy, urine changes in, 1, 2, 3
 Erysipelas, streptococci in, 120
 Erythema bullosum, eosinophilia with, 96
 nodosum, eosinophilia with, 92
 Erythro-dextrin, 162
 Esbach's albuminometer, 53
 fluid, 53
 for serous effusions, 158
 Ewald's test breakfast, 139
 Exercise, effect on urine, 1
 urate deposit after, 13
 Eyes, embedding in celloidin, 170

 Fæces, examination of (and see Stools), 140
 Fat globules in urine, 23
 osmic acid stain for, 23
 Sudan III. stain for, 23, 99
 Fatty stools, 141
 Favus, 156

- Fehling's solution, formulary for,
 34
 reduced after the ad-
 ministration of various
 drugs, 35, 51, 52
 reduced by alkapton, 35,
 49
 by cerebro - spinal
 fluid, 161
 by glycuronic acid,
 41
 test, 34
 fallacies of, 35, 36
 influence of creatinin on,
 35, 36
 Fermentation method for estimat-
 ing sugar, 56, 57
 test for glucose, 38
 Ferments in pancreatic cyst fluid,
 162
 Ferric chloride, precipitation of
 phosphates by, 49
 reaction in alkaptonuria,
 49
 in carboluria, 41, 49,
 52
 in melanuria, 49
 in urine after various
 drugs, 41, 51, 52
 with diacetic acid, 41
 Ferrous sulphate, test for nitric
 acid, 178
 Fevers, chlorides in urine reduced
 in, 44
 gastric HCl in, 135
 urate deposit in, 13
 urine decreased in, 1
 highly acid in, 3
 Fibroid lung, hæmoptysis in,
 107
 polycythæmia from, 71
 sputum in, 107
 Filaria, chyluria from, 24
 sanguinis hominis, 97, 98
 Filariasis, eosinophilia in, 92, 96
 parasites in blood in, 96
 Films, fixation of, 79
 Fish, bothriocephalus in, 147
 Fixing methods, 79, 145, 166, 170
 Flagella, 121, 125
 Formaldehyde method of esti-
 mating ammonia, 64
 Formalin as tissue preservative,
 165
 method of fixing blood films,
 79
 Fraenckel, pneumococci of, 115,
 120
 Fractured skull, cerebro - spinal
 fluid from ear in, 161
 Fractures, lipæmia after, 100
 Friedländer, pneumobacilli of,
 115
 Fungating endocarditis, blood
 cultures in diagnosing,
 102
 colour index in, 77
 gonococci in, 120
 no leucocytosis with, 95
 pneumococci in, 121
 simulated by pernicious
 anæmia, 77
 streptococci in, 120
 Fusiform bacillus, Vincent's, 127
 Gaertner's bacillus, 146
 Gall-stones, cholesterin in, 144
 in stools, 144
 stools with, 140
 Gallic acid, ferric chloride reaction
 in urine after, 51
 Gangrene of lung, elastic fibres in
 sputum in, 110
 Gans' apparatus for estimating
 sugar, 56
 Gastric carcinoma, hydrochloric
 acid changes with, 134
 contents, examination of, 128
 hydrochloric acid in. See
 Hydrochloric acid
 lactic acid in, 131, 135
 sarcinæ in, 138, 139
 ulcer, acetoneuria in, 41
 blood in stools from, 142
 hyperacidity with, 136
 General paralysis of the insane,
 cerebro-spinal fluid in, 161
 Gentian violet stain, 118
 Germain Sée's test lunch, 139
 Giemsa's stain, 126
 Gigantoblast, 88
 Glanders bacilli, 124
 from horses, 125
 Gleet, pus cells in urine from, 17

- Glucose in urine, diacetic acid and acetone with, 39
 methods of estimating, 54, 55, 56, 57
 significance of, 39
 tests for, 34, 35, 36, 37, 38, 39
- Glycerin, reduction of Fehling's solution after, 35
- Glycosuric acid, reduction of Fehling's solution by, 35
- Glycuronic acid, fallacy with phenyl-hydrazine test, 38
 from chloral, 51
 from chloroform, 51
 in urine, drugs causing, 42
 reduction of Böttger's solution by, 37, 41
 of Fehling's solution by, 35, 41
- Glycyrrhizin, reduction of Fehling's solution after, 35
- Gmelin's test for bile pigments, 42
- Gonococci, characters of, 120
 Gram-negative, 119
 in urine, 21, 22
 lesions due to, 120
 medico-legal importance, 120
- Gonorrhœa, prostatic threads after, 25
 pus cells in urine after, 17
- Gout, glycosuria in, 39
 relation of uric acid to, 8
- Gower's hæmoglobinometer, 73
- Gram-negative organisms, 119
 -positive organisms, 119
- Gram's iodine solution, 118
 in section staining, 168
 stain, 117, 118, 119
- Granular kidney, urine changes in, 1, 2, 3
- Grass-green stools, 141
- Growth of bladder, particles in urine, 16
 of lung, red - currant - jelly sputum with, 108
 of kidney, renal cells in urine in, 15
- Guaiacol, greenish - black urine from, 4
- Guaiacum test, fallacies of, 29, 30
- Guaiacum test for blood in stools, 143
 for hæmaturia, 29
 for iodides, 51
 for pus, 32
- Guanin, 63
- Gulland's solution for fixing amœba, 145
- Gunsburg's reagent, 128
 test, 132
- Hæmatin, 143
- Hæmatoporphyrin, spectrum of, 52
- Hæmatoporphyrinuria from sulphonal, 52
 simulating bile in urine, 42
- Hæmatoxylin, Delafield's, 82
 Ehrlich's, 169
 red-brown urine from, 3
- Hæmaturia, albuminuria with, 29
 colour of urine in, 3, 8
 conditions simulating, 42, 49
 discoloration of phosphates in, 5, 13
 from bilharzia, 24
 in Bright's disease, 19
 leucocytes in urine in, 17
 sources of, 18, 32
 tests for, 29, 30, 31
- Hæmocytometer in calcium estimation, 105
 Thoma-Zeiss, 65
- Hæmoglobin, estimation of, 73, 74, 75, 76, 99
 specific gravity measurement of, 99
 spectrum, 9, 31
- Hæmoglobinometer, Gower's, 73
 Haldane's, 73
 Tallqvist's, 76
- Hæmoglobinuria, paroxysmal, 31
 simulating bile in urine, 42
- Hæmophilia, importance of calcium in, 105
- Hæmoptysis, causes of, 107
- Hæmorrhage, colour index after, 77
- Haines's test for glucose, 34
- Hair, arsenic in, 176
- Haldane's hæmoglobinometer, 73
- Hayem's fluid, 67

- Head injury, glycosuria after, 39
 pallor of urine after, 3
 polyuria after, 1
- Heart disease, gastric HCl in, 135
 failure, albuminuria with, 29
 high specific gravity of urine in, 2
 occasional tube-cast in urine in, 19
 renal cells in urine in, 15
 urate deposit in, 13
 urine decreased in, 1
- Heat method of fixing films, 79
 test for albumin, fallacies of, 27
 effervescence in, 27
 for urates, 12
- Heller's test for albumin, 25
 fallacies of, 26
 for hæmaturia, 30
- Helmitol, reduction of Fehling's solution after, 35
- Hepatic abscess, amœbic, 125
 anchovy sauce pus from, 108
 chocolate pus in, 125
 leucocytosis with, 73
- Herpes gestationis, eosinophilia with, 92, 96
- Hides, anthrax from, 123
- Hippuric acid, 35
- Hodgkin's disease, blood changes in, 93
 colour index in, 77
 lymphocytes in, 91
 myelocytes in, 92
- Homogentisic acid, 49
- Hooklets, hydatid, 162
 of tapeworms, 147, 148
- Hopkins's method of estimating uric acid, 60
- Horses, glanders from, 125
- Hyaline lymphocytes, 89
- Hydatid cyst fluid, 162
 disease, 148
 eosinophilia with, 96
 hooklets, 162
- Hydroa herpetiformis, eosinophilia with, 92, 96
- Hydrocele fluid, 163
- Hydrochloric acid, gastric, combined, and free, 131
 free, tests for, 132, 133, 134
 significance of, 134, 135
 test for, 177
- Hydrocyanic acid, test for, 178
- Hydroquinone, 49
- Hyperacidity of urine, 8, 9
 of vomit, estimation of, 136
- Hypobromite solution, 57
- Hypoxanthin, 35, 63
- Hysteria, urine changes in, 1, 2, 3
- Impetigo, absence of leucocytosis with, 95
- Indican, brown - red ring with nitric acid, 25
 greenish-black urine from, 4
 tests for, 45
- Indicanuria, darkening with potash, 37
 significance and causes of, 46
 simulating bile in urine, 42
- Indol, 46
- Infants, small lymphocytes in, 90
 sodium urate in, 11
- Infarction of kidney, renal cells in urine in, 15
 tube-casts in, 19
 of lung, prune-juice sputum with, 108
- Infective endocarditis. See Fungating endocarditis
- Inferior vena cava thrombosis, albuminuria with, 29
- Influenza bacilli in cerebro-spinal fluid, 160
 in pneumonia, 115
 in sputum, 109
- Injury, chyluria from, 24
- Intestinal casts, 141
 putrefaction, ethereal sulphates in urine in, 44
 indicanuria from, 46
 sand, 142
- Intussusception, blood and mucus in stool with, 140, 142

- Iodide test for corrosive sublimate, 177
 Iodides, fallacies in testing for indican, 45
 fallacy in guaiacum test, 29
 in urine, tests for, 51
 Iodine, sarcinæ stained by, 138
 solution, Gram's, 118
 test for bile pigments, 42
 for starch grains, 14, 138
 Iodoform test for acetone, 40
 Irritability of bladder, 8, 9
 Iron alum in chloride estimation, 61
 black motions from, 142

 Jaundice, clay - coloured stools with, 141
 Jaw, actinomycosis of, 124
 Jenner's stain, 80
 Joseph's stain for malaria parasites, 85

 Kaiserling's solution, 166
 Kidney growth cells in urine, 15
 infarction, 15, 19
 tuberculous, pyuria from, 17, 22
 Klebs-Loeffler bacilli, 122
 Klemperer's test, breakfast, 139
 Knife-rester crystals, 9

 Lactic acid in gastric contents, 131, 135
 Uffelmann's test for, 129, 133
 Lactose in urine, 33, 39
 reaction with picric acid, 37
 Böttger's solution, 37
 Lardaceous disease, albuminuria in, 29
 tissue, stain for, 171
 Laryngitis, pneumococci in, 121
 Lead in urine, tests for, 50
 reduction of Fehling's solution by, 35
 Leishman's stain, 83
 Lemon-pips, parasitic ova simulated by, 150
 Leprosy bacilli, acid-fast, 112
 Leucin, 11
 Leucin simulated by starch grains, 11, 14
 Leucocytes, differential count of, 88, 93, 94
 enumeration of, 65, 71, 73
 in cachexia, 91, 95
 in children, 93
 in urine, 17
 normal numbers of, 72
 simulating fragment of growth in urine, 16
 renal cells, 15
 Leucocytosis in abscess formation, 95
 in appendicitis, 73
 in cachexia, 95
 in empyema, 91
 in hepatic abscess, 73
 in leukæmia, 91, 94
 in pneumonia, 73, 95
 in pyosalpinx, 73, 92
 in suppuration, 72, 92, 95
 Leucopenia in malaria, 72, 95
 in typhoid fever, 73, 95
 Leukæmia, basophile cells in, 93
 blood changes in, 91, 92, 94
 Charcot-Leyden crystals in, 116
 colour index in, 77
 differential leucocyte count in, 93
 leucocytes in, 73
 purpura in, 32
 Life insurance, relation of nucleoproteid to, 48
 Lipæmia, causes and significance of, 99, 100
 Liquor potassæ in examining ringworm fringe, 155, 156
 test for glucose, 37
 for pus, 32
 Liverabscess. See Hepatic abscess
 actinomycosis of, 124
 affections, leucin and tyrosin in, 11
 cirrhosis. See Cirrhosis
 Lung, abscess of, sputum in 110
 gangrene of, 110
 glanders of, 125
 growth of, red-currant jelly sputum in, 108

- Lung infarction, prune-juice sputum from, 108
- Lymphatic leukæmia. See Leukæmia
- Lymphocytes, conditions in which increased, 90, 91
- in cachexia, 91, 95
 - in cerebro-spinal fluid, 161
 - in leukæmia, 91, 94
 - in malaria, 95
 - in tuberculous effusion, 160
 - in typhoid fever, 90, 95
 - large or hyaline, 89
 - small, 88
- Magnesium phosphate, 10, 13
- sulphate and nitric acid, test for albumin, 28
 - wire test for lead, 50
- Malaria, differential leucocyte count in, 93
- leucopenia in, 73, 95
 - lymphocytes in, 91, 95
 - parasites, effect of quinine on, 91, 97
 - in blood, 96, 97
 - Joseph's stain for, 85
 - Leishman's stain for, 83
 - Romanowski's stain for, 85
- Malfatti's method of estimating ammonia, 64
- Malignant disease, lymphocytes in, 91
- endocarditis. See Fungating endocarditis
 - pustule, 123
- Manganese, black motions from, 142
- Marsh's test, 175
- Mathison, *re* estimation of ammonia, 64
- Meat-eaters, acidity of urine in, 3
- uric acid in, 9
- Megaloblasts, 87
- Megalocytes, 87
- in pernicious anæmia, 88, 95
- Melæna, causes of, 142
- conditions simulating, 142
- Melanin, 4
- Melanin, sodium nitroprusside test for, 49
- Melanogen, 49
- Melanuria, significance of, 49, 50
- simulating bile in urine, 42, 49
 - hæmaturia, 49
 - tests for, 49, 50
- Meningitis, cerebro-spinal fluid in, 160
- meningococci in, 121
 - pneumococci in, 121
- Meningococci, in cerebro-spinal fluid, 161
- Weichselbaum's, 121
- Menorrhagia, importance of calcium in, 105
- Menstruation, albuminuria from, 29
- hæmaturia simulated by, 18
- Mercury, reduction of Fehling's solution after, 35
- Metalbumin, 163
- Methæmoglobin, spectrum of, 31
- Methæmoglobinuria, 31, 32
- simulating bile in urine, 42
- Methyl green, green urine from, 4
- mercaptan after asparagus, 4
 - orange, 131
 - violet solution, 131
 - stain for lardaceous change, 171
 - test for free HCl, 134
- Methylene blue alcoholic solution, 84
- and eosin stain, 84
 - aqueous solution, 84
 - urine coloured from, 4
 - normal saline, 5
- Microblast, 87
- Micrococcus ureæ, 3, 20
- Microcyte, 87
- Microscope section, fixing of, 166
- sections, mounting of, 168
 - test for blood, 31
 - for pus, 33
- Microsporon furfur, 156
- Microtome, 168
- freezing, 169
- Micturition frequent from oxaluria, 8
- Milk, undigested in stools, 143

- Mitral stenosis, blood changes in, 96
 hæmoptysis in, 107
 polycythæmia from, 71
 Mohr's method of estimating chlorides, 62
 Moore's test for glucose, fallacies of, 37
 Morbid anatomy specimens, preservation of, 165
 Morbus cæruleus, blood changes in, 96
 polycythæmia in, 71
 Mörner Sjöqvist, method of estimating urea, 60
 Morphia, test for, 174
 Morphine, reduction of Fehling's solution after, 35
 Motions. See Stools
 Mucous colitis, intestinal sand with, 142
 stools, 141
 Mucus, cylindroids of, 25
 effect of acetic acid on, 13
 in urine, spermatozoa with, 20
 in stools, 140, 141
 microscopic appearances, 25
 simulated by phosphates, 13
 simulating pus, 32
 tube-casts, 25
 white deposit from, 5
 with oxaluria, 20
 Mulberry calculus, 8
 Müller-formol fixing fluid, 170
 Müller's fluid, 170, 172
 Myelocytes, 90, 92, 94
 Myeloid sarcoma, albuminuria from, 27
- Nails, arsenic in, 176
 Naphthalene, greenish-black urine from, 4
 Naphthol, greenish-black urine from, 4
 Neisser's stain, 122
 Nematodes, 148
 Nervousness, effect on specific gravity of urine, 2
 pallor of urine from, 3
 Nephritis, albuminuria in, 29
 Nephritis, anuria in, 1
 chlorides diminished in urine in, 44
 chyluria in, 24
 gastric HCl in, 135
 hæmaturia in, 19
 leucocytes in urine in, 17
 renal epithelial cells in, 15
 suppurative, pus cells in urine in, 17
 urine decreased in, 1
 varieties of casts in, 19, 20
 Neutrophile cells, 89
 Nitric acid, albumin precipitated by, 13
 method of estimating albumin, 54
 phosphates dissolved by, 13
 ring after copaiba, 26, 51
 cubeb, 51
 with albumose, 26
 balsams, 51
 indican, 25
 test for, 178
 for albumin, 25, 28
 fallacies of, 26, 51
 significance of brown-red ring with, 26
 for albumose, 47
 for bile, 42
 Nitro - benzol, glycuronic acid from, 42
 -hydrochloric acid, test for indican, 45
 Nitroprusside test for acetone, 39
 for melanin, 49
 Nocturnal enuresis from coli bacilluria, 22
 from oxaluria, 8
 Normoblasts, 87
 Normocytes, 87
 Nucleated red cells, 87
 Nuclei brought out by acetic acid 17, 33
 Nucleo-proteid, cloud with heat test, 27
 differentiation from albumin, 27, 48

- Nucleo-proteid, precipitated by
 acetic acid, 48
 spermatozoa with, 20
 with oxaluria, 8, 20, 48
- Oatmeal, oxyuris simulated after,
 150
- Occult blood, detection of, 143
- Old persons, glycosuria in, 39
- Onions, oxyuris simulated after,
 150
- Operation, anuria after, 1
- Ophthalmia, gonococci in, 120
- Opium, reduction of Fehling's
 solution after, 35
 test for, 174
- Opsonic index, 102, 103
- Orange-pips, parasitic ova simu-
 lated by, 150
- Osmic acid stain for fat globules,
 23
- Osteomyelitis, streptococci in,
 120
- Otitis media, pneumococci in,
 121
- Otto's alkaloid method, 174
- Ova in fæces, 150
- Ovarian cyst fluid, 163
- Ovum of ascaris lumbricoides,
 152
 of bilharzia hæmatobia, 24
 of bothriocephalus latus, 153
 of tænia mediocanellata, 152
 of tænia solium, 152
 of trichocephalus dispar, 152
- Ox, tænia mediocanellata in, 147
- Oxalate crystals, agglomeration
 of, 8
 confirmative of calculus,
 8
 deposit, powdered-wig ap-
 pearance of, 7
 striæ in, 7
 test for lead, 50
- Oxalic acid, test for, 178
- Oxaluria, characters of deposit in,
 5, 7
 frequent micturition from, 8
 mucus with, 20
 nocturnal enuresis from, 8
 nucleo-proteid with, 8, 20, 48
 significance of, 7
- Oxaluria, spermatorrhœa from, 8
 spermatozoa with, 8, 20
- Oxyhæmoglobin, spectrum of, 31
- Oxyphile cells, finely granular, 89
- Oxyuris vermicularis, 148
 no eosinophilia with, 92
 simulated by vegetable
 fibres, 150
- Ozonic ether, bubbling with blood,
 32
 test for pus, 32
- Pancreas, stools in affections of,
 141
- Pancreatic cyst fluid, 161
 ferments in, 162
- Paraffin, embedding in, 167
- Parovarian cyst fluid, 163
- Parasites, filarial, 96
 intestinal, 146-153
 things simulating, 150
 in trypanosomiasis, 96
 malarial, 91, 96, 97
- Parasitic affections, eosinophilia
 in, 92, 96
- Paratyphoid bacillus, 146
- Paroxysmal hæmoglobinuria, 31
- Patent interventricular septum
 blood changes in, 96
- Pavy's solution, 54
- Pea-soup stool, 140
- Pemphigus, eosinophilia with, 92,
 96
- Pentose in urine, 33
- Pepsin, test for, 130, 137
- Pericarditis, pneumococci in, 121
- Periostitis, staphylococci in, 120
- Peritoneal fluid, characters of, 158
- Peritonitis, chlorides absent from
 urine in, 44
 micro-organisms causing, 120,
 121
 no leucocytosis with, 95
- Pernicious anæmia, ankylosto-
 miasis simulating, 150
 blood changes in, 95
 colour index in, 77
 leucin and tyrosin in, 11
 megalocytes in, 88, 95
 myelocytes in, 92
 red corpuscle changes in,
 87, 88

- Pernicious anæmia, simulated by carcinoma of stomach, 77
by infective endocarditis, 77
urobilin in, 46
- Perspiration, effect on urine, 1, 2, 13
- Pettenkofer's test for bile salts, 42
- Phenolphthaleïn, 64, 130
- Phenyl compounds, ferric chloride reaction in urine after, 41
greenish - black urine from, 4
- Phenyl-glucosazone crystals, 11, 38
-hydrazine test for glucose, 38, 137
-lactosazone crystals, 38
- Phloroglucin, 128
test for urotropin, 52
- Phosphate fallacy in Fehling's test, 36
of calcium, 10, 13
of magnesium, 10, 13
of spermin, 116
- Phosphates, acetic acid test for, 13, 27
ammonio-magnesian, 9, 11
amorphous, 13
brown from blood, 5, 13
dissolved by nitric acid, 13
earthy, 10, 13
estimation of, 62
forming cloud on heating, 13, 27
mistaken for pus, 9, 13, 32
precipitation by ferric chloride, 49
of blood pigment by, 30
simulated by mucus, 13
by urates, 13
stellar, 10, 13
white deposit of, 1, 5, 13
- Phosphorus poisoning, leucin and tyrosin in, 11
lipæmia in, 100
test for, 178
- Phthisis, Charcot-Leyden crystals in, 116
diazo-reaction in, 47
elastic fibres in sputum in, 110, 113
- Phthisis, gastric HCl in, 135
læmptysis in, 107, 109
pneumococci in, 121
sarcinæ in, 114
secondary infection in, 113
slow decomposition of urine in, 21
sputum in, 107, 109
streptococci in, 120
tetrads in, 114
- Picramic acid, 37
- Picric acid in Esbach's fluid, 53, 54
precipitation of urine pigments by, 54
test for albumin, 28
for glucose, 37
- Pig, tænia solium in, 147
- Pigments, precipitation by picric acid, 54
- Pike, bothriocephalus in, 147
- Piles, blood-stained stools from, 142
- Pine-apple, oxyuris simulated after, 150
- Pipettes, method of cleaning, 68
- Plague bacilli, 125
Gram-negative, 119
- Plastic bronchitis, 108
- Pleuritic effusion, characters of, 158
- Pneumo - bacilli, Gram - negative, 119
in pneumonia, 115
- Pneumococci, capsule stain for, 115
characters of, 120
Gram-positive, 119
in cerebro-spinal fluid, 161
in pneumonia, 115, 121
in sputum, 109, 113
occurrence in health, 121
various lesions due to, 121
- Pneumonia, acetoneuria in, 41
albuminuria in, 47
chlorides absent from urine in, 44
diazo-reaction in, 47
leucocytosis in, 73, 95
micro-organisms producing, 114, 115
polymorphonuclear cells in, 91, 95

- Pneumonia, sputum in, 107, 108, 114, 115
 urine decreased in, 1
 highly acid in, 3
- Poikilocytes, 87, 88
- Poisons, tests for, 174
- Polychromasia, 87
- Polychromatophilia, 87
- Polychrome methylene blue, 86
- Polycythæmia, various causes of, 71, 96
- Polymorphonuclear cells, 89
 in abscess formation, 73, 91, 95
 in cerebro-spinal fluid, 161
 in pneumonia, 91, 95
 relatively increased, 91
- Polyuria, various causes of, 1
- Potash, darkening of urine by, 37, 49
 in examining ringworm hairs, 155, 156
 test for acetone, 40
 for alkaptonuria, 37, 49
 for carboluria, 37, 49
 for glucose, 37
 for melanuria, 49
 for pyuria, 32
 for rhubarb in urine, 51
 for santonin, 52
- Potassium bichromate test for morphia, 174
 test for strychnine, 175
 chromate in estimating chlorides, 62
 test for lead, 50
 citrate in preventing thrombosis, 102
 ferro-cyanide in estimating phosphates, 62
 test for albumin, 28
 for copper, 177
 iodide in actinomycosis, 124
 in urine, 29, 45, 51
 test for lead, 50
 oxalate in estimation of ammonia, 64
 permanganate method of estimating uric acid, 60
 sulpho-cyanide in estimating chlorides, 61
- Potassium, urate deposit, 12
- Powdered - wig appearance of oxalates, 7
- Pregnancy, lactose in urine in, 39
- Prostatic threads, 24, 25
- Prostatitis, pus cells in urine in, 17
- Pine-juice sputum, 108
- Prussic acid, test for, 178
- Pseudo-leukæmia, lymphocytes in 91
 -mucin, 163
 -parasites, intestinal, 150
- Psoriasis, no eosinophilia with, 92
- Ptomaine poisoning, 146
- Pulmonary abscess, elastic fibres in sputum in, 110
 stenosis, blood changes in, 96
- Punctate basophilia, 87
- Purin bases, estimation of, 63
- Purinometer, 63
- Purpura, hæmaturia in 32
 in leukæmia, 32
- Pus cells in sputum, 114
 in urine, 17
 anchovy-sauce coloured, 108
 chocolate coloured, 125
 effect of acetic acid on, 13
 in stools, 141
 in urine. See Pyuria
 under pressure, albumosuria with, 47
- Pyæmia, streptococci in, 120
- Pyelitis, bacillus coli in, 121
 tailed renal pelvic cells in, 16
- Pyelonephritis, bacillus coli in, 121
- Pyonephrosis, pyuria with, 33
- Pyorrhœa alveolaris, streptococci in, 120
- Pyosalpinx, leucocytosis with, 73, 92
- Pyriform cells from bladder, 15
- Pyrocatechin, 49
 in cerebro-spinal fluid, 160
- Pyuria, albuminuria with, 17, 29, 33
 causes and significance of, 33
 effect on methylene blue in urine, 4
 fallacies in guaiacum test, 29
 from gonorrhœa or gleet, 17
 hæmaturia with, 29

- Pyuria, mucus simulating, 32
 phosphates simulating, 9, 13, 32
 tests for, 32, 33
 urates simulating, 32
 white deposit from, 5
 with coli bacilluria, 17, 33
 with cystitis, 17, 22, 33
 with tuberculous bladder, 22
 kidney, 22
- Quinine effect on malarial parasites, 91, 97
- Ray fungi, 124
- Reaction, amphoteric, 3
- Receptaculum chyli, chylous ascites from injury to, 160
 chyluria from injury to, 24
- Red corpuscles, crenated, 86
 enumeration of, 65, 71
 pathological changes in, 87
- Red-currant jelly sputum, 108
- Reinsch's test, 176
- Relapsing fever, spirillum of, 98
- Renal calculus, pyuria from, 17
 cells, 15, 16, 17
 growth, particles in urine, 16
 pyuria from, 17
 tube-casts, significance of, 19
 varieties of, 18, 19
 tuberculosis, pyuria from, 17, 22
- Rennin, test for, 137
- Resin, white ring with nitric acid, 26
- Resorcin, 131
 greenish-black urine from, 4
- Retina, lipæmia detected in, 99
- Rhubarb, chrysophanic acid from, 52
 oxalates after, 7
 oxyuris simulated after, 150
 red-brown urine from, 3, 51
 source of fallacy in Heller's test for blood, 30
 tests for, in urine, 51
- Rice-water stools, 125, 140
- Rings, malarial, 97
- Ringworm, 154
 potash in examining hairs in, 155, 156
- Roberts's method of estimating albumin, 54
- Rochelle salt in Böttger's test, 36
 in Fehling's solution, 34
 in Pavy's solution, 54
- Romanowski's stain, 85
- Rosettes of uric acid, 8
- Rothera's test for acetone, 40
- Salicin, ferric chloride reaction of urine after, 41
- Salicylates, ferric chloride reaction in urine after, 41, 52
 Fehling's solution reduced after, 35, 52
 glycuronic acid from, 42
- Salicyl-sulphonic acid test for albumin, 28
- Saliva fallacy of guaiacum test, 29
- Salol, ferric chloride reaction in urine after, 41, 52
 Fehling's solution reduced after, 52
 greenish-black urine from, 4
- Sand, intestinal, 142
- Santonin, bright yellow urine from, 4, 52
 source of fallacy in Heller's test for blood, 30
 test for, in urine, 52
- Sarcinæ in sputum, 114
 ventriculi, 138, 139
- Sarcoma, melanotic, melanuria in, 50
 of bone, albumosuria with, 27, 47
- Sarcoptes hominis, 156
- Scabies, 156
- Scurvy, importance of calcium in, 105
- Season, effect on urine, 1
- Sections, staining of, 168
- Semen, examination of stains due to, 172
- Seminal emission, albuminuria after, 29
- Senna, chrysophanic acid from, 52

- Senna, source of fallacy in Heller's
 test for blood, 30
 urine changes from, 3, 52
 Septicæmia, pneumococci in, 121
 staphylococci in, 120
 streptococci in, 120
 Serous effusions, characters of,
 158, 159, 160
 Serum treatment of anthrax, 123
 Shiga's bacillus, 126, 146
 Shock, specific gravity of blood in,
 99
 Silver nitrate, method of estimat-
 ing chlorides, 62
 test for arsenic, 176
 for chloral, 176
 for chlorides, 43
 for hydrochloric acid,
 177
 for prussic acid, 178
 Skatol, 46
 Skin diseases, eosinophilia with,
 92
 Skins, glanders from, 125
 Skull, fractured, cerebro-spinal
 fluid from ear in, 161
 Slides, method of albuminizing, 21
 Stools in typhoid stools, 144
 Smegma bacilli, acid-fast, 112
 simulating tubercle ba-
 cilli, 22, 23
 Soda solution, decinormal, 129
 Sodium acetate in estimating
 phosphates, 62
 biurate, 9
 hydrobromite method of esti-
 mating urea, 57
 nitrate in diazo-reaction, 46
 nitroprusside test for acetone,
 39
 test for melanin, 49
 phosphate causing acidity of
 urine, 3
 in gastric contents, 131
 quadriurate, 9
 urate, 10, 11, 12
 Sore throat, diphtheria bacilli in,
 122, 123
 South Africa, bilharzia in, 24
 Southall's ureometer, 108
 Specific gravity method of esti-
 mating sugar, 57
 Specific gravity of blood, 98, 99
 relation to hæmc-
 globin, 99
 of serous effusions, 158
 of urine, 2
 Spectroscope test for blood, 30,
 143
 Spectrum of hæmatoporphyrin,
 52
 of methæmoglobin, 31
 of oxy hæmoglobin, 31
 of urobilin, 46
 Spermatocoele fluid, 163
 Spermatozoa in spermatocoele, 163
 in stains, 172
 in urine, 20
 with oxalates, 8, 20
 Spermin phosphate, 116
 Spirals, Curschmann's, 116
 Spirillum of relapsing fever, 98
 Vincent's, 127
 Spirochæta, stains for, 126
 Splenic anæmia, blood changes
 in, 94
 colour index in, 77
 myelocytes in, 92
 Splenomedullary leukæmia. See
 Leukæmia
 Splenomegalic polycythæmia, 71
 Spores in tinea, 155
 Sputum anchovy-sauce coloured,
 108
 black, 108
 casts in, 108
 Charcot-Leyden crystals in,
 116
 Curschmann's spirals in, 116
 elastic fibres in, 109, 110,
 113
 films made from, 110
 from empyema, 107
 in bronchiectasis, 107
 in bronchitis, 107
 in fibroid lung, 107
 in lung abscess, 110
 in phthisis, 107, 109
 in plastic bronchitis, 108
 in pneumonia, 107, 108, 114,
 115
 influerza bacilli in, 109
 microscopical characters, 109-
 116

- Sputum, naked-eye characters,
 107, 108
 nummular, 107, 109
 prune-juice type, 108
 pus cells in, 114
 red-currant jelly type, 108
 rusty, 108
 sarcinæ in, 114
 staphylococci in, 109, 113
 streptococci in, 109, 113, 120
 tetrads in, 109, 113, 114
 tubercle bacilli in, 109, 110,
 111, 112, 113
 various bacteria in, 109
- Squamous cells, 14
- Staining of blood films, 80, 86
- Stains, examination of seminal,
 172
 gonococci in, 120
- Staphylococci, Gram-positive, 119
 in cerebro-spinal fluid, 161
 in pneumonia, 114
 in sputum, 109, 113
 in urine, 21
- Staphylococcus, lesions due to,
 119, 120
- Starch grains in urine, 5, 11, 14
 iodine test for, 14, 138
 mistaken for leucin, 11,
 14
- Starvation, acetonuria in, 41
 chlorides absent from urine
 in, 44
 influence of sulphates in
 urine, 44
- Stellar phosphate, 10, 13
- Stercobilin, 151
- Stools, *œmæba* of dysentery in,
 145
 bacteria in, 141
 black, 142
 blood and mucus in, 140
 in, 142
 clay coloured, 140
 fat in, 141
 gall-stones in, 144
 in biliary catarrh, 140
 milk in, 143
 ova in, 150
 pus in, 141
 rice-water, 125, 140
 typhoid bacilli in, 142
- Stools, vegetable remains in,
 150
- Streptococci, Gram-positive, 119
 in cerebro-spinal fluid, 161
 in pneumonia, 114
 in sputum, 109, 113, 120
 in urine, 21
 lesions due to, 120
- Striæ in oxalate deposit, 7
- Strychnine, test for, 175
- Sucking discs, 147
- Sudan III. stain for fat, 23, 99
- Sugar in urine. See Glucose in
 urine
- Sulphanilic acid in diazo-reaction,
 46
- Sulphate test for lead, 50
 test for nitric acid, 178
- Sulphates, ethereal, 44
 in urine, estimation of, 44
 significance of, 44
- Sulphonal, hæmatoporphyria
 from, 52
- Sulphur test for bile salts, 43
- Sulphuric acid, test for, 178
- Summer, urate deposit in, 13
- Suppuration, leucocytosis with,
 72, 92, 95
 polymorphonuclear cells in-
 creased in, 91
- Sweets, eosinophilia from, 3
 green urine after, 4
- Synovitis, gonococci in, 120
- S. philis, lymphocytes in, 90, 91
 spirochæta in, 126
 Wassermann's reaction in,
 104, 161
- Tabes dorsalis, cerebro-spinal
 fluid in, 61
- Tænia *echinococcus*, 148
mediocanellata, eosinophilia
 with, 96
 occurrence in ox, 147
 ovum of, 152
 solium, 147
 eosinophilia with, 96
 ovum of, 152
- Tailed cells from renal pelvis, 16
- Tallqvist's hæmoglobinometer, 76
- Tannic acid, ferric chloride re-
 action in urine after, 51

- Tapeworms, eosinophilia with, 92
 Test meals, 139
 Tetanus bacillus, 123
 Tetany, acetonuria in, 41
 Te'rads in sputum, 109, 113, 114
 Thallin, greenish-black urine from, 4
 Thoma's fluid, 67
 Thoma-Zeiss's hæmocytometer, 65
 in calcium estimation, 105
 Thoracic duct, obstruction, chylous ascites from, 160
 chyluria from, 24
 Thorn-apple crystals, 10, 11
 Threads, prostatic, 24, 25
 Thrombosis in kidney, renal tube-casts from, 19
 prevention by citrate of potash, 102
 Thymol, 149
 in preserving urine, 21
 Tinea circinata, 154
 spores of, 155
 stains for, 155
 tonsurans, 154
 no eosinophilia with, 92
 spores of, 155
 versicolor, 155
 Torulæ, 138
 Transfusion indicated by blood specific gravity, 99
 Trichinosis, eosinophilia in, 92, 96
 Trichocephalus dispar, 150
 no eosinophilia with, 92
 ovum of, 152
 Trichophyton megalosporon ectothrix, 154
 microsporon endothrix, 154
 Triple stain, Ehrlich's, 85
 Trommer's test for glucose, 36
 Tropæolin OO, 131
 test, 133
 Trypanosomiasis parasites in blood, 96
 Tube-casts, simulated by mucus, 25
 Tubercle bacilli, antiformin process for detecting, 110
 characters of, 112
 in cerebro-spinal fluid, 161
 Tubercle bacilli in sputum, 109, 110, 111, 112, 113
 in urine, 21, 22
 Ziehl-Neelsen stain for, 111
 Tuberculosis, lymphocytes in, 91
 Tuberculous bladder, pyuria from, 22
 cachexia, colour index in, 77
 effusions, lymphocytes in, 160
 kidney, pyuria from, 17, 22
 renal cells in urine in, 15
 tailed renal pelvis cells in, 16
 without pyuria, 22
 toxæmias, colour index in, 77
 ulceration, blood in stools from, 142
 Turkish bath, urate deposit after, 13
 Turpentine, violet smell of urine after, 4
 Typhoid bacilli in urine, 21, 22
 bacillus, 119, 122, 146
 fever, blood cultures in diagnosing, 102
 in stools in, 142
 diazo-reaction in, 47
 leucopenia in, 73, 95
 lymphocytes in, 90, 95
 stools in, 140, 143
 Widal's reaction in, 100
 sloughs in stools, 144
 Tyrosin, 11
 Uffelmann's reagent, 129
 test, 133
 Ulcerative endocarditis. See Fungating endocarditis
 Uræmia, cerebro-spinal fluid in, 161
 Uranium nitrate in estimating phosphates, 62
 Urate of ammonium, 10, 11
 of potassium, 12
 of sodium, 10, 11, 12
 Urates, colourless, 5
 coloured by uroerythrin, 5, 12
 effect of acetic acid on, 13
 heat test for, 12

- Urates, microscopical appearances, 12
 phosphates simulating, 13
 pink deposit of, 1, 5, 12
 significance of, 13
 simulating pus, 32
 white deposit of, 12
 Urea in cerebro-spinal fluid, 161
 methods of estimating, 57, 58, 59, 60
 nitrate, 26
 Ureometer, Southall's, 58
 in fermentation test for glucose, 38
 Urethritis from uric acid, 9
 gonococci in, 120
 leucocytes in urine in, 17
 Uric acid, calculus from, 9
 cayenne-pepper deposit of, 5, 8
 crystals, 8
 estimation of, 60
 hyperacidity of urine with, 3
 irritability of bladder from, 8
 non-reduction of Böttger's solution by, 37
 normal total excreted, 60
 physiology of precipitation of, 9
 relation to gout, 8
 urethritis from, 9
 Urine, abnormal colours in
 alkaptonuria, 49
 in carboluria, 49
 in melanuria, 49
 alkapton in, 4, 35, 37, 42, 49, 50
 ammoniacal fallacy in Fehling's test, 35
 on standing, 4
 with pyuria, 4, 33
 amounts passed, 1
 bacillus coli in, 4, 17, 21, 22
 typhosus in, 21, 22
 bile in, 4, 49
 bilharzia eggs in, 24
 black, 3
 blood in. See Hæmaturia
 blue, 4
 bright red, 3
 yellow from santonin, 4, 52
 brown, 3
 carbolic acid in. See carboluria
 casts in, 19
 cayenne-pepper deposit in, 5, 8
 cells in, 15, 16
 chlorides in, 43, 60, 62
 colour of normal, 3
 contamination with milk, 23
 with oil, 23, 24
 dark red, 3
 darkening on exposure to air, 49
 with potash, 37
 diacetic acid in, 39, 41
 dichroic, 4
 drug reactions in, 3, 4, 41, 49, 51, 52
 eosin in, 3
 estimation of total solids in, 2
 fæcal smell with, 4
 fat globules in, 23
 ferric chloride reactions in, 41, 49, 51, 52
 fragments of new growth in, 16, 17
 glycuronic acid in, various causes of, 42
 gonococci in, 21, 22
 green, 4
 greenish-black, various causes of, 4
 yellow from bile pigments, 4
 from drugs, 4
 hyperacidity of, 3, 8
 iodides in, 29, 51
 lactose in, 33, 39
 leucocytes in, 17
 method of microscopical examination of, 5, 6
 methylene blue in, 4
 milky from phosphates, 10
 mucus in, 5, 13, 20, 25, 32
 oxalate crystals in, 8
 pallor of, 3
 pentose in, 33
 pink deposit in, 1, 5, 12

- Urine, port-wine coloured, 3
 from hæmatoporphy-
 rin, 52
 phosphates in. See Phos-
 phates
 preservation by boric acid,
 21
 by thymol, 21
 prostatic threads in, 24, 25
 pus cells in. See Pyuria
 reaction of, 3, 8, 9
 red-brown, from drugs, 3, 51
 red corpuscles in, 8, 17, 18
 renal cells in, 15, 16
 scarlet, 3
 slow decomposition in phthi-
 sis, 21
 smell of, 4
 specific gravity of, 1, 2
 starch grains in, 5, 11, 14
 sulphates in, 44
 urotropin in, 52
 various micro-organisms in,
 21
 veronal in, 179
 violet-smelling, 4
 white deposit in, 1, 5, 9, 12,
 13, 32
 yeasts in, 23, 138
- Urinometer, 2
 in estimating blood specific
 gravity, 98
- Urobilin, 4, 46
 spectrum of, 46
 zinc chloride test for, 46
- Urochrome, 3, 46
 in uric acid crystals, 8
- Uroerythrin, absence from urine,
 5
 urates coloured by, 5, 12
- Urotropin, Fehling's solution
 reduced after, 35, 52
 phloroglucin test for, 52
- Uterine discharge, albuminuria
 from, 29
 sepsis, streptococci in, 120
- Uva ursi, Fehling's solution re-
 duced after, 52
 ferric chloride reaction
 after, 52
 greenish - black urine
 from, 4
- Vagina, squamous cells from, 14
 Vaginitis, albuminuria from, 29
 Vanillin, 128
- Vegetable marrow pips, parasiti
 ova simulated by, 150
- Vegetables, oxalates after, 7
- Vegetarians, acidity of urine in, 3
 phosphaturia in, 10
- Veronal, test for in urine, 179
- Vesuvium, 122
- Vibrio of cholera, 125
- Vincent's angina, 126
- Violet smell of urine after turpen-
 tine, 4
- Volatile oils, reduction of Feh-
 ling's solution after, 35
- Volhard's method of estimating
 chlorides, 60, 61
- Vomit, causes of acidity of (and
 see Hydrochloric acid), 131
 food particles in, 137, 138
 methods of examining, 128,
 131
 yeasts in, 138
- Vomiting, urate deposit after, 13
- Von Jaksch's disease, lympho-
 cytes in, 91
- Vulvitis, bacillus coli in, 121
- Wassermann reaction in cerebro-
 spinal fluid, 161
 serum reaction, 104
- Wasting diseases, melanuria in,
 50
- Weichselbaum, meningococci of,
 121
- Whetstone crystals of uric acid, 8
- White corpuscles, enumeration of,
 65, 71, 73
- Whooping-cough, hæmoptysis in,
 107
 lymphocytes in, 90
- Widal reaction in typhoid fever,
 100
- Women, high specific gravity of
 urine in, 2
 mucus in urine of, 1, 25
- Wool, anthrax from, 123
- Xanthin, 63
 bases, non-reduction of Bött-
 ger's solution by, 37

Xanthin, reduction of Fehling's solution by, 35	Yeast method of estimating sugar, 56, 57
Xylol in clearing sections, 169	test for glucose, 38
tissues cleared by, 167	Ziehl-Neelsen method of staining tubercle bacilli, 111
Yeast fungi in urine, 23	Zinc chloride test for urobilin, 46
in vomit, 138	Zymotic enteritis, bacteria of, 146
	stools in, 141

THE END

BAILLIÈRE, TINDALL & COX'S
'UNIVERSITY SERIES'
OF MANUALS FOR STUDENTS.

BUCHANAN'S ANATOMY. (Systematic and Practical.) Price 21s. net.

Pp. xxvi+1562. Complete in One Volume.
With 621 Illustrations, mostly original and in several colours.
Or in 2 Parts, price 10s. 6d. each, net.

CASTELLANI & CHALMERS' TROPICAL MEDICINE. Price 21s. net.

Second Edition. Pp. xxxii+1748. With 15 coloured Plates and 630 Illustrations, mostly original.

GREEN'S PATHOLOGY AND MORBID ANATOMY. Price 15s. net.

Eleventh Edition. Pp. x+642. With 350 Illustrations.

JELLETT'S MIDWIFERY. Price 21s. net.

Second Edition. Pp. xii+1210
Revised throughout, with 17 Plates and 557 Illustrations, plain and coloured, of which nearly 300 are new.

MACNAUGHTON-JONES' DISEASES OF WOMEN. Price 21s. net.

Ninth Edition. Pp. xl+1044.
With 122 Plates and 637 Illustrations.

MONRO'S MEDICINE. Price 15s. net.

Third Edition. Pp. xxii+1024.
With 42 Illustrations, plain and coloured.

MOOR & HEWLETT'S APPLIED BACTERIOLOGY. Price 12s. 6d. net.

Third Edition. Pp. x+476. 102 Figures, plain and coloured.

ROSE & CARLESS' SURGERY. Price 21s. net.

Eighth Edition. Pp. xii+1406.
With 12 Coloured Plates and 581 Illustrations.

STEWART'S PHYSIOLOGY. Price 18s. net.

Sixth Edition. Pp. xx+1064.
With 2 Coloured Plates and 449 Illustrations.

TURNER'S MEDICAL ELECTRICITY. Price 10s. 6d. net.

Fourth Edition. Pp. xxiv+434. With 205 Illustrations.

BAILLIÈRE, TINDALL & COX,
8, HENRIETTA STREET, COVENT GARDEN, LONDON.

BOOKS FOR STUDENTS.

- Aarons' Gynæcological Therapeutics.** Pp. xiv+178, with 46 illustrations. Price 5s. net.
- Bayly's Clinical Pathology of Syphilis and Parasyphilis.** Fcap. Svo. Pp. xiv+194, with 3 plates and 20 illustrations in the text. Limp leather, gilt top. Price 5s. net.
- Clarke's Refraction of the Eye.** Third Edition. Pp. x+230, with 88 illustrations. Price 5s. net.
- Cross and Cole's Modern Microscopy.** Fourth Edition. Pp. xvi+320, with 100 plates and figures. Price 6s. net.
- Dieulafoy's Textbook of Medicine.** Second Edition. Pp. xxiv+2082, with 7 plates and 105 illustrations. In two volumes. Price 25s. net.
- Elder's Ship-Surgeon's Handbook.** Second Edition. Pp. xii+388. Price 5s. net.
- French's Medical Laboratory Methods and Tests.** Third Edition. Pp. viii+190, illustrated. Price 5s. net.
- Gardner's Surgical Anæsthesia.** Pp. xii+240, with 12 plates, and 35 illustrations in the text. Price 5s. net.
- Gemmell's Chemical Notes and Equations, Inorganic and Organic.** Second Edition. Pp. xiv+266. Price 5s. net.
- Gray's Diseases of the Ear.** Pp. xii+388, with 53 plates, of which 37 are stereoscopic, and 70 other illustrations. Price, with Stereoscope, 12s. 6d. net.
- Ince's Latin Grammar of Pharmacy, including the reading of Latin Prescriptions.** Eighth Edition. Price 5s. net.
- Lake's Handbook of Diseases of the Ear.** Fourth Edition. Pp. x+242, with 3 plates and 70 figures. Price 7s. 6d. net.
- Lamb's Practical Guide to the Diseases of the Throat, Nose and Ear.** Third Edition. Pp. xvi+252, with 57 illustrations. Price 7s. 6d. net.
- Macewen's Surgical Anatomy.** Pp. xii+452, with 61 illustrations, plain and coloured. Price 7s. 6d. net.
- May & Worth's Manual of the Diseases of the Eye.** Third Edition. Pp. viii+418, with 336 illustrations, including 22 coloured plates. Price 10s. 6d. net.
- Mummery's After-Treatment of Operations.** Third Edition. Pp. x+251, with 38 illustrations. Price 5s. net.
- Richards' Practical Chemistry, including Simple Volumetric Analysis and Toxicology.** Second Edition. Pp. x+150. Price 3s. net.
- Schmieden's The Course of Operative Surgery.** Pp. xx+350, with 435 illustrations, many coloured. Price 12s. 6d. net.
- Wheeler's Student's Handbook of Operative Surgery.** Second Edition. Pp. xiv+296, with 157 figures. Price 7s. 6d. net.
- Wihlta's Pharmacy, Materia Medica, and Therapeutics.** Ninth Edition. Pp. xiv+674, with 23 figures. Price 9s. net.
- Whitla's Dictionary of Treatment.** Fifth Edition. Pp. x+1204. Price 16s. net.
- Williams' Minor Maladies and their Treatment.** Third Edition. Pp. viii+396. Price 5s. net.

BAILLIÈRE, TINDALL & COX,

8, HENRIETTA STREET, COVENT GARDEN, LONDON.

THE STUDENTS' AIDS SERIES.

Specially designed to assist Students in grouping and committing to memory the subjects upon which they are to be examined.

'As aids, not substitutes, these little books afford the means of refreshing the memory, and of economizing time.'—*Edinburgh Med. Journal.*

Aids to the Analysis and Assay of Ores, Metals, Fuels, etc. By J. J. MORGAN, F.I.C., F.C.S. With 8 Illustrations. Price, cloth, 2s. 6d., paper 2s.

Aids to the Analysis of Food and Drugs. By C. G. MOOR, M.A., F.I.C., and W. PARTRIDGE, F.I.C. Third Edition. Price, cloth, 3s. 6d. net, paper 3s. net.

Aids to Anatomy (Pocket Anatomy). By C. H. FAGGE, M.B., M.S., F.R.C.S. Sixth Edition. (Thirty-Second Thousand.) Price 3s. 6d. net.

Aids to Bacteriology. By C. G. MOOR, M.A., F.I.C., and W. PARTRIDGE, F.I.C. Second Edition. Price 3s. 6d. net and 3s. net.

Aids to Chemistry. By T. A. HENRY, D.Sc. Eighth Edition. Price 4s. 6d. and 4s.

Aids to the Diagnosis and Treatment of the Diseases of Children. By J. McCaw, M.D., L.R.C.P. Edin Fourth Edition. Price 4s. net and 3s. 6d. net.

Aids to the Feeding and Hygiene of Infants and Children. By the same Author. Price 2s. 6d. and 2s.

Aids to Dental Anatomy and Physiology. By ARTHUR S. UNDERWOOD, M.R.C.S., L.D.S. Eng. Second Edition. Price 2s. 6d. and 2s.

Aids to Dental Surgery. By ARTHUR S. UNDERWOOD, M.R.C.S., L.D.S. Eng., and DOUGLAS GABELL, M.R.C.S., L.R.C.P. Lond., L.D.S. Eng. Second Edition. Price 2s. 6d. and 2s.

Aids to Medical Diagnosis. By A. J. WHITING, M.D., C.M. Edin., M.R.C.S. Lond. Price 2s. 6d. and 2s.

Aids to Microscopic Diagnosis (Bacterial and Parasitic Diseases). By E. BLAKE KNOX, B.A., M.D., B.Ch., D.P.H. Dub., Capt., R.A.M.C. Price 2s. 6d. net and 2s. net.

Aids to Surgical Diagnosis. By H. W. CARSON, F.R.C.S. Eng. Price 3s. 6d. and 3s.

Aids to Practical Dispensing. By C. J. S. THOMPSON. Price 2s. 6d. and 2s.

Aids to Examinations; being Questions and Answers on Materia Medica, Medicine, Midwifery, Pathology, and Forensic Medicine. By DAVID WALSH, M.B., M.Ch. Edin. Price 2s. 6d. & 2s.

Aids to Examinations. Part II. Medicine and the Allied Sciences. By T. REVELL ATKINSON, M.D. Durham. Price 2s. 6d. and 2s.

Aids to Examinations: Replies to Questions in Therapeutics. By J. BRINDLEY JAMES, M.R.C.S. Price 1s. 6d. and 1s.

Aids to Forensic Medicine and Toxicology. By WILLIAM MURRELL, M.D., F.R.C.P. Seventh Edition. Price 2s. 6d. net and 2s. net.

Aids to Analytical Geometry.

I.—THE STRAIGHT LINE AND CIRCLE. By A. LE SUEUR, B.A. Cantab. Second Edition. Price 2s.

II.—THE CONIC SECTIONS. By GEORGE HEPPEL, M.A. Cantab. Price 2s.

BAILLIÈRE, TINDALL & COX,

8, HENRIETTA STREET, COVENT GARDEN, LONDON.

THE STUDENTS' AIDS SERIES—*continued.*

Aids to Gynæcology. Fifth Edition. By S. JERVOIS AARONS, M.D. Edin., M.R.C.P. Lond. Price 2s. 6d. net and 2s. net.

Aids to Histology. By A. GOODALL, M.D., F.R.C.P. Edin. With 20 Illustrations. Price 2s. 6d. net and 2s. net.

Aids to the Mathematics of Hygiene. By R. BRUCE FERGUSON, M.A., M.D., B.C. Cantab., D.P.H., M.R.C.S. Fourth Edition. Price 2s. 6d. net and 2s. net.

Aids to Materia Medica and Therapeutics. By W. MURRELL, M.D., F.R.C.P.
Part I.—Introduction and Inorganic Materia Medica. 2s. 6d. and 2s.
Part II.—Drugs of Vegetable Origin. Price 2s. 6d. and 2s.
Part III.—Synthetical Products; Drugs of Animal Origin; Glandular and Serum Therapeutics. Price 2s. 6d. and 2s.

Aids to Medicine. By B. HUDSON, M.D. Cantab., M.R.C.P. Lond. Price 3s. net and 2s. 6d. net.

Aids to Obstetrics. By SAMUEL NALL, B.A., M.B. Cantab., M.R.C.P. Lond. Seventh Edition. Revised by C. J. NEPEAN LONGRIDGE, M.D. Vict., F.R.C.S. Eng., M.R.C.P. Lond. Price 2s. 6d. net and 2s. net.

Aids to Ophthalmology. By N. BISHOP HARMAN, M.A., M.B. Cantab., F.R.C.S. Eng. Fifth Edition. Price 2s. 6d. net and 2s. net.

Aids to Osteology (Pocket Osteology). By P. TURNER, M.B., B.Sc. Lond., F.R.C.S. Eng. Price 3s. net.

Aids to Pathology. By HARRY CAMPBELL, M.D. Lond., B.S., F.R.C.P. Second Edition. Price 3s. 6d. net and 3s. net.

Aids to Practical Pharmacy. By A. CAMPBELL STARK, M.B., L.S.A. Second Edition. Price 2s. 6d. net and 2s. net.

Aids to Physiology. By PEYTON BEALE, F.R.C.S. Eng. Price 3s. 6d. and 3s.

Aids to Practical Physiology. By J. BRINDLEY JAMES, M.R.C.S., L.R.C.P. Price 1s. 6d. and 1s.

Aids to Public Health. By D. SOMMERVILLE, M.A., M.D., D.P.H. Price 2s. 6d. net and 2s. net.

Aids to Sanitary Law. By HARRY CRITCHLEY, M.A., M.D., D.P.H. Price 2s. 6d. and 2s.

Aids to Sanitary Science. By FRANCIS J. ALLAN, M.D., and R. A. FARRAR, M.A., M.D., D.P.H. Second Edition. Price 4s. 6d. and 4s.

Aids to Surgery. By JOSEPH CUNNING, M.B., B.S., F.R.C.S. Eng. Third Edition. Price 4s. net and 3s. 6d. net.

Aids to Rational Therapeutics. By J. MILNER FOTHERGILL, M.D., M.R.C.P. Lond. Price 2s. 6d. and 2s.

Aids to Tropical Medicine. By GILBERT E. BROOKE, M.A. Cantab., D.P.H., F.R.G.S. Price 2s. 6d. net and 2s. net.

Aids to Tropical Hygiene. By R. J. BLACKHAM, D.P.H. Lond., Major R.A.M.C. Price 3s. net and 2s. 6d. net.

BAILLIÈRE, TINDALL & COX,
8, HENRIETTA STREET, COVENT GARDEN, LONDON.

MP
E.

~~169360~~ Rep. by

168921

Author French, Herbert

Title medical laboratory methods (and tests.)

University of Toronto
Library

DO NOT
REMOVE
THE
CARD
FROM
THIS
POCKET

Acme Library Card Pocket
Under Pat. "Ref. Index File"
Made by LIBRARY BUREAU

W. Lancaster